

1947
53
1

PHYTOLOGIA

An international journal to expedite botanical and phytoecological publication

Vol. 53

March 1983

No. 1

FIFTIETH JUBILEE YEAR

CONTENTS

REVEAL, J. L., *Significance of pre-1753 botanical explorations in temperate North America on Linnaeus' first edition of Species Plantarum* 1

LIBRARY

MAR 25 1983

NEW YORK
BOTANICAL GARDEN

Published by Harold N. Moldenke and Alma L. Moldenke

303 Parkside Road
Plainfield, New Jersey 07060
U.S.A.

Price of this number \$3.00; for this volume \$13.00 in advance or \$14.00 after close of the volume; \$5.00 extra to all foreign addresses and domestic dealers; 512 pages constitute a complete volume; claims for numbers lost in the mails must be made immediately after receipt of the next following number for free replacement; back volume prices apply if payment is received after a volume is closed.

**SIGNIFICANCE OF PRE-1753 BOTANICAL EXPLORATIONS IN
TEMPERATE NORTH AMERICA ON LINNAEUS' FIRST EDITION
OF SPECIES PLANTARUM**

James L. Reveal

Department of Botany, University of Maryland,
College Park, MD 20742, U.S.A.

INTRODUCTION

When Carl Linnaeus published the first edition of *Species plantarum* in 1753 he discussed some 889 species of vascular plants from temperate North America, here defined as the present-day continental United States and Canada. The information relative to these plants came from the efforts of men and women who either visited the New World or who lived there and collected plants as amateur or professional naturalists.

In a few instances they were gentlemen correspondents of Linnaeus' or were men in Europe who never traveled but supplied Linnaeus with collections sent by others. A few were chance collectors, some gathered plants as a hobby, and others to gain political favors. Occasionally a collector was a person trained in medicine and used that knowledge to search for promising plants that might be used in the treatment of maladies. More often collectors were members of the clergy who collected potentially important horticultural plants for their gentlemen supporters in Europe. This led not only to the discovery of many important ornamental trees and shrubs, but many New World plants of agricultural import-

ance. The early Spanish explorators discovered a rich array of grains, fruits and vegetables that eventually found use in European culture and cuisine. The more notable were maize, potato, tomato, various squashes and of course a non-food plant, tobacco.

The early explorers who ventured into the New World were mainly interested in the riches of gold and silver. Even through natural history was of relatively minor concern, surprising numbers of books were published treating the native flora and fauna of the newly discovered lands. In 1526, Oviedo y Valdes published a book on the natural history of the West Indies based on his own observations in the Caribbean and Central America. Nicolas Monardes who never left Europe, published his book, in parts, on the natural history of the New World from 1569 to 1574. Both men were knowledgeable physicians and as such were particularly interested in medicinal plants and their local uses. Monardes' book was rewritten and published in Latin by Carolus Clusius in 1574. John Frampton translated that work into English and published it in 1577.

The exotic vegetation of the new lands across the ocean

The present paper was the subject of a presidential address to the Botanical Society of Washington, December 1982. The research reported here was supported by National Science Foundation grant DEB80-04628, the United States Fish and Wildlife Service's Office of Endangered Species, and the University of Maryland. This is Scientific Paper No. A3369, Contribution No. 6441 of the Maryland Agricultural Experiment Station.

proved exciting to the European naturalists, and a source of considerable confusion. Species somewhat similar to those in Europe which proved to be useful were often considered to be the same species, even if they were not, so as to encourage their use in the medical trade. Plants which were the same, but were of no use or interest to the European physicians, were either ignored or taken to be a different species. It is well to remember that in the late 1500s, the concept of species was essentially that proposed by the ancient Greeks, chiefly Pliny, and the unique New World species were proving difficult to fit into Pliny's scheme of the vegetable kingdom.

Plants which were of potential economic importance were sometimes ignored or not used by the Europeans. Maize was not used extensively in Europe as a table food, and is even today largely considered an animal fodder. Potato was not considered fit for the tables of the aristocratic, and was shunted off to the slaves of the New World or the poor of Europe. For some economically important plants, tradition indicated they were poisonous such as was the case with the tomato up to the end of the last century. Some plants, because they were used by the "heathens", were considered unchristian for if God had intended christians to have such foods, they would be found in Europe!

Jose d'Acosta, a missionary, lived in Peru from 1569 to 1588. His notes originally were published in 1589 and then expanded into a four-volume work in 1590. These volumes were translated into French, Flemish and German around 1600, and then into English in 1604. His exact

observations are still a major source of information on the uses of plants by the native people of South America. d'Acosta, like Ovieda and Monardes, talked of maize, cassava and potato among other species. He also noted the heavy use of chili peppers, and reported that the Peruvian Indians were exceedingly fond of garlic which had been introduced into the New World by the Europeans. He called attention to the fine grapes in the New World, and introduced several kinds into Europe.

Collectors to temperate North America came only a few years after those that visited the more tropical regions. Early French explorers gathered plants and sent seeds and fruits back to France where they were soon flourishing in the gardens. Many of these species were gathered in what is today southeastern Canada. There was a desire on the part of the French to introduce plants not only of medicinal but of potential horticultural value as well. Many temperate conifers and hardwoods were ideally suited to the French climate and quickly were adopted, unlike the more tropical species.

By the start of the 17th century interest in plants was beginning to slowly move away from the rigidity of the Dark Ages and into a more enlightened period. The nearly 500 species of plants described by Dioscorides in his work on *materia medica* were destined to give way to over 6000 species in the works of the Bauhin brothers, and especially Gaspard Bauhin. They described several new species from the New World in 1623. The great philosophical barrier of Old World versus New World speculation had finally been broken, and with that restriction lifted, the urge to

search and discover new plants from all corners of the world began to be felt by botanists and explorers.

As the 17th century began some 50 species of North American plants were known from cultivation in European gardens. The concept of the botanical garden, begun in Venice in 1545, was rapidly becoming the fashion, and soon no medical school was complete without a rich and varied garden of medicinal plants. Botanical gardens were established in England and France by the start of the 1600s, and with them came the beginnings of a golden age of botanical exploration throughout the world.

THE GARDEN CURIOSITIES

During the first half of the 17th century, botanical explorations in temperate North America were restricted to individuals who, from time to time, gathered seeds. Little concern was given to the preservation of dried voucher specimens, and almost none to recording the details of geographical location, plant association or similar information. John Tradescant the Younger came to Virginia in 1637 and collected plants along the York River. The purpose of his trip was to gather novelties for his natural history museum, known as Tradescant's Ark, which he and his father had established at Lambeth in London. Tradescant brought to England such species as the tulip tree, *Liriodendron tulipifera*, the red maple, *Acer rubrum*, sycamore, *Platanus occidentalis*, black walnut, *Juglans nigra*, and the bald cypress, *Taxodium distichum*. Although herbarium specimens of these species were not made in Virginia by Tradescant, various collectors in later years made voucher spec-

imens of John Tradescant's original introductions. These specimens are still extant and may be examined in the Sloane Herbarium at the British Museum (Natural History) in London. The museum specimens shown in Tradescant's Ark were eventually obtained by Elias Ashmole, and they became a part of the Ashmolean Museum, and are now a part of the University of Oxford Museum at Oxford, England.

To be sure the early discoveries of temperate North American plants were finding their way into the literature. As already mentioned, Gaspard Bauhin reported several in 1623. However, many were also described, illustrated and evaluated by herbalists. Maize and other New World plants were well known to European naturalists early in the 16th century, and a number of herbals contain illustrations of these species. The tradition of herbals was slowly being replaced by works of botany more concerned with the classification and identification of species, and less so with the medicinal uses of plants. Nonetheless, John Parkinson, the last of the English herbalist, was destined to describe several American species in his 1640 herbal, *Theatricum botanicum*. Plants reportedly found originally in Canada, New England and Virginia were included in Parkinson's book.

As for the Canadian references, Parkinson took most of his information from Jacques Philippe Cornuti's *Canadensium plantarum* published in 1635. This book described species of plants found in cultivation in and around Paris. A total of 68 species were described, but only 38 were from Canada or the New England area. Cornuti's book was the most scientific effort yet produced devoted to New

World botany, and nearly all of the 38 copper plate illustrations of Canadian plants executed by Cornuti were ultimately cited by Linnaeus when he wrote the first edition of *Species plantarum* in 1753. Linnaeus referred to Parkinson's book also, but much less frequently than Cornuti's.

Before leaving this period Thomas Hariot, friend and mentor of Sir Walter Raleigh, must be mentioned. Hariot came to Virginia in the 1580s and described what he found in a famous pamphlet published in 1588. Entitled "A briefe and true report of the new found land of Virginia", his remarks are valuable because of the detail he devotes to the forest resources and medicinal plants. He describes the native uses of grapes, beans, potatos, melons, sunflowers, tobacco, and a wide variety of small grains including oats, barley and what he called wheat. Hariot reported on the several different kinds of acorns used by the Indians as well as walnuts, strawberries, mulberries and crab apples. Among the identifiable trees were species of oak, elm, ash, walnut, fir, juniper, maple, witch hazel, willow, beech, sassafras and pines. Hariot's report on Virginia -- which was also based on observations made in North Carolina -- was an exceedingly glowing one, and while not consulted by Linnaeus, it is of interest today for its early information on plant ecology and ethnology.

There were many other similar publications on the general aspects of the flora of temperate North America published by travelers prior to the 1680s. Most were in the vein of Hariot's. To Linnaeus they were of little or no interest. Even to contemporary naturalists they

were little more than descriptive catalogues of vague but wonderous species of plants which may or may not have much of a factual basis in reality. In many cases such publications were "flowery" so as to attract colonists to the New World. To the English, however, there was no question that great botanical treasures awaited discovery in the New World and that skilled and knowledgeable men were needed to survive the rigors of the new lands. The rapid growth of scientific organizations and thought in England and France was fomenting desires to explore, discover, and to answer questions about nature.

After the establishment of the Oxford Club in 1648, other scientific organizations were soon formed. Upon the final restoration of the Stuart monarchy in 1660, the Royal Society of London was established. This was, at least by modern standards, the first truly scientific organization in England. The first twenty years of the organization was a period of maturing and promoting scientific endeavors in various parts of the world, including the subtropical and tropical regions of the New World.

The growth of botanical gardens in Europe was significant during the last half of the 17th century. What initially had begun only as a desire of the nobility to possess exotic plants and animals from strange new lands soon escalated beyond the royal estates and gardens into the homes and estates of not only the more wealthy families of Europe, but even to the gardens of the common people. The exploration of the new lands across the sea, in Africa, the New World, and the Far East was creating new wealth. These new men of means were often

well educated, skilled in the arts, and were impressed by exotica acquired by of their fellows. They were also willing to associate with persons of ability, be they in art or literature, or persons trained as natural philosophers interested in the classification and identification of vascular plants. With the growth of botanical gardens, the craze for exotic garden plants increased, and friendships with knowledgeable botanists soon became paramount for nearly every gentlemen of means who wished to have a well-curated cabinet of natural objects. The classifiers and identifiers of plants suddenly became highly esteemed in the gentlemen's circle of close friends.

The Royal Society of London provided a common meeting place for scholars, noblemen and men of means. The union was perfect for land was needed for gardens, money and connections were needed to provide the means to obtain plants. What better a relationship to foster than that between the hopeful botanist, the newly established exporter of foreign curiosities, and the landed gentry who wished to have the exotics in their garden?

With the beginning of efforts to prepare a detailed review of the plants of the world, especially by English workers such as John Ray and Robert Morison, to say nothing of other workers in Europe, the need for more collectors in temperate North America grew. More and more members of the Royal Society began to call for an expedition to North America. Finally, it was decided to fund an expedition and a search was begun to find a suitable naturalist. The results produced John Banister.

THE VIRGINIA BEGINNINGS

John Banister was born in 1650 and educated at Magdalen College at the University of Oxford where he received his B.A. in 1671 and his M.A. in 1674. He continued to serve the University after graduation, first as a clerk and then as a chaplain. At Oxford he met Dr. Robert Morison, the University's botany professor, and came to know the Bishop of London, Dr. Henry Compton, a noted gardener and botanical enthusiast. Under the guidance of these two men, Banister began to botanize and make collections of dried plants in the Oxford area. His skills soon manifested themselves, and Compton encouraged Banister to accept a ministerial position in Virginia. His duties included collecting objects of natural history while serving the needs of the Church of England.

Banister accepted the position and left for Virginia in 1678 under the sponsorship of a number of members of the Royal Society. William Byrd I of Virginia, served as treasurer of this group.

At first Banister stayed with Byrd at James River Falls and then at Westover, the Byrd estate. Not only did Banister send back to England a large number of seeds, fruits, dried specimens, and illustrations of native plants, he also received plants from England and plant them in Virginia. The European plants were coming from Jacob Bobart, the gardener at the University of Oxford, who, altogether with his father, served the University for over 60 years.

Financial support for Banister's efforts was never really substantial. His English patrons included such men as Dr. Hans Sloane, Dr. Henry

Compton, Dr. Martin Lister, Robert Morison, Jacob Bobart, George London, James Petiver and others, but none seemed capable of providing him with sufficient funding so that he could devote his full efforts to botanical explorations in Virginia.

In the 1680s, Banister entered into a lively correspondence with John Ray, and sent to him a large number of plant specimens. At the same time Banister was also providing specimens to a second major English botanist, Leonard Plukenet, Ray's rival. Ray took a catalogue of Virginia plants sent to him by Banister and published it in 1688 as an appendix to his second volume of *Historia plantarum*. Plukenet, on the other hand, took his Banister specimens, and in particular the Banister illustrations, and began to publish the figures in 1691, and a series of new scientific names (polynomials) in his first book, *Almagestum botanicum*, published in 1696.

To a degree Banister was depressed by these publishing activities. He himself began preparing manuscripts with the idea of publishing his findings under his own name. His previous attempts to publish his own works had failed for whenever he sent a manuscript to Europe, it was taken and published by others under their own name. Morison, Ray and Plukenet were all guilty of this, from time to time, but without their efforts to present the Virginian discoveries at once, others, especially workers in France and Holland, would have described the species anyway when they observed the uniqueness of the plants in their garden.

In spite of John Banister's difficulties, his fortunes changed somewhat when he

married a wealthy widow, and suddenly found a ready source of modest support. In addition, his station in Virginia was changing, and while still a minister, he accepted other duties and these often kept him out-of-doors collecting plants. Soon, the number of Virginia plants Banister sent to England surpassed 300 species, and was destined to reach about 340 species of vascular plants, 100 insects and about 20 mollusks before a rifle ball ended his life in an accidental shooting while he was out botanizing. The year was 1692, and his passing would close the opening period of temperate North American botany.

To Morison, Bobart, Ray and Plukenet, the collections of John Banister were the mainstay of their knowledge of temperate North American botany. Bobart took up the writing of Morison's *Plantarum historiae universalis oxoniensis* when Morison died in 1683, and began to describe and illustrate Banister species based on a combination of garden-grown material raised at Oxford and herbarium specimens. Banister's catalogue, which was published by Ray in 1688, was not illustrated, but Plukenet's books were illustrated. Unlike Bobart, however, Plukenet based his illustrations almost entirely upon herbarium specimens or figures executed by Banister which Plukenet obtained. Between these men, and others, notable Joseph Pitton de Tournefort in France and Paul Hermann of Leyden, Banister's plants were rapidly described and characterized.

Of perhaps even greater importance were the many species of vascular plants Banister introduced into western Europe. He left a legacy of exotic trees, shrubs and herbs

which generations of naturalists in the future would examine in even greater detail. Linnaeus would not only see and grow many of the Virginia species introduced into Europe by Banister, but would come to acknowledge Banister as an important collector of temperate North American species.

THE BOTANY CLUB

The death of John Banister was a serious blow to English botany and its attempt to classify temperate North American plants and to obtain seeds and fruits to introduce such species into cultivation. The loss of Banister produced a significant gap in the active program of plant explorations in the British colonies of North America, and it was immediately agreed that a new person had to be found to replace him.

The Royal Society had been growing, expanding and changing over the three decades from its foundation in 1660 to 1690. A new generation of men were coming into power within the Society, and even within English society as well. Among the leaders of this new group was Hans Sloane. Sloane was a physician who had gone to the New World with the new Royal Governor of Jamaica, the Duke of Albemarle. He left England in September of 1687 and returned in late May of 1689 laden with several hundred new plants. Within a short period of time, Sloane increased significantly his knowledge of botany by studying with Tournefort and Magnol in France, and by working on his own collections. Sloane soon obtained a number of Royal appointments including physician to Her Majesty the Queen, and established a large and profitable practice. Sloane married a wealthy widow, whom he likely met in Jamaica,

and was soon in a position to devote much of his time to the gathering of curiosities from throughout the world. Sloane would also rise to power in the Royal Society, holding the important post of treasurer which saw to the publication of the *Philosophical Transactions*, the scientific journal of the Society. It was due to Sloane that the journal saw a significant rise in the number of papers dealing with botany. All in all, Sloane would live for 93 years, dying in 1753. In the years of his activities in the natural sciences, Sir Hans Sloane would obtain the largest collection of objects in the world, and because he lived so long, was able to obtain -- and thus save for future generations -- the volumes of herbarium specimens made by a majority of his contemporaries.

Others who rose to prominence during the 1690s were men who can be roughly divided into two groups: scientists and the supporters of scientists. The distinction is not absolute, as the life of Sir Hans Sloane witnesses. Foremost of the scientists was John Ray. His efforts during the 1680s to produce a two-volume work on the classification of all the plants of the world was, and still is, considered the finest produced of this period. Ray is best known for his flora of England which was the standard that Linnaeus had to overcome a half century later to win the critical support Linnaeus needed in England to carry his ideas of plant classification and nomenclature. Even so, the arrangement of plants proposed by Ray was still preferred by most English workers over Linnaeus' so-called "sexual" system. Today, many consider Ray to be the "Father of systematic botany". Close to Ray was Samuel Doody, Keeper

of the Chelsea Physic Garden, the medicinal garden of the Society of Apothecaries. A neighbor was Samuel Dale who would be an important intermediary between Ray and other English naturalists, and the men and women who collected plants in North America. Dale would amass a large collection of dried plants, and it would be to Dale that the Ray herbarium and library would go when Ray died. Ray was modestly but consistently supported by a number of gentlemen who were interested in botany. Most important was Charles Hatton, an aristocratic promoter, who funded many of Ray's publications.

In the opposing botanical camp were the followers of Leonard Plukenet. Plukenet, like Ray, was a professional botanist, and equally skilled in the art of systematic botany. Unlike Ray who was a man of ill health and abject poverty, Plukenet was at least healthy. His temperament was less conducive to good will, and the Society was badly divided into factions based on the supporters of Ray versus Plukenet. While Ray had a broad base of support, Plukenet's was largely restricted to the even more ill-tempered John Woodward. Woodward, who was even less well thought of by his contemporaries, was not as skilled as a naturalist as Plukenet, and he tended to be ignored by most of the membership of the Royal Society.

Some of the men of note involved in the promotion of botanical science in the Royal Society were James Ayrey, a Quaker merchant; Charles Du Bois, secretary of the East India Company; and Dr. Henry Compton. Compton is a special case. He was a major supporter of the Society's efforts to obtain plants, but as an avid gardener and the Bishop of

London, he was ideally positioned to make certain convenient appointments which benefited both systematic botany and his personal garden. In short, it was no accident that many naturalists that were sent into the field by the Royal Society were ministers for the Church of England.

Bordering both groups were two men. George London was initially the gardener for Henry Compton, but from the late 1690s to the early 1700s, he was gardener to William and Mary, and after her death, to William. As the Royal Gardener, London was ideally suited to ensure a proper place for the exotic plants coming into England from foreign lands. A skilled naturalist in his own right, London did much to foster the international exchange of seeds and bulbs, and to maintain some degree of cooperation between the various factions within the systematic community in London.

A second person is of considerable importance, especially in the history of the discovery of vascular plants in temperate North America. James Petiver was born in about 1663, and established an apothecary shop in London out of which he maintained a world wide correspondence with collectors of natural objects. Interested mainly in insects and shells, Petiver relied mainly upon ship's captains and surgeons, soldiers, naval officers, local physicians, assorted farmers, and even the bored wives of many gentlemen to collect curious plants and animals for him, and send them to London so that he might describe them in his own publications.

Petiver was not only a major promoter of the natural sciences to the amateur, he

often saw to it that their discoveries were rapidly published. Unlike most of his contemporaries, Petiver was willing to provide his correspondents with practical information including sets of instructions on how to collect and preserve dried specimens, ship seeds and seedlings, and to record useful information to go with the collections. As an author, Petiver was respected. He was not of the stature of Ray, or even Plukenet, for his observations were often with little regards to what had already been published and he had a tendency to redescribe species others had already proposed. His pamphlets and articles are often filled with valuable historical information. As Petiver was receiving so much material from causal collectors, it is only by reading his publications that one can find who was collecting where at any one time in history.

All of these men were members of the Royal Society, but as a whole, the members of the Society were not devote to the subject of botany to the degree satisfactory to those who wished to concentrate on the plant kingdom. The stature of the Society was such that causal discussions were often difficult, and the mere matters of the day-to-day discussions of botanical nomenclature, the latest book on the mosses of England, or the problems of shipping seeds from China were not the type of heady matters one ought to present before the entire membership of the Society. Thus, those gentlemen interested in botany met informally at what must have been one of the first botanical societies in the world. It was known as the Temple Coffee House Botany Club.

Members of the Temple

Coffee House Botany Club met every Friday evening. There are no minutes of the meetings, for it was an informal gathering, but contemporary correspondence indicates that the membership consisted of such men as Sloane, London, Compton, Petiver, Doody, Dale, Ray, Plukenet, Woodward, and when in London, William Sherard and the younger Jacob Bobart of Oxford. From time to time field trips were taken, usually on Sundays, to gardens or places of botanical interest. In the relaxed comforts of the Temple Coffee House, the more detailed aspects of botany could be discussed and it was not unusual for members to bring recently acquired collections of exotic species for demonstration before the membership or to ask for help in establishing the correct scientific name.

It was here, no doubt, that the death of John Banister was discussed, and the need for a replacement reviewed. Compton likely proposed that he, as the Bishop of London and a proponent of the expansion of the Church of England in the New World, could find a position within a parish if the Society could find a naturalist willing to take vows. Funding support was likely discussed as well, and a group of supporters would be established and a treasurer appointed. It would be a duty of the treasure to secure a governmental post so that the colonial government would support the naturalist when he went about his collecting activities. Once the details were worked out, and the plans finalized, then the matter was brought before the membership of the Royal Society for approval. The system worked well.

It was William Byrd I who urged the Royal Society to

send a new collector to Virginia, and called for a search. This was initiated in 1693, but no one was found that was willing or qualified to fill the position. Funding was finally found in the form of a position from the Royal Governor of Maryland, Francis Nicholson, who had previously been Vice-Governor in Virginia. A strong supporter of the expansion of the Church of England, Nicholson's duties in the Catholic dominated colony of Maryland were to transform that colony to the Anglican faith. He moved the capitol from St. Marys to Annapolis, and saw, in the needs of the Royal Society, a means of adding another clergyman.

In late 1694, Edward Lhwyd, Keeper of the Ashmolean Museum at Oxford, wrote to Compton that he had a young man that might be qualified. Lhwyd indicated that he could train the young man in natural history, but it was up to Compton to provide training in the ways of the Church. The young man he spoke of was his deputy keeper, Hugh Jones.

BOTANY IN MARYLAND

Hugh Jones came to Oxford in 1694 on a pauper's scholarship. Born in Wales, he was probably only 24 when Lhwyd reported to Compton that he might be qualified. Over the course of 1695, Lhwyd provided Jones with a basic education in the natural sciences, and in December of that year was able to send him to London armed with glowing letters of introduction and instructions to take his religious vows. Over the course of the month, Jones was schooled on the matters of the Church of England, and in the end was ordained a priest and became a deacon. He was now ready to go to the New World and begin a new life in Maryland.

The Byrd resolution proposed and accepted by the Royal Society was to send a naturalist to Virginia. Funding, however, could not be found in that colony but was obtained from Maryland's governor. When Jones completed his training at the end of December, 1695, he set out for the coast from London to board a ship for the Chesapeake Bay, and the new capitol of Maryland, Annapolis. Rough weather made it impossible for the ship to depart for several weeks, and from time to time, Jones would travel back to London. He could not afford to stay long in the city, but apparently did visit the Temple Coffee House Botany Club where he fell into the Sloane circle of influence. It was James Petiver who took command of the Reverend Jones and began to educate him in the fine art of practical botany. Through him and other members of the Botany Club, Jones received instructions, various supplies, and an occasional piece of equipment. Most importantly were the personal contacts he made. From Petiver, Jones received a request that he should send to him insects and fossils; for Ayrey and Doody he should gather seeds and fruits; and for London and others he should make herbarium specimens. Petiver further urged that Jones should send everything to him for dispersal. He, in turn, would provide Jones with new publications in the field of natural history, help with his medical needs, and provide news about London.

When Jones' ship finally departed in the spring of 1696, Jones was as educated as could be expected. He eventually landed in Maryland, arriving in Calvert County in the heat of August and rode horseback to Annapolis -- a city of less than 40 buildings

and considerable mud -- where he resided with Governor Nicholson for five weeks.

The initial proposal was for Jones to serve as Nicholson's chaplain, but even before Jones left for Maryland, this proposal had been altered to allow Jones to assume duties in one of Maryland's parishes. After five weeks in Annapolis, Jones left for Christ Church Parish in southern Calvert County where he assumed the rectorship of one of Maryland's largest and most wealthy parishes.

During the summer and fall of 1696, Jones collected what he could, and continued to look for objects of interest for his London friends over the winter. In March of 1697, Jones sent letters and two boxes of specimens to Petiver in London. During the growing season of 1697, Jones searched among the meadows and woods near his church, along the edges of the Patuxent River, and the cliffs along the shore of the Chesapeake. He found a wide variety of herbs, shrubs and trees, assorted insects and butterflies, and fossils. Soon he had specimens of birds, small mammals and minerals to go along with the seeds and young plants we was sending to London.

What, no doubt, Jones sent in great expectation, and awaited in the same fashion, was not received with the same level of enthusiasm previously experienced by Petiver and others when a shipment from Banister arrived in the Old World. The seedlings were ill-packed and did not survive the sea voyage. The seeds were poorly labeled, the shells unsorted, the plant specimens small and fragmentary, and the insects broken or damaged. The animals were

poorly preserved and few could be adequately identified or even characterized. For the members of the Temple Coffee House Botany Club, the Reverend Hugh Jones was a profound disappointment.

For Jones, Maryland was not all one could wish for either. The winter of 1696-1697 was especially hard. He writes that snow came in November and remained until March, and at times the snow exceed two feet in depth. The Chesapeake Bay became frozen with ice so thick that sailors could walk to the shore from their ice-bound ships. The cold rains were hard on Jones and members of his parish. Still, for the young minister, his ministry was a challenge.

For Jones, as with most members of the Church of England in Maryland, the long history of Catholic rule represented the past and it was his duty to overcome the errors of religious views previously imposed upon the people. That Maryland had been settled as a Catholic colony was of no immediate importance to the Crown, the Royal Governor, or even the Reverend Hugh Jones. His directions were to conduct all services according to the dictates of the Church of England, no matter the local religious preferences in the community. Similarly, the obvious threat imposed by the growing number of Quakers coming southward from Pennsylvania was a matter of concern, and Jones was a signatory to a document demanding governmental control over the teachings of the Quakers. As a minister in Maryland's most wealthy parishes, Jones was well placed in the social fabric of the Colony. His yearly income was such that he was a wealthy man, that coming from a tax, in tobacco, levied by the sheriff for the minister. The

levy was based on the number of people in the parish, with each adult white man or woman equal to one share per person while any black slave, no matter the age or sex, was equal to two-thirds of a share. Good year or bad, the minister received his levy.

For Petiver the failings of Hugh Jones in Maryland to carry out his mandate to collect was particularly troublesome. As his strongest supporter in London, the plight of trying to collect, and survive in the New World, were understood by Petiver. Petiver stood to lose his own investment of time and energy if Jones failed, but it was clear even to the apothecary that the hopes for a significant improvement in Jones' abilities were ill-founded. Petiver himself expressed his disappointment when he reviewed his broken insects, yet he attempted to describe what he could. The bulk of Jones' Maryland plants went directly to James Ayrey and George London, and while Petiver received a few, these were not a significant part of his botanical holdings. However, in late 1697, George London gave to Petiver two large volumes of dried plants gathered in the New World, including a significant set of Jones' plants. Armed with these two volumes filled with Maryland plants, Petiver set at once to characterize them and to prepare an article for publication in the *Philosophical Transactions*.

To further this effort, Petiver wrote to Jones urging him to send all collections that he could immediately. In addition he told Jones of the disappointment his collection had caused in London and informed him of a move to send others to the Colony. If Jones was to receive credit

for his work in Maryland, Petiver warned, he had to send his plant and animal collections before the end of 1698, for in that year, others would be collecting in Maryland.

At the November, 1697, meeting of the Royal Society, William Byrd II moved that the Society should find "a Fitt person" to be sent to "Virginia in order to make observations and Descriptions of all the Naturall productions of those parts and to write the History thereof." Although Byrd urged that the naturalist be sent to Virginia, the remaining part of his motion stated that for "such a fitt person the charge of his passage and 25 pounds per Ann. would be allowed him by the Governor of Maryland." Once again, Francis Nicholson offered to fund the collector, and the naturalist was to be in Maryland, not Virginia.

Nearly all of the necessary arrangements were completed at the Temple Coffee House Botany Club before the vote and William Vernon, a fellow of Peterhouse at Cambridge University, was determined to be that "fitt" person called for in Byrd's resolution. He was approved at the December meeting of the Royal Society, and set out for Maryland in January, 1698.

Vernon arrived in the Maryland in April and he began to collect almost immediately. Vernon's proposal was to remain in Maryland for three years, and to concentrate upon the bryophytes, lichens and similar non-vascular plants. He had received permission from the University to be gone from Cambridge with the proviso that he study botany, not marry, and report each year that he was alive!

The warnings sent to

Jones were heeded, and a large number of plant collections arrived in London in early 1698. Immediately Petiver set to work upon them, in concert with the Geogre London collections already in hand, but by this time Petiver was somewhat less concerned about his interests in Maryland for he had played a trump card in the form of Dr. David Krieg.

David Krieg was a Prussian friend of James Petiver's who had been living with him for much of 1697. Krieg had been illustrating many of Petiver's scientific papers during the year he was staying in London, and as a physican and skilled naturalists, Krieg was an ideal person to venture into the field. Petiver had been a party to Byrd's proposal to send Vernon to Maryland, but the opportunity to send another collector, especially one devoted to him, could not be passed by. During the conversations relative to Jones, Petiver must have convinced Krieg that he should go to Maryland as well. Krieg could not obtain support from the Royal Society, or even Francis Nicholson, so it was necessary for him to find a position as a ship's surgeon. This accomplished, Krieg departed in March of 1698, nearly two months after Vernon, and arrived in April just before Vernon.

Krieg's close and personal friendship with Petiver is well known, and their letters of affection continued well after their year together at Petiver's home in 1697. Perhaps as a part of that relationship, Petiver gave to Krieg a young man named Issac who was to assist Krieg with his explorations. Upon his departure for the New World, Petiver presented Issac with a rather famous set of instructions:

When ever you goe ashore take with you a Quire of Brown Paper or Collection Book. An Insect Box, Pins & a small Viall halfe fil'd with Spirit in which you drown all your supernumery Flies, Beetles, Catterpillars, & other Insects especially such you find in water. Also a Booke for Butterflies & Moths of each wch get al you can find, with a paper bag or two to put all ripe seed, ffruit & berries as also all ye shells you meet with both land & water & as many of each sort as you can find; such as are thin & brittle you must put into a Pocket by themselves with moss or any soft leaves to keep them from breaking.

These instructions were exact and perhaps a bit overbearing, especially if Issac had actually carried them out fully to the letter. As fate would have it, Issac was never allowed to leave the ship, and thus was of no value to Krieg as an assistant.

Petiver was not the only one trying to get a personal collector into Maryland. John Woodward -- the vowed enemy of Sloane and Petiver -- and his botanical associate, Plukenet, also wanted material from Maryland. At first, apparently, Woodward and Plukenet did not receive Jones material, and were not slated to receive Vernon collections either. Fearful of this, John Woodward apparently wrote to the Royal Governor, Francis Nicholson, and complain of Sloane and others in London. He implied that there were collectors in Maryland who were taking objects of nature without his permission, and that such person -- Jones, Krieg and Vernon -- were associated with such

"Rogues and Rascalls" as Sloane, Ray and Petiver. This correspondence must not have greatly impressed Nicholson for he knew of Woodward from others. Still, Petiver was concerned and in his letters urged Jones to send Woodward specimens which, apparently, Jones did. Vernon wrote Sloane from Maryland that although Woodward had claimed over 100 correspondents in America, he had met none and suspected Woodward's honesty. Vernon ended his commentary on Woodward by calling him "an abominable Villanous & Silly fellow".

The Reverend Hugh Jones knew that Krieg and Vernon were coming, and both men carried letters of introduction from Petiver. It is likely that all three men met each other in Maryland. It is certain that Krieg and Vernon collected jointly, and likely traveled together on a few occasions.

The collecting activities of Krieg and Vernon were limited to the coastal plain of Maryland, and an examination of their extant collections clearly shows that Krieg was able to collect plants more early in the year than Vernon. This was of little initial concern to Vernon as he was planning to stay in the Colony for three years. In July of 1698, Vernon wrote to Sloane that he was having to return to London. This was most likely because he had learned that Francis Nicholson had been appointed Royal Governor to Virginia, and would shortly leave Maryland. Without Governor Nicholson to support his position, Vernon would have to depart. Krieg, who planned to spend only the growing season in Maryland, collected more rapidly, gathering plants, insects, birds and mammals, fossils, shells and assorted

other items of natural curiosity. In October of 1698, Krieg and Vernon departed the Royal Colony of Maryland and arrived together in England less than two months later.

Almost immediately the collections of Krieg and Vernon were divided into sets and sent out to the supporters of Vernon, with duplicates of the Krieg collections going to many of the same people as well. Sloane received a large set of these collections, and these, in turn, were sent to John Ray for naming. Ray was in the process of completing a supplement to his *Historia plantarum*, and the new Maryland collections would prove a valuable addition to the world's flora. Krieg specimens went to Plukenet, probably through Petiver, and Petiver provided Sloane with additional material which was not sent to Ray. Duplicates were accepted by Ayrey from both Krieg and Vernon, and Sloane received a set of Vernon specimens directly from the collector. Some material was sent to William Sherard at Oxford. Seeds and fruits were distributed among the various growers, and soon Maryland plants were flourishing in English gardens.

BOOKS AND ARTICLES

The first of the major papers dealing with the natural history of Maryland was published by James Petiver. It appeared as an article in the last 1698 issue of *Philosophical Transactions* and was a catalogue of the plants and animals found in Maryland by Hugh Jones. In all 54 vascular plants were reported by Petiver. The majority were species previously known to naturalists and had been found in Virginia by Banister and other naturalists. A few were considered to be new species

by Petiver, and described as such. None of the species was illustrated.

A small collection of plants had been gathered in Maryland prior to the arrival of Hugh Jones in 1696. A few species from the Colony were described by Plukenet in 1696, and one was illustrated as early as 1691 in *Phytographia*, his book of illustrations. It is unclear who might have gathered these plants and when. A possible candidate is a ship's surgeon and later correspondent of James Petiver, Dr. John Smart. He gathered plants in Maryland in 1708. He is best known for his plants collection from the Hudson Bay region of Canada which he obtained in 1708. These specimens are found in the Petiver and Plukenet volumes of dried plants in the Sloane Herbarium.

In 1700 Plukenet published a second volume devoted to botany. His 1696 book, *Almagestum botanicum*, had been well received but was criticized as incomplete as a world flora. This was indeed true, and a supplement, *Almagesti botanici mantissa* was released in 1700. The *Mantissa* contained the description of over 200 species of Maryland plants, and several were illustrated in the *Phytographia* section included with the *Mantissa*. Most of the Maryland plants reported by Plukenet were those gathered by Jones and Krieg. A new book, *Amalthemum botanicum*, was published in 1705 by Plukenet. In that year only 16 Maryland species were described, but numerous plants previously described from the Colony were illustrated in 1705.

John Ray published a supplement to his two-volume *Historie plantarum* in 1704. Called the *Supplementum*, Ray ac-

counted for over 400 polynomials applied to Maryland plants. In addition to new species proposed by Ray, he listed, sometimes in synonymy, sometimes with comments, all previously published names which had been applied to Maryland plants by Petiver and Plukenet. William Sherard, and perhaps Jacob Bobart, described a small number of Maryland plants which Ray published in his *Supplementum*. Ray also published an index to James Petiver's herbarium in the *Supplementum*, and this contained descriptions of a few new species from Maryland.

From 1699 to 1702, Petiver added other new species of Maryland plants to the literature. These appeared in a series of pamphlets published by him, and many of the species were illustrated.

In addition to the Maryland plants reported in the published works of Plukenet, Petiver and Ray, these authors published new species from other regions of temperate North America as well. The majority came from Virginia, but a growing number of new collections were coming to Europe from the Carolinas and from the New England area. Unlike Banister and Vernon, who were professional naturalists in the sense that their positions were supported by the Royal Society, collections from other areas were being gathered by amateurs. Most of these people were ship's captains or surgeon, local residents, or the occasional visitor. None was destined to become a major supplier of plants to European gardens, however, and their impact upon the history of systematic botany is exceedingly limited.

Botanical explorations essentially ended in Maryland with the departure of Krieg

and Vernon in October of 1698. Jones remained in Maryland and continued to serve his parish. He gathered specimens in 1699, but by 1700 the effects of tuberculosis were beginning to sap his strength. His letters became less frequent and were more often filled with indications of his declining health. Finally, in January of 1702, Hugh Jones, minister and naturalist died. He was probably 31 years of age.

The failure of Jones, Krieg and Vernon in Maryland was a serious blow to the men of the Royal Society, and especially the members of the Temple Coffee House Botany Club. To be sure others were collecting along the eastern seaboard of North America, and James Petiver continued to receive specimens from a large number of persons from New England to Georgia. Several people were in Virginia and the Carolinas collecting specimens which Petiver would describe. Yet, death was taking its toll even of the membership of the Botany Club. Ray died in 1705 and Plukenet followed in 1706. Petiver himself would pass away in 1718, but by this time, Vernon and Krieg had both died.

Botany itself was changing. The desire to have exotic plants had not lessened in any fashion, but new and more exciting species were coming from the subtropical and tropical regions of the world. The growing diversity of flowering trees and shrubs from China and Japan entering Europe was proving to be far more exciting than similar species from temperate North America. Likewise, the succulent species being found in southern Africa were suddenly attracting a large number of enthusiasts, to say nothing of the bulbous species native to northern Africa and parts of

the Middle East. Horticultural interests in temperate North America were waning.

The nomenclatural confusion was rapidly getting out of hand, and the lack of systematic order and a useful classification scheme -- in spite of Ray and Tournefort -- was causing great difficulties in attempting to classify the hoards of species coming to Europe from foreign lands. Publication costs were rapidly escalating so rapidly that publishing became difficult. The idea of the "genus" had been established by Tournefort, who died in 1708, and Ray had proposed groups of genera which we would today call families, but it would remain for the Jussieu family of Paris to complete these ideas, and that would be after 1753. Botanical exploration continued in temperate North America, and many unique species became known in Europe. A twenty year period of neglect would exist from 1700 until 1720 although a review of extant herbaria collections in Europe reveals a period of considerable activity. Without these discoveries being published, however, their existence was not known and it would remain for others to complete the task of describing the native flora of temperate eastern North America.

THE CATESBY YEARS

Mark Catesby is perhaps the best known of the American naturalists. Born and educated at Essex, Catesby found that he had an early interest in the natural sciences due to his close friendship with Samuel Dale, the long-time friend of John Ray. Dale, the Brain-tree apothecary, was the author of the widely used *Pharmacologia* and had acquired a fine herbarium over years of plant collecting. He received

many specimens through his association with the Temple Coffee House Botany Club. With Dale's encouragement, but limited support, Catesby went to Virginia in 1712 to pursue a career as a naturalist. He went to live with his sister, Elizabeth, who had married Dr. William Cocke, an Englishman who had migrated to Virginia in 1710. A friend of the Royal Governor and the Byrd family, these associations proved to be exceedingly useful to Catesby, and he took full advantage of them. Within a week of his arriving in Virginia, Catesby met William Byrd II. Catesby spent a part of the summer and fall of 1712 at Westover, the Byrd estate, and gathered plants in the area where John Banister had roamed 20 years before.

Encouraged by his friendship with Byrd, Catesby began to collect everything he could find. Within a year Governor Spotswood sent to Henry Compton a large consignment of seeds collected by Catesby along with a number of herbarium specimens. The specimens went to Dale, and he informed the members of the Temple Coffee House Botany Club of the many interesting and finely collected plants he was receiving from Catesby. By 1715 James Petiver was writing to Catesby and asking the young naturalist to send him specimens.

When Catesby returned to England in 1719, he brought with him a large collection of vascular plants which he gave to Dale. Dale shared the Catesby material with William Sherard at Oxford who, from 1703 to 1717, had been consul at Smyrna, a city-state in Turkey. Sherard had been working on a revision of Bauhin's Pinax and had much of it completed prior to 1703. He had urged Tournefort to complete

the task when he assumed his diplomatic post, but as Tournefort died in 1708, the revision remained undone. With the arrival of Catesby collections from Virginia, coupled with the large number of species described since 1703, Sherard once again turned his attention to the Pinax revision.

An aspect that certainly aided Sherard in this matter was the annotations Sherard found on the Catesby specimens. Dale had attempted to keep up with the synonymy of his day, and many of Catesby's specimens he had been able not only to identify but assign synonymies as well. Nonetheless, fully half of what Mark Catesby brought with him was new.

Catesby was certainly in the mold of Banister. He made careful notes and field sketches -- which may still be examined today on his specimens -- and knew what had been gathered in Virginia previously as well. Catesby knew that the coastal plain had been well collected, not only by Banister and others who had been in Virginia, but Jones, Krieg and Vernon in Maryland. He proceeded to work further inland, and even went into the mountains.

Upon his return to England in 1719, Catesby was well known, but only to Dale, Petiver and a few other members of the Royal Society. Through a series of well placed letters with rather blunt hints, and a careful distribution of herbarium specimens, Dale managed to impress critical members of the Society with the value of Mark Catesby as a collector. Within a few months, the Society was discussing the idea of sending Catesby back to the New World. As before it was necessary to find a position,

and to arrange for a group of supporters.

The Society had worked closely with Francis Nicholson in the past, and once again the members turned to him for help. Nicholson was now the Royal Governor to South Carolina, having been appointed to that post in 1720. In October Nicholson informed the Society that he would permit Catesby to come to the Colony and would provide him with an annual sum of 20 pounds per year.

The history of botanical explorations in the Carolinas prior to the time Mark Catesby arrived was one marked with the occasional visitor and residents interested in natural philosophies. Even so South Carolina contributed little compared to Virginia prior to 1720, and North Carolina was even less significant during that period of time. Various people gathered plants in the Carolinas with the help of James Petiver. Robert Stevens, who lived near Charles Town, sent plants and animals to Petiver starting in 1698. From the same area Edmund Bohun and Robert Ellis sent more plants to Petiver, and these were described by Petiver, and Plukenet, prior to 1705. A British sea captain, William Halsted, also gathered plants from 1699 to 1703. The collections of Bohun and Ellis led to many significant additions to Petiver's volumes of dried plants.

One of the first women to actively collect in temperate North America was Hannah Williams of South Carolina. She sent a large collection of butterflies to Petiver in 1701 followed by a few plant specimens by 1705. These collections would have been more significant had they been published, but only a few ever

saw the printed page, and most were lost to the botanical community when Petiver died in 1718.

One of the finest collections of Carolina plants ever received from the Carolinas prior to 1720 was made by the Reverend Joseph Lord. American by birth and educated at Harvard University, Lord was well known to the small circle of New England scientists active in that area before he went to South Carolina in 1695. He settled in Dorchester on the banks of the Ashley River, and under the guidance of Hannah Williams, began to collect plants in 1701. Lord's large and well preserved specimens are readily noted in Petiver's extant volumes of plant specimens because of their elegantly hand written labels.

Lord wrote yearly to Petiver and sent him specimens with nearly every letter until 1711. Many of Lord's collections were described by Petiver in his *Gazophylacii naturae & artis*, a series of pamphlets that were published from 1702 to 1709. In 1712, Petiver described a few of Lord's fern species in his *Pterigraphia americana*.

Another of James Petiver's Carolina correspondents was John Lawson. An apprentice in the London Society of Apothecaries before he went to North Carolina in 1701, Lawson was knowledgeable in the art of collecting plants and securing information on the medicinal significance of exotic plant species. Although there was some initial correspondence between Lawson and Petiver in 1701, there was no major exchange until 1709 when Lawson published his own account of the natural history of North Carolina. This book was mainly an overview of the natural

history, topography and geology of the Colony, and although he traveled widely he seems to have made only a few collections prior to 1709. Lawson met Petiver when he was in London shortly before his book was published. He was the Surveyor General of the Colony, in a position to explore, and both Petiver and George London encouraged him to send them specimens and seeds. A gift of several books and assorted items necessary to gather animals, insects and plants helped to solidify their relationship. During Lawson's surveying of the Virginia and North Carolina boundary, he was able to gather many specimens and to enrich Petiver's holdings of plants, insects, snakes, birds and fossils. In 1711 Lawson was hoping to collect extensively and wrote Petiver of his plans. Unfortunately he was captured by Tuscarora Indians who were displeased with his attitude toward them and killed him. His last book of plants had been sent to Petiver but two months prior to his death in September of 1711.

John Lawson's *A new voyage to Carolina* proved to be a successful publication and greatly strengthened his place in history. Unfortunately this travel account does not contain information on the plants of the Colony in a form useful to subsequent generations of taxonomists and he was destined to play no role in the future of systematic botany.

When Catesby came to the New World in May of 1722, he was already well known for his knowledge of plants and animals. The community of Charles Town, South Carolina, was one place where naturalists were considered an asset.

Catesby left London well sponsored with the financial support of not only the Royal Governor but that of four noble peers of the realm and a number of English gentlemen naturalists, notably Sloane, Dale, William Sherard and Mr. Charles Du Bois, the secretary to the East India Company. His efforts were sanctioned by the Royal Society and carried out with the unofficial but nonetheless important approval of the Temple Coffee House Botany Club. As he had done in Virginia, Catesby quickly established himself in the best of social circles in Charles Town, and became an influential member of the Colony.

Catesby traveled widely in the Carolinas, visiting both North and South Carolina, portions of Georgia, and even northern Florida. He went westward as far as the Piedmont. In the winters he often went to Bermuda and Bahama collecting and illustrating the plants he encountered. He gathered many different kinds of seeds and fruits which he sent to England, and these were rapidly incorporated into the gardens of western Europe. He gathered all objects of nature he could find -- animals of all kinds, fossils, plants of nearly every type -- and managed to make notes on geology, anthropology and ecology. Pleased with his efforts in the Carolinas, he attempted to encourage his London supporters to finance an expedition to Mexico, but this proved impossible. When the term of Francis Nicholson ended in 1725, Catesby traveled to Bermuda where he spent nearly a year, and then, in 1726, continued on to England.

Upon his return to London, Catesby immediately set to the task of publishing his *New World discoveries*. He

proposed to his supporters that he begin to prepare a volume on the natural history of the areas he had visited. Unfortunately for Catesby, his supporters in London were no longer willing to provide him with the means to pursue his goal, now that he was not providing them with specimens, and he was forced to find other means. Faced with this problem, Catesby learned to etch his own copper plates for the proposed colored illustrations which he then had to paint himself. He found work in various nurseries where he was able to observe many of his own species growing in cultivation and was able to adjust his illustrations and descriptions accordingly. With a no-interest loan from Peter Collinson, Catesby was able to proceed and finally release, in parts, his *Natural history of Carolina, Florida, and the Bahama Islands*.

Initially, William Sherard at Oxford assisted Catesby in the preparation of his text, and especially with the complex nomenclature. Given that many of Catesby's American plants had previously been described based on Virginia, Maryland and New England collections, Sherard's knowledge of these plants was valuable.

Catesby's close relationship with Sherard caused him some minor difficulties in his friendship with Sir Hans Sloane. While Catesby had been in the New World collecting for his English patrons, they had been waring among themselves over his material. This battling was most fiercely waged between Sloane and Sherard. To a large degree this was only a manifestation of their longer and more serious disagreement over the access Sherard requested of the many collections of plants Sloane had acquired. In parti-

cular, Sherard wanted access to the collections formerly owned by James Petiver. Sherard's work on his revision of Bauhin's *Pinax* had been delayed by his inability to thoroughly study Petiver's many collections. By the early 1720s, this became a major source of difference between the two men.

In 1721, Sherard asked a German botanist, John Jacob Dillenius, to come to England and assist him in completing the *Pinax* project. Sherard lived in London at the time, and had asked Sloane on numerous occasions to be allowed to review the Petiver collections so he could complete his revision. Sloane initially refused saying that the Petiver material was too disorganized and as soon as he had it in order Sherard could see it. When Sherard finally did get an opportunity to examine Petiver's collections, he complained that Sloane had so ill treated them that they were essentially useless. Their dispute over access to the Petiver collections continued for years, and when Sherard died in 1728, the feelings of ill-will were passed -- in a less intensive form -- to Dillenius.

Dillenius did what he could with the Petiver polynomials, and other scientific names proposed by Plukenet (Sloane had his collections as well) and others. By using the herbarium at Oxford, and the expertise of Dillenius, Catesby was able to associate his material with species previously described by Ray, Plukenet, Petiver and others. This was aided by Samuel Dale's own annotations of Catesby specimens that had come to England while Catesby was still exploring. As a result, Dale, who had Ray's personal herbarium, had been able to

accurately name many of Catesby's specimens based on his own knowledge of the Virginia and Maryland flora for he had duplicates of many of those species in his own herbarium.

By the spring of 1729 Catesby had completed the first portion of his book and began to release it as sets of numbered plates with the idea that the book be assembled later when it was completed. He presented each set of plates to the Royal Society, and by November of 1732 had given the Society all the parts that constituted the first volume. It was Collinson who finally presented Catesby to the Society for membership which was granted in May of 1733.

Parts for the second volume began to appear in January of 1735, and Catesby continued to give each part to the Royal Society until all parts were completed in December of 1743. In 1747 he published an appendix and Collinson was able to write to Linnaeus that Catesby's book was completed. By this time the number of contributors to Catesby's project had grown to 166, well above the original 12 men who had provided him with his initial funding. Included among his large list of patrons were a number of royal families, ambassadors, botanical friends and colleagues. Perhaps most significant was the number of supporters from the American colonies. The final book contained 200 colored plates of which 171 plants were figured.

Catesby's three-year effort in South Carolina has come to be regarded as one of the most significant in American natural history explorations. He has come to be regarded as one of the more skilled observer. Catesby was

only an adequate illustrator, but the combination of his art work and his ability to observe plants and animals has been appreciated by many since 1747, including Linnaeus.

THE LINNAEAN ERA

For temperate North American botany, the Linnaean era began when Carl Linnaeus came to England. He came to examine the collections of dried plants that had been gathered in the New World by the host of collectors starting with John Banister and essentially ending with Mark Catesby. For Linnaeus, Catesby's work was highly significant. The full text of volume one and much of the second had been printed when Linnaeus visited Dillenius in August of 1736. At Oxford, Linnaeus could not only examine the plates, but the actual specimens as well for Sherard and Dillenius had received nearly a full set of Catesby's specimens. In addition, Linnaeus could examine collections made by earlier naturalists, now annotated by Sherard and Dillenius with the polynomials proposed by various workers up to the mid-1730s. Previously in Sweden, Holland and France Linnaeus had been able to see garden material, and had been able to characterize many of the New World species on the basis of these plants. He continued to examine garden material, including the botanical garden at Oxford, but it was his examination of herbarium material that proved critical to his understanding of the application of polynomials.

Linnaeus' visit to England made it possible for him not only to see many collections which he had been unable to study previously, but also gave him an opportunity to talk with many of the men who had been supporting

explorations throughout the world. He visited Sir Hans Sloane briefly but did not examine his herbarium. He formed a bond with Peter Collinson and James Ellis which was to carry him through the difficult period of acceptance of the Linnaean methodology in England after 1753.

While Linnaeus certainly concentrated upon the collections made by Catesby and others who collected in the southern portion of the eastern seaboard, he did see a few collections made by collectors from the New England area and eastern Canada. Botanical explorations had been occurring in the north while similar activities were being carried out in the south. John Josselyn was one of the first naturalists to visit this area, traveling to the present state of Maine in 1638 and 1639. On this trip he had little time to observe plants and animals, but on a much longer trip, lasting from 1663 to 1671, Josselyn had ample opportunity to make observations. He was primarily interested in medicinal plants of the area, and discovered that the herbals of his day were inadequate in dealing with the New England vegetation. He made no permanent collections and apparently failed to gather seeds of the plants he saw. Nonetheless, Josselyn prepared a small book with rather crude and poorly executed illustrations of the New England flora. Entitled *New-England's rarities discovered*, his book was published in 1672; it proved to be of little interest to Linnaeus and it is not mentioned in *Species plantarum*.

Other collectors came to the New England area, but few were of major importance. Thomas More, the so-called "Pilgrim Botanist", came to New England in 1722 and stayed

until 1724. When five Mohawk chiefs were sent to England in 1709, More apparently met them and received an invitation to visit them in the New World. He had worked with Bobart at Oxford in the 1690s, and had collected for Lhwyd in south-western England in the 1700s. He was an enthusiastic collector, and Petiver, Sloane and others encouraged him to go to America. It was Sherard who secured the funds for More to travel, but had some difficulties in finding supporters for a man in his 40s.

More proved not to be as successful a collector as William Sherard and his supporters had hoped. Nonetheless More's many collections were in the Sherardian Herbarium at Oxford University when Linnaeus visited it in 1736. So too were the Hudsons Bay collections of Richard Tilden who had visited the area in about 1700. These collections were likely the only plants from northern Canada that Linnaeus examined in the herbarium.

No doubt Linnaeus and Dillenius, who initially began their association with a profound distrust of the other, discussed Dillenius' book *Hortus elthamensis* which had been published in 1732. Dillenius had discussed nearly 400 species grown in the garden of James Sherard, the brother of William Sherard. Dillenius had served Sherard up to his death in 1728 with a high degree of devotion, and hoped to complete the revision of the *Pinax* which Sherard had so long struggled to see done. Shortly after William's death, James asked Dillenius to prepare a book on his garden. Dillenius, the first Sherardian Professor at Oxford was reluctant to do it, but bowed to the will of the wealthy James and devoted several years to the book. The *Hortus*

was well illustrated and well written with abundant information on the distribution and taxonomy of each species mentioned. It was a major work and Linnaeus recognized it as such almost immediately. At Oxford, Linnaeus was able to review both the garden and herbarium specimens Dillenius had used in arriving at his taxonomic disposition of the species he treated. Included in the Hortus were a number of Banister and Catesby species as well as a few grown from seed gathered in Maryland. The synonymy was rather complete, and Linnaeus could examine the herbarium specimens annotated by Sherard, Dale, Dillenius, and even Ray, Plukenet and Petiver, and trace the evolution of the species definitions Dillenius had arrived at. Linnaeus remained at Oxford for some eight days. It was here he studied the extant collections of American plants which aided him when he assisted Johann Friedrich Gronovius during the winter of 1737-1738 in classifying a large collection of plants from Virginia gathered by John Clayton.

CLAYTON AND KALM

Colonial Virginia of John Clayton's era was different from that Mark Catesby had found in 1712. The population was growing and the urban centers, once restricted to the Williamsburg area, had expanded far beyond the James River. There was a college, William and Maryland, newspapers in the Colony -- something that was not even yet in the city of Oxford in England. The intellectual center that had once been Williamsburg was now divided between Boston, Philadelphia and Charles Town. Virginia still held great families, and most were deeply interested in botany and the other natural philosophies.

Some maintained large gardens and were exchanging plants not only with European centers, but even with other colonists elsewhere in the New World.

Clayton came to Virginia in 1720, joining his father who was in the government and would eventually rise to be the Attorney General. Clayton acquired land and became an established, although modest, man of means. He too became a government official serving as the Clerk of Gloucester County where he lived. Like other Virginians, Clayton established a garden and entered into a correspondence with William Byrd II, Mark Catesby who he had met, Peter Collinson, and John Bartram. It was Catesby that sent Clayton's herbarium specimens to Gronovius, and others, and urged Gronovius to carefully review them with the hopes of classifying them.

In the 1730s Clayton prepared a manuscript entitled "A catalogue of plants, fruits, and trees native to Virginia" and sent it to Gronovius. Taking the manuscript, and the many Clayton specimens which had been gathered since the early 1730s, Gronovius and Linnaeus began to prepare a manuscript on the collection following Linnaeus' new system of classification rather than that of Ray's which Clayton had followed. Linnaeus worked closely with Gronovius on the nomenclature of the Virginia species he was publishing. The knowledge Linnaeus had gained from his visit to Oxford where he examined so many of the temperate North American plants proved to be useful to him in naming the plants gathered by John Clayton and assigning synonymies to them. The resulting book, *Flora virginica*, appeared in two parts. The first part was published in 1739, while the second was released in 1743. Linnaeus

was closely involved with the 1739 volume, but had little direct influence on the 1743 volume.

When Clayton saw Gronovius' *Flora virginica*, he was profoundly disappointed. Not only had Gronovius not followed his manuscript, he failed to acknowledge Clayton's contribution to the *Flora* except for a minor comment in the introduction. To remove this as a possibility from happening again, Clayton began to study the Linnaean methodology with the hopes of writing his own flora of Virginia. Within a few years he was able to write his fellow naturalists commenting on various technical points in their descriptions, and he turned his attention to writing. In 1748, Clayton traveled into the high mountains collecting many new species which were as yet unknown to Linnaeus. In 1757, Clayton sent to Peter Collinson his revision of *Flora virginica*, but could not find a publisher. Collinson asked George Ehret to prepare plates for the volume, and some were actually prepared before Laurenz Gronovius, the son of Johann, published his father's long delayed second edition in 1762. The book forced Clayton and Collinson to abandon their project, and Clayton's revision was never published.

Clayton had proposed to use Linnaeus' binomial system of nomenclature, and this plus the Ehret plate would have made his flora a major publication. The younger Gronovius did not illustrate his work, nor did he use binomial nomenclature, and thus this post-1753 flora is of no systematic importance and is little used today.

A causal friend of John Clayton's was John Mitchell, a man of many talents and broad

scientific interests. He was in Virginia from 1735 to 1746, returning to England because of his ill health. He also maintained a large and active correspondence with many contemporary scientists including botanists such as Dillenius, Gronovius and Linnaeus. He knew John Bartram as well and communicated with Cadwallader Colden in New York, Peter Collinson in London, and even two of his fellow American collectors, Mark Catesby and Peter Kalm.

Mitchell's interests in botany were somewhat limited compared to his contributions made in other areas of the natural and physical sciences. Nonetheless he sent numerous herbarium specimens to Dillenius and Collinson, along with various manuscripts. One of those papers was published for him by Collinson in 1748; it was titled "*Nova genera plantarum Virginiensium*." This work was consulted by Linnaeus and various of Mitchell's new genera were mentioned or treated in the first edition of *Species plantarum*.

Of perhaps greater interest were Mitchell's comments regarding species and speciation. He proposed that plants and animals that can produce offsprings biologically ought to be considered members of the same species, whereas those that cannot produce offsprings or the off-spring are sterile, ought not to be regarded as the same species. Mitchell felt Linnaeus' artificial system was unreliable, and that Ray's system was much more natural as it was based on attributes of the whole plant rather than just the reproductive parts.

Even though Mitchell sent the bulk of his plant collection to England in 1738 and in 1742, he continued to botanize

in Virginia. When he departed in 1746, he took more than a thousand specimens. Unfortunately, his ship was stopped by a Spanish vessel carrying pirates and the contents plundered. Although Mitchell was eventually able to recover the majority of his collection, most of it was no longer useful and was discarded.

Both Mitchell and Clayton correspondent with Cadwallader Colden of New York. Colden had initially come to America in 1715. Born in Ireland of Scottish parents and educated at the University of Edinburgh, he studied medicine in London before going to Philadelphia where he practiced medicine among other enterprises. Colden became a correspondent of James Petiver but apparently never sent him any plant collections. He returned to Scotland in 1715, married, and returned to Philadelphia where he continued to practice. However, in 1720, Colden was appointed Surveyor General for New York, and he moved to that Colony where he entered into a life of politics.

As a result of his position, Colden was able to become wealthy through land speculation, and was slowly able to amass an excellent scientific library. His correspondents were the "who's who" of American and European science, with Collinson, Gronovius and Linnaeus among his European associates, and the Bartrams, Clayton, Mitchell, and a host of others who would be important after 1753 among his American friends.

In 1739, Colden and his family left New York City and moved to an estate he named Coldenham. Here he was able to write on a wide variety of subjects, including botany. Colden began to study the

Linnaean method in the early 1740s, and with a visit from John Bartram in 1742, Colden began to seriously study the local flora. He collected plants around Coldenham and sent them to Collinson, and even Linnaeus. Soon he was attempting to prepare a flora of New York, but mostly it was a treatment of plants in his local area. Published from 1743 to 1750, in parts, Colden's "Plantae Coldenhamiae in provincia Novaboracensi Americanae sponte Crescentes" was used by Linnaeus to characterize a few species in 1753.

Colden, like Mitchell, found fault in Linnaeus' sexual system, and voiced his disagreements about it to Gronovius. Within a few years, Colden's interest in botany had lessened, but he was pleased to discover that his curiosity in botany had passed onto his daughter, Jane.

Jane Colden began her studies of botany about the time *Species plantarum* appeared in print, and by 1755 her father wrote to Gronovius of her pronounced skills. What correspondence he was receiving on botany, Cadwallader passed on to Jane. She exchanged seeds and specimens with several American and European naturalists, and entered into her own correspondence with a number of new naturalists, such as Alexander Garden, of Charles Town, South Carolina, who, like herself, would be a significant contributor to Linnaeus' subsequent editions of *Species plantarum*. She prepared a "Flora of New York" with some 340 illustrations, but the work was never published during her lifetime. It now appears that this failure was more one of her sex than her ability.

Both of the Coldens knew John Bartram. Of all the

Linnaean era collectors in temperate North America, John, and his son William, collected more widely than any other naturalist. Born in Pennsylvania of Quaker parents in 1699, he moved at an early age to North Carolina where he remained until a young man. Upon the death of an uncle, Bartram inherited a small farm near Darby, Pennsylvania, where he moved. After the death of his first wife in 1827, John and his only surviving son moved to a new farm on the Schuylkill River, a short distance from Philadelphia, and established a small but significant botanical garden on about five acres.

Bartram prospered in his farming, married again, and he was able to expand his land holdings so that he could eventually devote much of his personal time to botany and horticulture. Bartram soon began to explore the field of botany, mainly from the aspect of medicinal plants, learned to read Latin, and to read Linnaeus' writings as they became available in America. He also learned to read and write English, and while neither his English or Latin were perfect, he managed, and became an active participant in the scientific circles that dominated the lives of a handful of men in the Philadelphia area.

Like so many others, Bartram began to correspond with Collinson in the 1730s, and to exchange specimens. Some of his collections of dried plants went to Dillenius at the University of Oxford, and these Linnaeus likely saw in August of 1736. Collinson became an enthusiastic supporter of Bartram and encourage all who went to America to meet him. He also encourage John Bartram to travel and collect plants which would be of potential value to the horticultural

gardens of Europe. By 1738 Bartram was proposing to Collinson to make a major botanical expedition into Virginia, and to this Collinson added even more encouragement in the form of letters of introduction, instructions and advice of the art of collection live specimens, seeds and herbarium specimens. Soon Bartram seeds were sprouting in the Oxford Botanical Garden as well as at the Chelsea Physic Garden. Within a few years Collinson had established a network of supporters for Bartram, and as the years past, Bartram made more and more exciting discoveries of unique species heretofore unknown in the scientific world.

Many of Bartram seeds found their way to Linnaeus' garden at Uppsala, and through them, Linnaeus was able to describe new species of American plants. Bartram was now traveling widely in the eastern part of temperate North America, visiting not only the more remote regions along the coastal plain but the high mountains to the west. He made repeated visits into the colonies of New Jersey, Delaware and Maryland, as well as numerous trips to various parts of Pennsylvania. He traveled into New York, met Colden, and collected in the Catskills. He even went into southern Canada.

The Bartrams continued to collect well past 1753, and John and his son William soon became well known for their fine plants and exotic travels to the more remote regions of the eastern seaboard. The two men rose to fame, with John being appointed the Royal Botanist in 1765. Their 1765 expedition to Florida has become famous due to John's book published the following year.

A visitor to the Bartram

farm in the summer and fall of 1748 was Peter Kalm. A student of Linnaeus', Kalm came to America in September of 1748 with the intention of collecting dried herbarium specimens and seeds for his major professor. During his first fall he gathered seeds and plants in Pennsylvania and New Jersey, while during the summer he turned northward to New York and southern Canada returning to Philadelphia for the winter of 1749-1750. Kalm then traveled from New Jersey across western Pennsylvania to the eastern Great Lakes, collecting numerous specimens before his return to the city of Brotherly Love in October of 1750. From there, in February, 1751, Kalm sailed for England, and then on to Sweden where he arrived, laden with many new and wondrous plants, upon his professor's door at Uppsala in June.

SPECIES PLANTARUM

For Carl Linnaeus, the idea of a treatment of all the species of the world began initially in 1733 when he first attempted the project. Almost at once he realized it was impossible to do at that time and turned to describing the genera instead. This effort resulted in the publication of *Genera plantarum* published in 1737. He then took a copy of his new book, added blank pages, and began to insert species. From 1746 to 1748 Linnaeus worked on a new draft of *Species plantarum* following the format he had used in his earlier (1738) *Hortus cliffortianus*. This proved equally impossible but for somewhat different reasons.

Much as Morison, Sherard and Dillenius realized the need for a new edition of Bauhin's *Pinax*, so too did Linnaeus when it came to eval-

uating the vast number of polynomials that had been published since 1623. The prospects of producing such a work was essentially nil. Dillenius had been unable to complete the work, and no doubt Linnaeus knew of the difficulties he had had and did not see any way of resolving them himself. Linnaeus did not see himself traveling now to London and reviewing the many collections or publications associated with the writings of Ray, Plukenet and Petiver, and it was these writers, in large part, whose nomenclature was uncertain and confusing. Certainly Linnaeus had been able to resolve some problems when working Gronovius' *Flora virginica*, but all of these men had published new species from all over the world. Particularly troublesome was Petiver who had described plants not only from the Carolinas and Florida of North America, but species from the interior of China, far off Japan, and remote regions of Africa. Linnaeus could not have a complete list of synonymy for each and every name; he had to rethink his manuscript and what he hoped to accomplish.

Linnaeus completed about a third of his proposed *Species plantarum* from 1746 to 1748 when he once again set the manuscript aside. He had not yet adopted the binomial system of nomenclature, and other duties prevented him from devoting the kind of time needed to do the work. Thus the book remained until June of 1751 when Kalm returned with his collections.

Perhaps spurred by Kalm, Linnaeus once again started on *Species plantarum*. By this time he had conceived of the handy index device of proposing a trival name -- what we today call the species epithet -- to go along with the cor-

rect scientific phrase name by which each species should be known. He realized that he had to shorten his synonymy and could economically treat the world's flora only by having brief statements as to the distribution of each kind and a phrase name that did not exceed two lines of print. Taking his 1746-1748 draft, Linnaeus managed to rewrite the first 67 printed pages of his book in eight days. By November of 1751 Linnaeus had completed the first part (or volume), and by March, 1752, three-quarters of the text was written. In July, 1752, or some 13 months after beginning, Linnaeus completed the text for *Species plantarum*.

It is obvious that Linnaeus went quickly through the early sections of part one, and took most of his information from his previously written manuscript. Only a few Kalm collections were cited, and occasionally the same species is described later in the book -- in its correct position according to Linnaeus' scheme of classification -- with Linnaeus not realizing he had already treated that plant. As one proceeds through the text, Linnaeus became more exact in the citation of geographical location, his dependency upon Kalm collections for species from temperate North America increased, and he became more critical of his circumscription of genera. However, toward the latter third of the book, as Linnaeus' health began to fail and he began to hurry, he once again is not as careful and more errors appear. *Species plantarum* is filled with errors, omissions and an occasional strange statement. Linnaeus occasionally miscites places of publication, wrong page numbers or the incorrect figure or plate. In some instances Linnaeus gives seem-

ingly bizarre polynomials in synonymy, as in the case of the two Maryland-based polynomials he cites under *Osmunda cinnamomea*. A simple reading of the Latin would quickly show that these names ought to apply to *O. claytoniana*, yet Linnaeus puts them where he did. In large part this was likely due to his observation of specimens of *O. cinnamomea* at the Sherardian Herbarium at the University of Oxford where the names are given on the labels. However, there are also collections, from Maryland, of *O. claytoniana* with these same names also listed, so it is difficult to understand his treatment of these polynomials.

When the first edition of *Species plantarum* appeared in 1753, the botanical community did not immediately jump onto the Linnaean binomial bandwagon. His system of classification had long been criticized, and his critics were not immediately willing to embrace all of his radical departures from the established norm. It is important for modern workers to remember that even Linnaeus did not regard his trival names as anything more than a convenient means of index his book. Philip Miller, the English gardener who had published his *Gardners dictionary* for years, did not use binomials until 1768 -- and then only in this one edition -- mainly because his book was already alphabetically arranged and did not need an index.

The impact of Linnaeus' *Species plantarum* was slowly felt. When copies finally arrived on the American continent, some workers immediately adopted the idea of binomials. Jane Colden in her unpublished book used them, as did the Bartrams. Alexander Garden used binomials when the species were known to him, but

otherwise used phrase names in his correspondence. With the appearance of the second edition of *Species plantarum* in 1762 and 1763, the trend to binomials was largely established. This new edition contained many new and exciting plants that Linnaeus had received from Jane Colden, Alexander Garden, John and William Bartram, and others who were now exploring and collecting in temperate North America. Not only would Linnaeus describe these species, but others, working elsewhere in Europe, would pen validly published scientific names to these collections and they too would rise in the world of systematic botany.

With the opening of the vast lands to the west of the Appalachian Mountains, the end of French and Indian War in America, and the growth of resident naturalists in America, the field of plant taxonomy and discovery would expand rapidly soon leaving behind even Linnaeus for a new generation of taxonomists.

SUMMARY OF SPECIES PLANTARUM

One cannot understate the impact Linnaeus had upon systematic botany. His work, and especially his first edition of *Species plantarum* have, by international agreement, become highly significant to all modern-day workers in the field of systematic botany. Nonetheless, a careful evaluation and summation of Linnaeus' state of knowledge regarding the vascular plants of temperate North America as defined by Linnaeus in 1753 has not been attempted. In the following section such a review will be made. It is necessary first to prepare listings of species, workers and publications by categories, and then to comment on these findings. The following

categories will be given:

1--All species mentioned in the first edition of *Species plantarum* with direct or indirect indication as being in temperate North America. This will include obvious errors made by Linnaeus.

2--A breakdown of these species into distributional groups as reported by Linnaeus, listing first those which Linnaeus considered to occur only in temperate North America; those which occur in this area but also occur elsewhere in the New World; those that occur in temperate North America and elsewhere in the world; and finally, those which Linnaeus does not specifically state to occur in temperate North America, but gives a synonym or reference based upon a temperate North American collection.

3--A listing of vascular plants species named for geographical regions in temperate North America.

4--A listing, with the collection numbers insofar as possible, of the Clayton collections Linnaeus indirectly cites when he gives Gronovius' *Flora virginica* among the list of publications he cites. In some instances Gronovius is basing his information upon a collection made by someone other than Clayton, and this can be discovered only by consulting the Gronovius herbarium now at the British Museum (Natural History) in London. In perhaps a few other instances, Gronovius may not have had a collection in hand, but garden material or information about a particular species in temperate North America and reported it. Most of these particular kinds of instances can be seen for those species which lack a Clayton collection number. However, it must be immediately noted that Clayton did send some specimens without collection numbers, and in some

cases Gronovius failed to give the number in his publication, so it cannot be assumed that in all instances where there is no number that this reference is based on something other than a Clayton sheet. The first list is by species name; the second list is by collection number.

5--A listing of seemingly Kalm collections taken not only from statements in *Species plantarum*, but from Linnaeus' herbarium at the Linnean Society in London. This is to be regarded as an incomplete listing, as it will require a more careful examination of Kalm collections housed elsewhere, and a comparison of those specimens with others in Linnaeus' herbarium before a more finalized list can be completed.

6--A listing of major publications cited by Linnaeus for those species which, in Linnaeus' opinion, were endemic to temperate North America in their distribution. Species with a more widespread distribution are not reviewed at this time.

7--A list of synonymies for some of the vascular plant species reported for temperate North America in Linnaeus' first edition of *Species plantarum*.

The review that follows is based upon observations obtained from these listings. In all instances they should be considered somewhat tentative as it has not been possible to correct errors made by Linnaeus, especially in the citation of literature, nor to review fully the holdings in the British Museum (Natural History) of the Gronovius herbarium. Until these and other similar tasks are completed it will be impossible to fully report on all aspects relative to the first edition of Linnaeus *Species plantarum*.

Vascular Plants from Temperate North America Mentioned in Linnaeus' First Edition of *Species Plantarum*

Acalypha virginica
Acer negundo
Acer pensylvanicum
Acer rubrum
Acer saccharinum
Acnida cannabina
Acrostichum areolatum
Acrostichum platyneuros
Acrostichum polypodioides
Actaea racemosa
Actaea spicata var. *alba*
Adiantum pedatum
Aesculus pavia
Agave virginica
Ageratum altissimum
Agrimonia eupatoria
Agrostis virginica
Aletris farinosa
Alisma cordifolia
Alisma subulata
Allium canadense
Alsine media
Alyssum hyperboreum
Amaranthus graecizans
Amaranthus hybridus
Amaranthus hypocondriacus
Amaranthus lividus
Amaranthus retroflexus
Amaranthus spinosus
Amaryllis atamasca
Ambrosia artemisiifolia
Ambrosia elatior
Ambrosia trifida
Ammannia ramosior
Amorpha fruticosa
Anchusa virginiana
Andromeda arborea
Andromeda calyculata
Andromeda mariana
Andromeda paniculata
Andromeda racemosa
Andropogon alopecuroides
Andropogon divaricatum
Andropogon hirtum
Andropogon nutans
Andropogon virginicum
Anemone dichotoma
Anemone hepatica
Anemone quinquefolia
Anemone thalictroides
Anemone virginiana
Angelica atropurpurea
Angelica lucida
Angelica sylvestris
Annona glabra

<i>Annona muricata</i>	<i>Aster undulatus</i>
<i>Annona triloba</i>	<i>Aster vernus</i>
<i>Anthericum calyculatum</i>	<i>Astragalus canadensis</i>
<i>Antirrhinum canadense</i>	<i>Astragalus carolinianus</i>
<i>Antirrhinum elatine</i>	<i>Atriplex halimus</i>
<i>Aphanes arvensis</i>	<i>Atriplex laciniata</i>
<i>Apocynum androsaemifolium</i>	<i>Avena pensylvanica</i>
<i>Apocynum cannabinum</i>	<i>Avena spicata</i>
<i>Aquilegia canadensis</i>	<i>Azalea lutea</i>
<i>Arabis canadensis</i>	<i>Azalea viscosa</i>
<i>Arabis lyrata</i>	
<i>Aralia nudicaulis</i>	<i>Baccharis foetida</i>
<i>Aralia racemosa</i>	<i>Baccharis halimifolia</i>
<i>Aralia spinosa</i>	<i>Baccharis ivaefolia</i>
<i>Arbutus uva-ursi</i>	<i>Bartsia coccinea</i>
<i>Arenaria rubra</i> var. <i>marina</i>	<i>Betonica annua</i>
<i>Arethusa bulbosa</i>	<i>Betula lenta</i>
<i>Arethusa divaricata</i>	<i>Betula nigra</i>
<i>Arethusa ophioglossoides</i>	<i>Bidens bipinnata</i>
<i>Aristolochia arborescens</i>	<i>Bidens bullata</i>
<i>Aristolochia serpentaria</i>	<i>Bidens frondosa</i>
<i>Arnica maritima</i>	<i>Bidens nivea</i>
<i>Arum dracontium</i>	<i>Bidens pilosa</i>
<i>Arum triphyllum</i>	<i>Bignonia caerulea</i>
<i>Arum virginicum</i>	<i>Bignonia capreolata</i>
<i>Arundo phragmites</i>	<i>Bignonia catalpa</i>
<i>Asarum canadense</i>	<i>Bignonia crucigera</i>
<i>Asarum virginicum</i>	<i>Bignonia radicans</i>
<i>Asclepias amoenia</i>	<i>Bignonia sempervirens</i>
<i>Asclepias decumbens</i>	<i>Briza eragrostis</i>
<i>Asclepias incarnata</i>	<i>Bromus ciliatus</i>
<i>Asclepias nivea</i>	<i>Bromus purgans</i>
<i>Asclepias purpurascens</i>	<i>Buchnera americana</i>
<i>Asclepias rubra</i>	<i>Bunias cakile</i>
<i>Asclepias syriaca</i>	<i>Bupthalmum frutescens</i>
<i>Asclepias tuberosa</i>	<i>Bupthalmum helianthoides</i>
<i>Asclepias variagata</i>	<i>Burmanna biflora</i>
<i>Asclepias verticillata</i>	
<i>Ascyrum crux-andreae</i>	<i>Cacalia atriplicifolia</i>
<i>Ascyrum hypericoides</i>	<i>Cacalia porophyllum</i>
<i>Ascyrum villosum</i>	<i>Cacalia suaveolens</i>
<i>Asplenium rhizophyllum</i>	<i>Cactus opuntia</i>
<i>Aster annuus</i>	<i>Cactus pentagonus</i>
<i>Aster concolor</i>	<i>Caesalpinia brasiliensis</i>
<i>Aster cordifolius</i>	<i>Callicarpa americana</i>
<i>Aster divaricatus</i>	<i>Callitriche palustris</i>
<i>Aster dumosus</i>	<i>Caltha palustris</i>
<i>Aster ericoides</i>	<i>Campanula americana</i>
<i>Aster grandiflorus</i>	<i>Campanula perfoliata</i>
<i>Aster laevis</i>	<i>Canna glauca</i>
<i>Aster linariifolius</i>	<i>Cardamine virginica</i>
<i>Aster linifolius</i>	<i>Carduus altissimus</i>
<i>Aster miser</i>	<i>Carduus virginianus</i>
<i>Aster mutabilis</i>	<i>Carex folliculata</i>
<i>Aster novae-angliae</i>	<i>Carex pseudocyperus</i>
<i>Aster novi-belgii</i>	<i>Carex squarrosa</i>
<i>Aster puniceus</i>	<i>Carpinus betulus</i>
<i>Aster rigidus</i>	<i>Carpinus ostrya</i>
<i>Aster tenuifolius</i>	<i>Cassia chamaecrista</i>
<i>Aster tradescantii</i>	<i>Cassia ligustrina</i>

- Cassia marilandica*
Cassia nictitans
Cassine peragua
Ceanothus americanus
Celastrus bullatus
Celastrus myrtifolius
Celastrus scandens
Celosia paniculata
Celtis occidentalis
Cenchrus tribuloides
Cephalanthus occidentalis
Cerastium semidecandrum
Cercis canadensis
Chaerophyllum arborescens
Chelidonium glaucium
Chelone glabra
Chelone hirsuta
Chelone penstemon
Chenopodium album
Chenopodium anthelminticum
Chenopodium virginicum
Chionanthus virginica
Chironia angularis
Chironia campanulata
Chironia dodecandra
Chrysanthemum arcticum
Chrysanthemum serotinum
Chrysocoma graminifolia
Chrysogonum virginianum
Chrysosplenium oppositifolium
Cicuta bulbifera
Cicuta maculata
Cinna arundinacea
Circaea lutetiana
 var. *canadensis*
Cissampelos smilacina
Cistus canadensis
Claytonia virginica
Clematis crispa
Clematis viorna
Clematis vitalba
Clethra alnifolia
Clinopodium incanum
Clinopodium vulgatum
Clinopodium vulgare
Clitoria mariana
Clitoria virginiana
Coix dactyloides
Coix lacryma-jobi
Collinsonia canadensis
Commelina communis
Commelina erecta
Convallaria polygonatum
Convallaria racemosa
Convallaria stellata
Convolvulus carolinus
Convolvulus hederaceus
Convolvulus panduratus
Convolvulus repens
Convolvulus spithameus

Conyza asteroides
Conyza bifrons var. *flosculosa*
Conyza linifolia
Coreopsis alternifolia
Coreopsis angustifolia
Coreopsis auriculata
Coreopsis lanceolata
Coreopsis tripteris
Coreopsis verticillata
Cornus canadensis
Cornus florida
Cornus sanguinea
Cracca virginiana
Crataegus coccinea
Crataegus crus-galli
Crataegus tomentosa
Crataegus viridis
Crescentia cujete
Crotalaria alba
Crotalaria perfoliata
Crotalaria sagittalis
Cucubalus stellatus
Cupressus distichia
Cupressus thyoides
Cuscuta americana
Cynanchum hirtum
Cynanchum suberosum
Cynoglossum virginianum
Cynosurus aegyptius
Cyperus arundinacea
Cyperus odoratus
Cyperus strigosus
Cypripedium calceolus

Dactylis cynosuroides
Dalibarda repens
Datisca hirta
Datura stramonium
Daucus carota
Dianthera americana
Dianthus plumarius
Diodia virginiana
Dioscorea villosa
Diospyros virginiana
Dipsacus fullonum
Dirca palustris
Dodecatheon meadia
Dolichos polystachyus
Dolichos regularis
Draba verna
Dracocephalum virginianum
Dracontium foetidum
Drosera rotundifolia

Elatine hydropiper
Elephantopus scaber
Elephantopus tomentosus
Elymus canadensis
Elymus hystrix
Elymus virginicus

Epigaea repens	Gnaphalium plantaginifolium
Epilobium hirsutum	Gnaphalium purpureum
Equisetum arvense	Gomphrena interrupta
Equisetum hyemale	Gomphrena serrata
Erigeron camphoratum	Gratiola dubia
Erigeron canadense	Gratiola virginiana
Erigeron philadelphicus	Guilandina dioica
Eriocaulon decangulare	
Eriophorum virginicum	Hamamelis virginiana
Eryngium aquaticum	Hedera quinquefolia
Eryngium foetidum	Hedysarum canadense
Erythrina herbacea	Hedysarum canescens
Erythronium dens-canis	Hedysarum frutescens
Eupatorium altissimum	Hedysarum hirtum
Eupatorium aromaticum	Hedysarum marilandicum
Eupatorium coelestinum	Hedysarum nudiflorum
Eupatorium hyssopifolium	Hedysarum paniculatum
Eupatorium perfoliatum	Hedysarum repens
Eupatorium purpureum	Hedysarum violaceum
Eupatorium rotundifolium	Hedysarum virginicum
Eupatorium scandens	Hedysarum viridiflorum
Eupatorium sessilifolium	Hedysarum volubile
Eupatorium trifoliatum	Helenium autumnale
Euphorbia corollata	Helianthus angustifolius
Euphorbia ipecacuanhae	Helianthus atrorubens
Euphorbia maculata	Helianthus decapetalus
Euphorbia polygonifolia	Helianthus divaricatus
Euphorbia portulacoides	Helianthus giganteus
Euonymus americanus	Helianthus laevis
	Helianthus multiflorus
Fagus pumila	Helianthus strumosus
Ferula canadensis	Heliotropium indicum
Fraxinus americana	Helleborus trifolius
Fumaria cucullaria	Helonias bullata
Fumaria sempervirens	Heuchera americana
	Hibiscus moscheutos
Galax aphylla	Hibiscus palustris
Galium bermudense	Hibiscus virginicus
Galium tinctorium	Hieracium gronovii
Galium trifidum	Hieracium kalmii
Gaultheria procumbens	Hieracium paniculatum
Gaura biennis	Hieracium venosum
Gentiana ciliata	Hippophae canadensis
Gentiana quinquefolia	Holcus laxus
Gentiana saponaria	Holcus striatus
Gentiana villosa	Holosteum succulentum
Geranium carolinianum	Hordeum jubatum
Geranium maculatum	Horminum virginicum
Gerardia flava	Houstonia caerulea
Gerardia pedicularia	Houstonia purpurea
Gerardia purpurea	Hydrangea arborescens
Geum virginianum	Hydrocotyle americana
Gleditsia triacanthos	Hydrocotyle umbellata
Glycine apios	Hydrophyllum virginianum
Glycine bracteata	Hyoseris virginica
Glycine comosa	Hypericum ascyron
Glycine frutescens	Hypericum canadense
Glycine tomentosa	Hypericum kalmianum
Gnaphalium margaritaceum	Hypericum lasianthus
Gnaphalium obtusifolium	Hypericum mutilum

- Hypericum setosum*
Hyssopus nepetoides

Ilex aquifolium
Ilex cassine
Impatiens noli-tangere
Ipomoea carolina
Ipomoea lacunosa
Ipomoea nyctelea
Ipomoea tannifolia
Iris verna
Iris versicolor
Iris virginica
Isnardia palustris
Itea virginica
Iva frutescens

Juglans alba
Juglans nigra
Juncus bulbosus
Juncus campestris
Juncus effusus
Juncus filiformis
Juniperus virginiana
Jussiaea erecta

Kalmia angustifolia
Kalmia latifolia

Lactuca canadensis
Lamium amplexicaule
Laurus aestivalis
Laurus benzoin
Laurus borbonia
Laurus indica
Laurus sassafras
Laurus winterana
Lechea major
Lechea minor
Leontice thalictroides
Leontodon dandelion
Lepidium virginicum
Ligusticum scothieum
Lilium camschatcense
Lilium canadense
Limodorum tuberosum
Linnaea borealis
Linum virginianum
Liquidambar peregrina
Liquidambar styraciflua
Liriodendron tulipifera
Lithospermum virginianum
Lobelia cardinalis
Lobelia cliffortiana
Lobelia inflata
Lobelia kalmii
Lobelia siphilitica
Lonicera marilandica
Lonicera sempervirens
Lonicera symphoricarpos

Ludwigia alternifolia
Lupinus perennis
Lycopodium alopecuroides
Lycopodium apodum
Lycopodium carolinianum
Lycopodium complanatum
Lycopodium obscurum
Lycopodium rupestre
Lycopsis virginica
Lycopus virginicus
Lysimachia ciliata
Lysimachia punctata
Lysimachia quadrifolia
Lythrum lineare
Lythrum petiolatum
Lythrum verticillatum

Magnolia virginiana
Magnolia virginiana
 var. *acuminata*
Magnolia virginiana
 var. *foetida*
Magnolia virginiana
 var. *glauca*
Magnolia virginiana
 var. *grisea*
Magnolia virginiana
 var. *tripetala*
Malva caroliniana
Medeola virginiana
Medicago virginica
Melanthium virginicum
Melica altissima
Melissa nepeta
Melissa pulegioides
Melothria pendula
Menispermum canadense
Menispermum carolinum
Menispermum virginicum
Mentha canadensis
Mentha spicata var. *viridis*
Mentzelia aspera
Mespilus arbutifolia
Mespilus canadensis
Mimulus ringens
Mitchella repens
Mitella diphylla
Mollugo verticillata
Monarda ciliata
Monarda clinopodia
Monarda didyma
Monarda fistulosa
Monarda punctata
Monotropa hypopithys
Monotropa uniflora
Morus rubra
Myosotis virginiana
Myrica asplenifolia
Myrica cerifera

Napaea dioica	Philadelphus inodorus
Napaea hermaphrodita	Phlox divaricata
Nepeta virginica	Phlox glaberrima
Nicotiana rustica	Phlox maculata
Nymphaea alba	Phlox ovata
Nymphaea lotus	Phlox paniculata
Nymphaea lutea	Phlox pilosa
Nymphaea nelumbo	Phlox setacea
Nyssa aquatica	Phlox subulata
	Phryma leptostachya
Obolaria virginica	Physalis pruinosa
Oenothera biennis	Physalis pubescens
Oenothera fruticosa	Physalis viscosa
Oenothera mollissima	Phytolacca americana
Oldenlandia uniflora	Pinus balsamea
Oncoclea sensibilis	Pinus strobus
Ophiorrhiza mitreola	Pinus taeda
Ophrys cernua	Pistacia simaruba
Ophrys lilifolia	Plantago virginica
Orchis ciliaris	Platanus occidentalis
Orchis flava	Poa capillaris
Orchis psycodes	Poa compressa
Orchis spectabilis	Poa flava
Origanum vulgare	Podophyllum diphyllum
Ornithogalum bivale	Podophyllum peltatum
Ornithogalum canadense	Polemonium caeruleum
Ornithogalum hirsutum	Polemonium dubium
Orobanche uniflora	Polemonium rubrum
Orobanche virginiana	Polygala cruciata
Orontium aquaticum	Polygala incarnata
Osmunda cinnamomea	Polygala lutea
Osmunda claytoniana	Polygala sanguinea
Osmunda regalis	Polygala senega
Osmunda virginiana	Polygala verticillata
Osteospermum uvedalia	Polygala viridescens
Othonna cineraria	Polygonum arifolium
Oxalis longiflora	Polygonum articulatum
Oxalis stricta	Polygonum aviculare
Oxalis violacea	Polygonum convolvulus
	Polygonum erectum
Panax quinquefolius	Polygonum maritimum
Panax trifolius	Polygonum pensylvanicum
Pancreatum carolinianum	Polygonum persicaria
Panicum capillare	Polygonum sagittatum
Panicum clandestinum	Polygonum scandens
Panicum crusgalli	Polygonum virginianum
Panicum dichotomum	Polypodium bulbiferum
Panicum dissectum	Polypodium lonchitis
Panicum filiforme	Polypodium marginale
Panicum glaucum	Polypodium noveboracense
Panicum italicum	Polypodium phegopteris
Panicum latifolium	Polygonum virginianum
Panicum sanguinale	Polypremum procumbens
Panicum virgatum	Populus balsamifera
Parthenium integrifolium	Populus heterophylla
Passiflora incarnata	Populus nigra
Passiflora lutea	Polymnia canadensis
Penthorum sedoides	Pontederia cordata
Phalaris oryzoides	Portulaca oleracea
Phaseolus helvulus	Potamogeton nutans

Potentilla canadensis	Sagina virginica
Potentilla norvegica	Sagittaria sagittifolia
Potentilla reptans	Salicornia virginia
Prenanthes alba	Salsola kali
Prenanthes altissima	Salsola prostrata
Prinos glaber	Salvia lyrata
Prinos verticillatus	Salvia urticifolia
Proserpinaca palustris	Sambucus canadensis
Prunus lusitanica	Samolus valerandii
Prunus virginiana	Sanguinaria canadensis
Ptelea trifoliata	Sanguisorba canadensis
Pteris atropurpurea	Sanicula canadensis
Pulmonaria virginica	Sanicula europaea
Pyrola maculata	Sanicula marilandica
Pyrola rotundifolia	Saponaria officinalis
Pyrola umbellata	Sarothra gentianoides
Pyrus coronaria	Sarracenia flava
	Sarracenia purpurea
Quercus alba	Satureja origanoides
Quercus nigra	Satureja virginiana
Quercus phellos	Saururus cernuus
Quercus prinus	Saxifraga nivalis
Quercus rubra	Saxifraga pensylvanica
Queria canadensis	Scandix cerefolium
	Scandix procumbens
Ranunculus abortivus	Schwalbea americana
Ranunculus repens	Schoenus glomeratus
Renealmia usneoides	Scirpus capitatus
Rhexia mariana	Scirpus glomeratus
Rhexia virginica	Scirpus retrofractus
Rhinanthus virginica	Scirpus spadiceus
Rhododendron maximum	Scrophularia marilandica
Rhus copallinum	Scrophularia nodosa
Rhus glabra	Scutellaria hyssopifolia
Rhus radicans	Scutellaria integrifolia
Rhus toxicodendron	Scutellaria lateriflora
Rhus vernix	Senecio aureus
Ribes cynosbati	Senecio canadensis
Ribes nigrum	Senecio hieracifolius
Ribes oxyacanthoides	Serratula glauca
Robinia pseudoacacia	Serratula noveboracensis
Rosa carolina	Serratula praealta
Rubus caesius	Serratula scariosa
Rubus canadensis	Serratula spicata
Rubus fruticosus	Serratula squarrosa
Rubus hispidus	Sicyos angulata
Rubus odoratus	Sida abutilon
Rubus occidentalis	Sida crispa
Rudbeckia hirta	Sida rhombifolia
Rudbeckia laciniata	Sida spinosa
Rudbeckia oppositifolia	Sigesbeckia occidentalis
Rudbeckia purpurea	Silene antirrhina
Rudbeckia triloba	Silene nocturna
Ruellia biflora	Silene virginica
Ruellia strepens	Silphium asteriscus
Rumex acetosella	Silphium helianthoides
Rumex britannica	Silphium laciniatum
Rumex persicarioides	Silphium solidaginoides
Rumex sanguineus	Silphium trifoliatum
Rumex verticillatus	Sison canadense

Sisymbrium nasturtium-aquaticum	Thalictrum purpurascens
Sisyrinchium bermudiana	Thaspia trifoliata
Sium rigidius	Thesium umbellatum
Sloanea emarginata	Thuja occidentalis
Smilax bona-nox	Tiarella cordifolia
Smilax caduca	Tilia americana
Smilax herbacea	Tradescantia virginiana
Smilax lanceolata	Tragopogon virginicum
Smilax laurifolia	Trichostema brachiatum
Smilax pseudo-china	Trichostema dichotomum
Smilax rotundifolia	Trifolium arvense
Smilax sarsaparilla	Trifolium biflorum
Smilax tannoides	Trifolium comosum
Smyrniium aureum	Trifolium reflexum
Smyrniium integerrimum	Trifolium repens
Solanum carolinense	Trillium cernuum
Solanum diphyllum	Trillium erectum
Solanum mammosum	Trillium sessile
Solanum nigrum	Triosteum angustifolium
var. virginicum	Triosteum perfoliatum
Solanum nigrum	
var. vulgare	Ulmus americana
Solanum tomentosum	Uniola paniculata
Solanum verbascifolium	Uniola spicata
Solanum virginianum	Urtica canadensis
Solidago altissima	Urtica capitata
Solidago caesia	Urtica cylindrica
Solidago canadensis	Urtica divaricata
Solidago flexicaulis	Urtica pumila
Solidago lateriflora	Utricularia gibba
Solidago latifolia	Utricularia subulata
Solidago noveboracensis	Uvularia perfoliata
Solidago rigida	Uvularia sessilifolia
Solidago sempervirens	
Sonchus canadensis	Vaccinium album
Sonchus floridanus	Vaccinium corymbosum
Sophora tinctoria	Vaccinium frondosum
Sparganium erectum	Vaccinium hispidulum
Spermacoe tenuior	Vaccinium ligustrinum
Spiraea aruncus	Vaccinium mucronatum
Spiraea hypericifolia	Vaccinium stamineum
Spiraea opulifolia	Valeriana cornucopiae
Spiraea tomentosa	Valeriana locusta var. radiata
Spiraea trifoliata	Veratrum luteum
Staphylea trifolia	Verbena hastata
Statice armeria	Verbena nodiflora
Statice limonium	Verbena spuria
Stewartia malacodendron	Verbena urticifolia
Stipa avenacea	Verbesina alba
Swertia corniculata	Verbesina virginica
Swertia difformis	Veronica anagallis-aquatica
	Veronica arvensis
Taxus baccata	Veronica beccabunga
Tetracera volubilis	Veronica marilandica
Tetragonotheca helianthoides	Veronica serpyllifolia
Teucrium canadense	Veronica virginica
Teucrium chamaepitys	Viburnum acerifolium
Teucrium virginicum	Viburnum dentatum
Thalictrum cornutii	Viburnum lentago
Thalictrum dioicum	Viburnum nudum

Viburnum prunifolium
Viola canadensis
Viola lanceolata
Viola palmata
Viola pedata
Viola primulifolia
Viscum purpureum
Viscum rubrum
Viscum terrestre
Vitex agnus-castus
Vitis arborea
Vitis labrusca
Vitis laciniosa
Vitis vinifera
Vitis vulpina

Xanthium strumarium
Xyris indica

Yucca filamentosa
Yucca gloriosa

Zannichellia palustris
Zanthoxylum clava-herculis
Zizania aquatica

Vascular Plants from Temperate North America Mentioned in Linnaeus' First Edition of Species Plantarum as Occurring only in Temperate North America

America

Actaea spicata var. *alba*
Aletris farinosa
Alyssum hyperboreum
Andropogon alopecuroides
Andropogon virginicum
Annona muricata
Arabis canadensis
Arethusa divaricata
Aristolochia arborescens
Arum dracontium
Asclepias amoena
Asclepias tuberosa
Asclepias variagata
Aster dumosus
Aster ericoides
Aster grandiflorus
Aster laevis
Aster linariifolius
Aster linifolius
Aster miser
Aster mutabilis
Aster puniceus
Aster tenuifolius
Aster undulatus
Bidens bullata
Bidens frondosa
Bidens pilosa

Bignonia capreolata
Bignonia radicans
Bupthalmum helianthoides
Cacalia porophyllum
Cactus pentagonus
Campanula americana
Celosia paniculata
Cephalanthus occidentalis
Chionanthus virginica
Chrysanthemum serotinum
Circaea lutetiana
 var. *canadensis*
Clitoria mariana
Coix dactyloides
Commelina communis
Convolvulus repens
Conyza asteroides
Cynanchum hirtum
Cyperus odoratus
Diospyros virginiana
Dracocephalum virginianum
Eriocaulon decangulare
Eupatorium purpureum
Euphorbia maculata
Fagus pumila
Galium tinctorium
Gomphrena interrupta
Gomphrena serrata
Hedysarum volubile
Helenium autumnale
Helianthus divaricatus
Hydrocotyle americana
Hydrocotyle umbellata
Jussiaea erecta
Limodorum tuberosum
Liriodendron tulipifera
Mentzelia aspera
Mitella diphylla
Myrica asplenifolia
Nyssa aquatica
Osmunda virginiana
Panicum filiforme
Panicum latifolium
Phlox paniculata
Phryma leptostachya
Physalis pruinosa
Platanus occidentalis
Podophyllum peltatum
Polygonum canadense
Populus balsamifera
Pyrola maculata
Quercus nigra
Quercus phellos
Quercus prinus
Rhus copallinum
Rhus glabra
Sanguinaria canadensis
Sarracenia flava
Sarracenia purpurea
Schwalbea americana

Senecio hieracifolius	Cistus canadensis
Serratula spicata	Collinsonia canadensis
Silphium laciniatum	Convallaria racemosa
Sison canadense	Convallaria stellata
Smyrniun aureum	Conyza bifrons
Solanum diphyllum	var. flosculosa
Solanum tomentosum	Conyza linifolia
Solanum verbascifolium	Coreopsis alternifolia
Solanum virginianum	Cornus canadensis
Solidago altissima	Cracca virginiana
Solidago caesia	Crataegus coccinea
Solidago lateriflora	Cucubalus stellata
Solidago noveboracensis	Cupressus thuyoides
Trichostema brachiatum	Dalibarda repens
Trifolium comosum	Elymus canadensis
Triosteum perfoliatum	Epigaea repens
Uniola spicata	Erigeron philadelphicus
Vaccinium corymbosum	Eupatorium rotundifolium
Vaccinium frondosum	Euphorbia corollata
Vaccinium mucronatum	Euphorbia ipecacuanhae
Vaccinium stamineum	Euphorbia polygonifolia
Vitis labrusca	Fumaria cucullaria
	Fumaria sempervirens
Canada	Galium trifidum
Actaea racemosa	Gaultheria procumbens
Adiantum pedatum	Gerardia flava
Ageratum altissimum	Gerardia pedicularia
Allium canadense	Gerardia purpurea
Ambrosia elatior	Guilandina dioica
Ambrosia trifida	Hedera quinquefolia
Anemone quinquefolia	Hedysarum canadense
Anemone thalictroides	Helianthus decapetalus
Angelica atropurpurea	Helianthus giganteus
Angelica lucida	Helianthus strumosus
Antirrhinum canadense	Hibiscus moscheutos
Apocynum androsaemifolium	Hibiscus palustris
Apocynum cannabinum	Hieracium paniculatum
Aquilegia canadensis	Hippophae canadensis
Arabis lyrata	Holcus laxus
Aralia racemosa	Hordeum jubatum
Arethusa bulbosa	Hypericum canadense
Arethusa ophioglossoides	Hypericum mutilum
Asarum canadense	Hyssopus nepetoides
Asclepias incarnata	Lactuca canadensis
Aster annuus	Lechea major
Astragalus canadensis	Lechea minor
Betula lenta	Lilium canadense
Betula nigra	Liquidambar peregrina
Bromus ciliatus	Lobelia cliffortiana
Bromus purgans	Lobelia inflata
Buchnera americana	Lobelia kalmii
Cacalia atriplicifolia	Lycopodium alopecuroides
Carex folliculata	Lysimachia ciliata
Carex squarrosa	Melissa pulegioides
Celastrus scandens	Menispermum canadense
Chelone glabra	Mentha canadensis
Chironia campanulata	Mespilus canadensis
Chrysocoma graminifolia	Mimulus ringens
Cicuta bulbifera	Monarda fistulosa
Cinna arundinacea	Monotropa uniflora

Ophrys cernua	Veratrum luteum
Orchis ciliaris	Verbena hastata
Orchis psychodes	Verbena spuria
Ornithogalum hirsutum	Verbena urticifolia
Orontium aquaticum	Viburnum lentago
Othonna cineraria	Viburnum prunifolium
Oxalis violacea	Viola canadensis
Panax quinquefolius	
Pinus balsamea	Carolina
Pinus strobus	Amorpha fruticosa
Pinus taeda	Andromeda arborea
Poa capillaris	Annona glabra
Polygala incarnata	Annona triloba
Polygonum articulatum	Asarum virginicum
Polypodium bulbiferum	Asclepias purpurascens
Polypodium marginale	Astragalus carolinianus
Polypodium noveboracense	Bidens nivea
Polymnia canadensis	Bignonia caerulea
Potentilla canadensis	Callicarpa americana
Prenanthes altissima	Canna glauca
Prinos glaber	Carduus altissimus
Queria canadensis	Ceanothus americanus
Ranunculus abortivus	Cissampelos smilacina
Rhus radicans	Clematis crispa
Rhus toxicodendron	Clematis viorna
Ribes cynosbati	Clethra alnifolia
Ribes oxycanthoides	Convolvulus carolinus
Rubus canadensis	Coreopsis lanceolata
Rubus hispidus	Crotalaria alba
Rubus odoratus	Crotalaria perfoliata
Rubus occidentalis	Cupressus distichia
Rudbeckia hirta	Erythrina herbacea
Rudbeckia laciniata	Eupatorium coelestinum
Sanguisorba canadensis	Fraxinus americana
Saxifraga pensylvanica	Geranium carolinianum
Scutellaria integrifolia	Glycine frutescens
Scutellaria lateriflora	Gnaphalium purpureum
Senecio aureus	Hedysarum marilandicum
Senecio canadensis	Horminum virginicum
Smilax caduca	Ilex cassine
Smilax rotundifolia	Ipomoea carolina
Solidago canadensis	Ipomoea lacunosa
Solidago flexicaulis	Ipomoea tannifolia
Solidago latifolia	Juniperus virginiana
Solidago sempervirens	Laurus borbonia
Sonchus canadensis	Laurus sassafras
Sonchus floridanus	Lonicera marilandica
Spiraea trifoliata	Lonicera symphoricarpos
Teucrium canadense	Lycopodium apodum
Thalictrum cornutii	Lycopodium carolinianum
Thalictrum dioicum	Magnolia virginiana
Thalictrum purpurascens	var. acuminata
Tilia americana	var. foetida
Tragopogon virginicum	var. glauca
Trifolium biflorum	var. grisea
Urtica capitata	var. tripetala
Urtica divaricata	Malva caroliniana
Urtica pumila	Menispermum carolinum
Uvularia perfoliata	Menispermum virginicum
Uvularia sessilifolia	Mitchella repens

Myrica cerifera
 Phaseolus helvulus
 Philadelphus inodorus
 Polemonium rubrum
 Polyprenum procumbens
 Prenanthes alba
 Quercus rubra
 Rosa carolina
 Rudbeckia purpurea
 Ruellia biflora
 Ruellia strepens
 Serratula glauca
 Serratula praealta
 Silene antirrhina
 Silphium asteriscus
 Sloanea emarginata
 Smilax bona-nox
 Smilax laurifolia
 Smilax tannoides
 Solanum carolinense
 Spermacoe tenuior
 Spiraea hypericifolia
 Spiraea opulifolia
 Trillium cernuum
 Trillium sessile
 Uniola paniculata
 Viscum purpureum
 Viscum rubrum
 Vitis arborea

Florida

Actaea racemosa
 Dioscorea villosa
 Laurus sassafras
 Polygonum arifolium

Maryland

Acrostichum areolatum
 Asarum virginicum
 Cassia marilandica
 Iris versicolor
 Juglans nigra
 Kalmia latifolia
 Lonicera marilandica
 Mitchellia repens
 Monotropa uniflora
 Osmunda cinnamomea
 Polygala senega
 Polygonum sagittatum
 Rhexia mariana
 Sanicula marilandica
 Saururus cernuus
 Serratula glauca
 Smilax herbacea
 Valeriana locusta
 var. radiata

Mississippi

Erythrina herbacea
 Silphium laciniatum

New England

Aster novae-angliae

New Jersey

Kalmia angustifolia

New York

Bartsia coccinea
 Holosteum succulentum
 Kalmia angustifolia
 Monarda didyma
 Solidago sempervirens

Pennsylvania

Acer pensylvanicum
 Acer rubrum
 Acer saccharinum
 Amaranthus retroflexus
 Ambrosia artemisiifolia
 Andromeda racemosa
 Aster novi-belgii
 Avena pensylvanica
 Clethra alnifolia
 Datisca hirta
 Eupatorium altissimum
 Euphorbia portulacoides
 Gaura biennis
 Gentiana quinquefolia
 Gnaphalium obtusifolium
 Gnaphalium purpureum
 Helonias bullata
 Hieracium gronovii
 Hieracium kalmii
 Iris versicolor
 Kalmia angustifolia
 Kalmia latifolia
 Linum virginianum
 Lycopodium apodum
 Lycopodium obscurum
 Monarda didyma
 Myrica cerifera
 Panax quinquefolius
 Polygala senega
 Polygonum erectum
 Polygonum pensylvanicum
 Prenanthes alba
 Sarothra gentianoides
 Saxifraga pensylvanica
 Serratula praealta
 Smilax tannoides
 Solidago rigida
 Spiraea tomentosa
 Thesium umbellatum
 Trichostema dichotomum
 Vaccinium album
 Vaccinium ligustrinum
 Viscum terrestre

Virginia

Acer negundo

- Acer rubrum
Acnida cannabina
Acrostichum areolatum
Acrostichum platyneuros
Actaea racemosa
Adiantum pedatum
Agave virginica
Ageratum altissimum
Agrostis virginica
Alisma cordifolia
Alisma subulata
Amaranthus graecizans
Amaranthus hybridus
Amaranthus hypocondriacus
Amaranthus lividus
Amaryllis atamasca
Ambrosia artemisiifolia
Ambrosia elatior
Ambrosia trifida
Ammannia ramosior
Anchusa virginiana
Andromeda arborea
Andromeda mariana
Andromeda paniculata
Andropogon divaricatum
Anemone quinquefolia
Anemone thalictroides
Anemone virginiana
Antirrhinum canadense
Apocynum androsaemifolium
Apocynum cannabinum
Aquilegia canadensis
Aralia spinosa
Arethusa bulbosa
Arethusa ophioglossoides
Aristolochia serpentaria
Arum virginicum
Asarum virginicum
Asclepias decumbens
Asclepias incarnata
Asclepias rubra
Asclepias syriaca
Asclepias verticillata
Ascyrum crux-andreae
Ascyrum hypericoides
Ascyrum villosum
Aster concolor
Aster divaricatus
Aster novi-belgii
Aster rigidus
Aster tradescantii
Aster vernus
Astragalus canadensis
Azalea lutea
Azalea viscosa
Baccharis foetida
Baccharis halimifolia
Bartsia coccinea
Betula lenta
Betula nigra
Bidens bipinnata
Bignonia sempervirens
Buchnera americana
Burmannia biflora
Cacalia atriplicifolia
Cacalia suaveolens
Callicarpa americana
Campanula perfoliata
Cardamine virginica
Carduus virginianus
Cassia marilandica
Cassia nictitans
Ceanothus americanus
Celastrus bullatus
Celtis occidentalis
Cenchrus tribuloides
Cercis canadensis
Chaerophyllum arborescens
Chelone glabra
Chelone hirsuta
Chelone penstemon
Chenopodium virginicum
Chironia angularis
Chironia dodecandra
Chrysogonum virginianum
Cicuta bulbifera
Cicuta maculata
Claytonia virginica
Clematis vionna
Clethra alnifolia
Clitoria virginiana
Collinsonia canadensis
Commelina erecta
Convallaria racemosa
Convolvulus panduratus
Convolvulus spithameus
Coreopsis alternifolia
Coreopsis angustifolia
Coreopsis auriculata
Coreopsis tripteris
Coreopsis verticillata
Cornus florida
Cracca virginiana
Crataegus coccinea
Crataegus tomentosa
Crataegus viridis
Cucubalus stellatus
Cupressus distichia
Cuscuta americana
Cynoglossum virginianum
Cyperus arundinacea
Dianthera americana
Diodia virginiana
Dioscorea villosa
Dirca palustris
Dodecatheon meadia
Dolichos polystachyus
Dolichos regularis
Dracontium foetidum
Elephantopus tomentosus

<i>Elymus virginicus</i>	<i>Hibiscus virginicus</i>
<i>Epigaea repens</i>	<i>Hieracium gronovii</i>
<i>Erigeron camphoratum</i>	<i>Hieracium venosum</i>
<i>Eriophorum virginicum</i>	<i>Holcus laxus</i>
<i>Eryngium aquaticum</i>	<i>Holcus striatus</i>
<i>Eupatorium aromaticum</i>	<i>Horminum virginicum</i>
<i>Eupatorium coelestinum</i>	<i>Houstonia caerulea</i>
<i>Eupatorium hyssopifolium</i>	<i>Houstonia purpurea</i>
<i>Eupatorium perfoliatum</i>	<i>Hydrangea arborescens</i>
<i>Eupatorium rotundifolium</i>	<i>Hydrophyllum virginianum</i>
<i>Eupatorium scandens</i>	<i>Hyoseris virginica</i>
<i>Eupatorium sessilifolium</i>	<i>Hypericum kalmianum</i>
<i>Eupatorium trifoliatum</i>	<i>Hypericum mutilum</i>
<i>Euphorbia corollata</i>	<i>Hypericum setosum</i>
<i>Euphorbia ipecacuanhae</i>	<i>Hyssopus nepetoides</i>
<i>Euphorbia polygonifolia</i>	<i>Ipomoea nyctelea</i>
<i>Euonymus americanus</i>	<i>Iris verna</i>
<i>Ferula canadensis</i>	<i>Iris versicolor</i>
<i>Fraxinus americana</i>	<i>Iris virginica</i>
<i>Fumaria cucullaria</i>	<i>Itea virginica</i>
<i>Fumaria sempervirens</i>	<i>Juglans alba</i>
<i>Galax aphylla</i>	<i>Juglans nigra</i>
<i>Galium bermudense</i>	<i>Juniperus virginiana</i>
<i>Gaura biennis</i>	<i>Jussiaea erecta</i>
<i>Gentiana saponaria</i>	<i>Kalmia latifolia</i>
<i>Gentiana villosa</i>	<i>Laurus aestivalis</i>
<i>Geranium carolinianum</i>	<i>Laurus benzoin</i>
<i>Gerardia flava</i>	<i>Laurus borbonica</i>
<i>Gerardia pedicularia</i>	<i>Laurus indica</i>
<i>Gerardia purpurea</i>	<i>Laurus sassafra</i>
<i>Geum virginianum</i>	<i>Leontice thalictroides</i>
<i>Gleditsia triacanthos</i>	<i>Leontodon dandelion</i>
<i>Glycine apios</i>	<i>Linum virginianum</i>
<i>Glycine bracteata</i>	<i>Lithospermum virginianum</i>
<i>Glycine tomosa</i>	<i>Lobelia cardinalis</i>
<i>Glycine tomentosa</i>	<i>Lobelia cliffortiana</i>
<i>Gnaphalium obtusifolium</i>	<i>Lobelia inflata</i>
<i>Gnaphalium plantaginifolium</i>	<i>Lobelia siphilitica</i>
<i>Gnaphalium purpureum</i>	<i>Lonicera marilandica</i>
<i>Gratiola dubia</i>	<i>Lonicera symphoricarpos</i>
<i>Gratiola virginiana</i>	<i>Ludwigia alternifolia</i>
<i>Hamamelis virginiana</i>	<i>Lupinus perennis</i>
<i>Hedysarum canadense</i>	<i>Lycopodium alopecuroides</i>
<i>Hedysarum frutescens</i>	<i>Lycopodium apodum</i>
<i>Hedysarum hirtum</i>	<i>Lycopsis virginica</i>
<i>Hedysarum marilandicum</i>	<i>Lycopus virginicus</i>
<i>Hedysarum nudiflorum</i>	<i>Lysimachia ciliata</i>
<i>Hedysarum paniculatum</i>	<i>Lysimachia quadrifolia</i>
<i>Hedysarum repens</i>	<i>Lythrum lineare</i>
<i>Hedysarum violaceum</i>	<i>Lythrum petiolatum</i>
<i>Hedysarum virginicum</i>	<i>Lythrum verticillatum</i>
<i>Hedysarum viridiflorum</i>	<i>Magnolia virginiana</i>
<i>Helianthus angustifolius</i>	var. <i>acuminata</i>
<i>Helianthus atrorubens</i>	var. <i>foetida</i>
<i>Helianthus giganteus</i>	var. <i>glauca</i>
<i>Helianthus laevis</i>	var. <i>grisea</i>
<i>Helianthus multiflorus</i>	var. <i>tripetala</i>
<i>Heuchera americana</i>	<i>Medeola virginiana</i>
<i>Hibiscus moscheutos</i>	<i>Medicago virginica</i>
<i>Hibiscus palustris</i>	<i>Melanthium virginicum</i>

Melissa pulegioides	Polygala sanguinea
Menispermum canadense	Polygala senega
Menispermum virginicum	Polygala verticillata
Mespilus arbutifolia	Polygala viridescens
Mespilus canadensis	Polygonum arifolium
Mimulus ringens	Polygonum sagittatum
Mitchella repens	Polygonum virginianum
Monarda ciliata	Polypodium virginianum
Monarda clinopodia	Polypremum procumbens
Monarda punctata	Populus heterophylla
Monotropa uniflora	Pontederia cordata
Morus rubra	Prenanthes alba
Myosotis virginiana	Prenanthes altissima
Myrica cerifera	Prinos verticillatus
Napaea dioica	Proserpinaca palustris
Napaea hermaphrodita	Prunus virginiana
Napaea virginica	Ptelea trifoliata
Obalaria virginica	Pteris atropurpurea
Oenothera fruticosa	Pulmonaria virginica
Oldenlandia uniflora	Pyrus coronaria
Onoclea sensibilis	Quercus alba
Ophrys cernua	Quercus rubra
Orchis ciliaris	Queria canadensis
Orchis flava	Ranunculus abortivus
Orchis spectabilis	Rhexia virginica
Ornithogalum bivale	Rhinanthus virginica
Ornithogalum hirsutum	Rhododendron maximum
Orobanche uniflora	Rhus radicans
Orobanche virginiana	Rhus toxicodendron
Orontium aquaticum	Robinia pseudoacacia
Osmunda claytoniana	Rudbeckia hirta
Osteospermum uvedalia	Rudbeckia laciniata
Oxalis longiflora	Rudbeckia oppositifolia
Oxalis stricta	Rudbeckia purpurea
Oxalis violacea	Rudbeckia triloba
Panax quinquefolius	Ruellia strepens
Panax trifolius	Rumex britannica
Panicum dichotomum	Rumex persicarioides
Panicum virgatum	Rumex sanguineus
Parthenium integrifolium	Rumex verticillatus
Penthorum sedoides	Sagina virginica
Phalaris oryzoides	Salvia lyrata
Phlox divaricata	Salvia urticifolia
Phlox glaberrima	Sambucus canadensis
Phlox maculata	Sanicula canadensis
Phlox ovata	Sanicula marilandica
Phlox pilosa	Sarothra gentianoides
Phlox setacea	Satureja origanoides
Phlox subulata	Satureja virginiana
Pinus balsamea	Saururus cernuus
Pinus strobus	Saxifraga pennsylvanica
Pinus taeda	Scandix procumbens
Plantago virginica	Schoenus glomeratus
Poa capillaris	Scirpus capitatus
Poa flava	Scirpus retrofractus
Podophyllum diphyllum	Scrophularia marilandica
Polemonium dubium	Scutellaria hyssopifolia
Polygala cruciata	Scutellaria integrifolia
Polygala incarnata	Scutellaria lateriflora
Polygala lutea	Senecio aureus

Serratula glauca
 Serratula praealta
 Serratula scariosa
 Serratula squarrosa
 Sigesbeckia occidentalis
 Silene antirrhina
 Silene virginica
 Silphium asteriscus
 Silphium helianthoides
 Silphium solidaginoides
 Silphium trifoliatum
 Sium rigidius
 Smilax herbacea
 Smilax lanceolata
 Smilax laurifolia
 Smilax tamnoides
 Smyrnum integerrimum
 Solidago canadensis
 Sonchus floridanus
 Spiraea opulifolia
 Spiraea trifoliata
 Staphylea trifolia
 Stewartia malacodendron
 Stipa avenacea
 Swertia difformis
 Tetragonotheca helianthoides
 Teucrium virginicum
 Thaspia trifoliata
 Thesium umbellatum
 Tilia americana
 Tradescantia virginiana
 Tragopogon virginicum
 Trichostema dichotomum
 Trifolium biflorum
 Trifolium reflexum
 Trillium erectum
 Trillium sessile
 Triosteum angustifolium
 Ulmus americana
 Urtica divaricata
 Utricularia gibba
 Utricularia subulata
 Uvularia perfoliata
 Vaccinium hispidulum
 Veratrum luteum
 Verbena nodiflora
 Verbena spuria
 Verbena urticifolia
 Verbesina virginica
 Veronica marilandica
 Veronica virginica
 Viburnum acerifolium
 Viburnum dentatum
 Viburnum nudum
 Viburnum prunifolium
 Viola palmata
 Viola pedata
 Vitis arborea
 Vitis vulpina
 Yucca filamentosa

**Vascular Plants from Temperate
 North America Mentioned in
 Linnaeus' First Edition of
 Species Plantarum as Occurring
 only in the New World**

Acrostichum polypodioides
 Aesculus pavia
 Andropogon nutans
 Annona muricata
 Arum triphyllum
 Asclepias nivea
 Baccharis ivaefolia
 Bignonia crucigera
 Buphthalmum frutescens
 Caesalpinia brasiliensis
 Cassia chamaecrista
 Cassia ligustrina
 Celastrus myrtifolius
 Chenopodium anthelminticum
 Crescentia cujete
 Crotalaria sagittalis
 Cynanchum suberosum
 Cyperus strigosus
 Eryngium foetidum
 Hedysarum canescens
 Hypericum lasianthus
 Iva frutescens
 Laurus winterana
 Lepidium virginicum
 Liquidambar styraciflua
 Lonicera sempervirens
 Melothria pendula
 Pancratium carolinianum
 Panicum capillare
 Panicum clandestinum
 Passiflora incarnata
 Passiflora lutea
 Physalis viscosa
 Phytolacca americana
 Pistacia simaruba
 Renealmia usneoides
 Sicyos angulata
 Sida crispa
 Sisyrrinchium bermudiana
 Smilax pseudo-china
 Smilax sarsaparilla
 Solanum mammosum
 Sophora tinctoria
 Tetracera volubilis
 Urtica cylindrica
 Verbesina alba
 Yucca gloriosa

**Vascular Plants from Temperate
 North America Mentioned in
 Linnaeus' First Edition of
 Species Plantarum as Occurring
 in Temperate North America and
 Elsewhere in the World**

- Acalypha virginica*
Andromeda calyculata
Anemone dichotoma
Aralia nudicaulis
Arbutus uva-ursi
Arnica maritima
Asplenium rhizophyllum
Aster cordifolius
Atriplex laciniata
Bunias cakile
Cactus opuntia
Carpinus betulus
Carpinus ostrya
Cassine peragua
Chelidonium glaucium
Chrysanthemum arcticum
Chrysoplenium oppositifolium
Clematis vitalba
Clinopodium rugosum
Clinopodium vulgare
Convolvulus hederaceus
Cornus sanguinea
Cynosurus aegyptius
Cypripedium calceolus
Dactylis cynosuroides
Datura stramonium
Dianthus plumarius
Drosera rotundifolia
Erigeron canadense
Erythronium dens-canis
Gentiana ciliata
Geranium maculatum
Gnaphalium margaritaceum
Helleborus trifolius
Hypericum ascyron
Ilex aquifolium
Impatiens noli-tangere
Isnardia palustris
Juncus bulbosus
Lilium camschatcense
Linnaea borealis
Lycopodium complanatum
Lycopodium rupestre
Melica altissima
Mollugo verticillata
Monotropa hypopithys
Nicotiana rustica
Nymphaea alba
Nymphaea lotus
Oenothera biennis
Ophrys lilifolia
Origanum vulgare
Panicum crusgalli
Panicum sanguinale
Poa compressa
Polemonium caeruleum
Polygonum maritimum
Polypodium lonchitis
Polypodium phegopteris
Portulaca oleracea
Potentilla norvegica
Prunus lusitanica
Pyrola rotundifolia
Pyrola umbellata
Rhus vernix
Ribes nigrum
Sagittaria sagittifolia
Salicornia virginica
Samolus valerandii
Saxifraga nivalis
Scirpus glomeratus
Serratula noveboracensis
Silene nocturna
Sisymbrium nasturtium-aquaticum
Statice armeria
Statice limonium
Swertia corniculata
Taxus baccata
Thuja occidentalis
Tiarella cordifolia
Trifolium arvense
Urtica canadensis
Valeriana cornucopiae
Veronica serphillifolium
Viola primulifolia
Xanthium strumarium
- Vascular Plants from Temperate North America Mentioned in Linnaeus' First Edition of Species Plantarum as Occurring Elsewhere but not Temperate North America**
Agrimonia eupatoria
Alsine media
Amaranthus spinosus
Andropogon hirtum
Anemone hepatica
Angelica sylvestris
Anthericum calyculatum
Antirrhinum elatine
Aphanes arvensis
Arenaria rubra var. marina
Arundo phragmites
Atriplex halimus
Betonica annua
Bignonia catalpa
Briza eragrostis
Callitriche palustris
Caltha palustris
Carex pseudocyperus
Cerastium semidecandrum
Chenopodium album
Clinopodium incanum
Coix lacryma-jobi
Convallaria polygonatum
Daucus carota
Dipsacus fullonum
Draba verna
Elatine hydropiper

Elephantopus scaber
 Epilobium hirsutum
 Equisetum arvense
 Equisetum hyemale
 Heliotropium indicum
 Juncus campestris
 Juncus effusus
 Juncus filiformis
 Lamium amplexicaule
 Ligusticum scothieum
 Lysimachia punctata
 Melissa nepeta
 Mentha spicata var. viridis
 Nymphaea lutea
 Nymphaea nelumbo
 Oenothera mollissima
 Ophiorrhiza mitreola
 Osmunda regalis
 Panicum dissectum
 Panicum glaucum
 Panicum italicum
 Physalis pubescens
 Polygonum aviculare
 Polygonum convolvulus
 Polygonum persicaria
 Populus nigra
 Potamogeton nutans
 Potentilla reptans
 Ranunculus repens
 Rubus caesius
 Rubus fruticosus
 Rumex acetosella
 Salsola kali
 Salsola prostrata
 Sanicula europaea
 Saponaria officinalis
 Scandix cerefolium
 Scirpus spadicus
 Scrophularia nodosa
 Sida abutilon
 Sida rhombifolia
 Sida spinosa
 Solanum nigrum var. vulgare
 Sparganium erectum
 Spiraea aruncus
 Teucrium chamaepitys
 Trifolium repens
 Veronica anagallis-aquatica
 Veronica arvensis
 Veronica beccabunga
 Vitex agnus-castus
 Vitis vinifera
 Xyris indica

**Vascular Plants Named for
 Certain Geographical Areas in
 Temperate North America by
 Linnaeus in the First Edition
 of Species Plantarum
 America**

Buchnera americana
 Callicarpa americana
 Campanula americana
 Ceanothus americanus
 Cuscuta americana
 Dianthera americana
 Euonymus americanus
 Fraxinus americana
 Heuchera americana
 Hydrocotyle americana
 Phytolacca americana
 Schwalbea americana
 Tilia americana
 Ulmus americana

Canada

Allium canadense
 Antirrhinum canadense
 Aquilegia canadensis
 Arabis canadensis
 Asarum canadense
 Astragalus canadensis
 Cercis canadensis
 Circaea lutetiana
 var. canadensis
 Cistus canadensis
 Collinsonia canadensis
 Cornus canadensis
 Elymus canadensis
 Erigeron canadense
 Ferula canadensis
 Hippophae canadensis
 Hypericum canadense
 Lactuca canadensis
 Lilium canadense
 Menispermum canadense
 Mentha canadensis
 Mespilus canadensis
 Ornithogalum canadense
 Polymnia canadensis
 Potentilla canadensis
 Queria canadensis
 Rubus canadensis
 Sambucus canadensis
 Sanguinaria canadensis
 Sanguisorba canadensis
 Sanicula canadensis
 Senecio canadensis
 Sison canadense
 Solidago canadensis
 Sonchus canadensis
 Teucrium canadense
 Urtica canadensis
 Viola canadensis

Carolina

Astragalus carolinianus
 Convolvulus carolinus
 Geranium carolinanum
 Ipomoea carolina

Lycopodium carolinianum
 Malva caroliniana
 Mensiperium carolinum
 Pancratium carolinianum
 Rosa carolina
 Solanum carolinense

Florida

Sonchus floridanus

Maryland

Andromeda mariana
 Cassia marilandica
 Clitoria mariana
 Hedysarum marilandicum
 Lonicera marilandica
 Rhexia mariana
 Sanicula marilandica
 Scrophularia marilandica
 Veronica marilandica

New England

Aster novae-angliae

New York

Aster novi-belgii
 Polypodium noveboracense
 Serratula noveboracensis
 Solidago noveboracensis

Pennsylvania

Acer pensylvanicum
 Avena pensylvanica
 Erigeron philadelphicum
 Polygonum pensylvanicum
 Saxifraga pensylvanica

Virginia

Acalypha virginica
 Agave virginica
 Agrostis virginica
 Anchusa virginiana
 Andropogon virginicum
 Anemone virginiana
 Arum virginicum
 Asarum virginicum
 Cardamine virginica
 Carduus virginianus
 Chenopodium virginicum
 Chionanthus virginica
 Chrysogonum virginianum
 Claytonia virginica
 Clitoria virginiana
 Cracca virginiana
 Cynoglossum virginianum
 Diodia virginiana
 Diospyros virginiana
 Dracocephalum virginianum
 Elymus virginicus
 Eriophorum virginicum

Geum virginianum
 Gratiola virginiana
 Hamamelis virginiana
 Hedysarum virginicum
 Hibiscus virginicus
 Hydrophyllum virginianum
 Hyoseris virginica
 Iris virginica
 Itea virginica
 Juniperus virginiana
 Lepidium virginicum
 Linum virginianum
 Lithospermum virginianum
 Lycopsis virginica
 Lycopus virginicus
 Magnolia virginiana
 Medeola virginiana
 Medicago virginica
 Melanthium virginicum
 Menispermum virginicum
 Myosotis virginiana
 Nepeta virginica
 Obolaria virginica
 Orobanche virginiana
 Osmunda virginiana
 Plantago virginica
 Polygonum virginianum
 Polypodium virginianum
 Prunus virginiana
 Pulmonaria virginica
 Rhexia virginica
 Rhinanthus virginica
 Sagina virginica
 Salicornia virginica
 Satureja virginiana
 Silene virginica
 Solanum virginianum
 Teucrium virginicum
 Tradescantia virginiana
 Tragopogon virginicum
 Verbesina virginica

**Collection Numbers and Species
 of Vascular Plants Gathered by
 John Clayton and Indirectly
 Reported in Linnaeus' First
 Edition of Species Plantarum**

201 Acalypha virginica
 530 Acer negundo
 000 Acer rubrum
 599 Acnida cannabina
 011 Acrostichum areolatum
 014 Acrostichum platyneuros
 685 Acrostichum polyodioides
 305 Actaea racemosa
 320 Adiantum pedatum
 321 Adiantum pedatum
 498 Agave virginica
 199 Ageratum altissimum

000	<i>Agrimonia eupatoria</i>	230	<i>Ascyrum hypericoides</i>
000	<i>Agrostis virginica</i>	607	<i>Aster concolor</i>
507	<i>Agrostis virginica</i>	143	<i>Aster divaricatus</i>
074	<i>Aletris farinosa</i>	767	<i>Aster divaricatus</i>
723	<i>Alisma subulata</i>	072	<i>Aster dumosus</i>
529	<i>Alsine media</i>	194	<i>Aster ericoides</i>
442	<i>Amaranthus graecizans</i>	239	<i>Aster grandiflorus</i>
000	<i>Amaranthus lividus</i>	072	<i>Aster linifolius</i>
569	<i>Amaranthus spinosus</i>	244	<i>Aster novae-angliae</i>
256	<i>Amaryllis atamasca</i>	009	<i>Aster rigidus</i>
512	<i>Ambrosia elatior</i>	391	<i>Aster vernus</i>
724	<i>Ambrosia trifida</i>	565	<i>Astragalus canadensis</i>
774	<i>Ammannia ramosior</i>	571	<i>Atriplex halimus</i>
304	<i>Anchusa virginiana</i>	000	<i>Atriplex laciniata</i>
000	<i>Andromeda arborea</i>	052	<i>Azalea lutea</i>
000	<i>Andromeda calyculata</i>	032	<i>Azalea viscosa</i>
030	<i>Andromeda mariana</i>	159	<i>Baccharis foetida</i>
073	<i>Andromeda paniculata</i>	240	<i>Baccharis halimifolia</i>
601	<i>Andropogon alopecuroides</i>	293	<i>Bartsia coccinea</i>
070	<i>Andropogon divaricatum</i>	271	<i>Betonica annua</i>
600	<i>Andropogon divaricatum</i>	000	<i>Betula lenta</i>
602	<i>Andropogon hirtum</i>	688	<i>Betula nigra</i>
621	<i>Andropogon nutans</i>	100	<i>Bignonia crucigera</i>
460	<i>Andropogon virginicum</i>	225	<i>Bignonia radicans</i>
606	<i>Andropogon virginicum</i>	582	<i>Briza eragrostis</i>
328	<i>Anemone hepatica</i>	142	<i>Buchnera americana</i>
294	<i>Anemone thalictroides</i>	732	<i>Bunias cakile</i>
529	<i>Anemone virginiana</i>	242	<i>Bupththalmum frutescens</i>
125	<i>Angelica sylvestris</i>	208	<i>Bupththalmum helianthoides</i>
058	<i>Annona muricata</i>	248	<i>Burmannia biflora</i>
269	<i>Anthericum calyculatum</i>	133	<i>Cacalia atriplicifolia</i>
256	<i>Antirrhinum canadense</i>	099	<i>Cactus opuntia</i>
435	<i>Antirrhinum elatine</i>	000	<i>Callicarpa americana</i>
374	<i>Aphanes arvensis</i>	378	<i>Callitriche palustris</i>
438	<i>Apocynum cannabinum</i>	522	<i>Caltha palustris</i>
338	<i>Aquilegia canadensis</i>	020	<i>Campanula perfoliata</i>
400	<i>Arabis canadensis</i>	462	<i>Cardamine virginica</i>
745	<i>Arabis canadensis</i>	193	<i>Carduus virginianus</i>
056	<i>Arabis lyrata</i>	259	<i>Carex pseudocyperus</i>
394	<i>Arabis lyrata</i>	000	<i>Carpinus ostrya</i>
042	<i>Aralia nudicaulis</i>	156	<i>Cassia chamaecrista</i>
233	<i>Aralia spinosa</i>	146	<i>Cassia ligustrina</i>
475	<i>Arenaria rubra</i>	069	<i>Ceanothus americanus</i>
	var. <i>marina</i>	311	<i>Ceanothus americanus</i>
000	<i>Arethusa bulbosa</i>	000	<i>Celastrus scandens</i>
472	<i>Arethusa bulbosa</i>	576	<i>Celosia paniculata</i>
635	<i>Arethusa divaricata</i>	624	<i>Celtis occidentalis</i>
077	<i>Arethusa ophioglossoides</i>	206	<i>Cenchrus tribuloides</i>
000	<i>Aristolochia serpentaria</i>	106	<i>Cephalanthus occidentalis</i>
066	<i>Arum triphyllum</i>	342	<i>Cerastium semidecandrum</i>
228	<i>Arum virginicum</i>	000	<i>Cercis canadensis</i>
581	<i>Arundo phragmites</i>	276	<i>Chelidonium glaucium</i>
288	<i>Asarum canadense</i>	010	<i>Chelone glabra</i>
704	<i>Asarum virginicum</i>	039	<i>Chelone hirsuta</i>
083	<i>Asclepias decumbens</i>	000	<i>Chenopodium album</i>
222	<i>Asclepias incarnata</i>	046	<i>Chionanthus virginica</i>
065	<i>Asclepias nivea</i>	120	<i>Chironia dodecandra</i>
263	<i>Asclepias rubra</i>	298	<i>Chrysogonum virginianum</i>
000	<i>Asclepias syriaca</i>	215	<i>Cicuta bulbifera</i>
216	<i>Asclepias verticillata</i>	013	<i>Cicuta maculata</i>

123	<i>Cicuta maculata</i>	439	<i>Eriocaulon decangulare</i>
251	<i>Claytonia virginica</i>	461	<i>Eriophorum virginicum</i>
411	<i>Clematis viorna</i>	500	<i>Eryngium aquaticum</i>
270	<i>Clematis vitalba</i>	282	<i>Eryngium foetidum</i>
112	<i>Clitoria virginiana</i>	691	<i>Erythronium dens-canis</i>
067	<i>Coix lacryma-jobi</i>	603	<i>Eupatorium aromaticum</i>
093	<i>Commelina communis</i>	179	<i>Eupatorium coelestinum</i>
093	<i>Commelina erecta</i>	000	<i>Eupatorium perfoliatum</i>
335	<i>Convallaria polygonatum</i>	162	<i>Eupatorium purpureum</i>
035	<i>Convallaria racemosa</i>	147	<i>Eupatorium scandens</i>
504	<i>Convolvulus hederaceus</i>	620	<i>Eupatorium trifoliatum</i>
641	<i>Convolvulus panduratus</i>	155	<i>Euphorbia corollata</i>
665	<i>Convolvulus repens</i>	000	<i>Euphorbia ipecacuanhae</i>
553	<i>Convolvulus spithameus</i>	152	<i>Euphorbia polygonifolia</i>
667	<i>Coreopsis angustifolia</i>	075	<i>Euonymus americanus</i>
298	<i>Coreopsis auriculata</i>	000	<i>Fagus pumila</i>
308	<i>Coreopsis verticillata</i>	548	<i>Ferula canadensis</i>
057	<i>Cornus florida</i>	000	<i>Fraxinus americana</i>
102	<i>Cracca virginiana</i>	625	<i>Fumaria cucullaria</i>
043	<i>Crataegus coccinea</i>	004	<i>Galax aphylla</i>
000	<i>Crataegus crus-galli</i>	313	<i>Galium bermudense</i>
055	<i>Crataegus tomentosa</i>	010	<i>Gentiana saponaria</i>
526	<i>Crataegus viridis</i>	605	<i>Gentiana villosa</i>
126	<i>Crotalaria sagittalis</i>	372	<i>Geranium carolinianum</i>
245	<i>Cucubalus stellatus</i>	307	<i>Geranium maculatum</i>
384	<i>Cupressus distichia</i>	009	<i>Gerardia flava</i>
215	<i>Cuscuta americana</i>	192	<i>Gerardia pedicularia</i>
001	<i>Cynanchum suberosum</i>	000	<i>Gerardia purpurea</i>
223	<i>Cynanchum suberosum</i>	199	<i>Geum virginianum</i>
257	<i>Cynoglossum virginianum</i>	000	<i>Gleditsia triacanthos</i>
562	<i>Cyperus arundinacea</i>	127	<i>Glycine apios</i>
509	<i>Cyperus odoratus</i>	592	<i>Glycine bracteata</i>
040	<i>Cypripedium calceolus</i>	182	<i>Glycine comosa</i>
577	<i>Dactylis cynosuroides</i>	113	<i>Glycine tomentosa</i>
583	<i>Dactylis cynosuroides</i>	203	<i>Gnaphalium obtusifolium</i>
000	<i>Datura stramonium</i>	287	<i>Gnaphalium plantaginifolia</i>
444	<i>Daucus carota</i>	385	<i>Gnaphalium purpureum</i>
408	<i>Dianthera americana</i>	164	<i>Gratiola dubia</i>
277	<i>Diodia virginiana</i>	594	<i>Gratiola dubia</i>
094	<i>Dioscorea villosa</i>	379	<i>Gratiola virginiana</i>
080	<i>Diospyros virginiana</i>	544	<i>Hamamelis virginiana</i>
632	<i>Diospyros virginiana</i>	673	<i>Hamamelis virginiana</i>
267	<i>Dipsacus fullonum</i>	116	<i>Hedera quinquefolia</i>
000	<i>Dirca palustris</i>	209	<i>Hedysarum canescens</i>
568	<i>Dolichos polystachyus</i>	174	<i>Hedysarum frutescens</i>
121	<i>Dolichos regularis</i>	510	<i>Hedysarum hirtum</i>
525	<i>Draba verna</i>	516	<i>Hedysarum marilandicum</i>
017	<i>Dracontium foetidum</i>	124	<i>Hedysarum nudiflorum</i>
003	<i>Drosera rotundifolia</i>	184	<i>Hedysarum paniculatum</i>
646	<i>Elatine hydropiper</i>	085	<i>Hedysarum repens</i>
655	<i>Elephantopus scaber</i>	103	<i>Hedysarum violaceum</i>
148	<i>Elephantopus tomentosus</i>	564	<i>Hedysarum virginicum</i>
446	<i>Elymus virginicus</i>	614	<i>Hedysarum virginicum</i>
250	<i>Epigaea repens</i>	190	<i>Hedysarum viridiflorum</i>
586	<i>Epilobium hirsutum</i>	202	<i>Helenium autumnale</i>
341	<i>Equisetum arvense</i>	013	<i>Helianthus angustifolius</i>
657	<i>Equisetum hyemale</i>	136	<i>Helianthus atrorubens</i>
165	<i>Erigeron camphoratum</i>	109	<i>Helianthus giganteus</i>
449	<i>Erigeron canadense</i>	195	<i>Helianthus laevis</i>
234	<i>Eriocaulon decangulare</i>	301	<i>Heuchera americana</i>

424	<i>Heuchera americana</i>	185	<i>Lycopus virginicus</i>
122	<i>Hibiscus moscheutos</i>	433	<i>Lysimachia punctata</i>
567	<i>Hibiscus virginicus</i>	419	<i>Lysimachia quadrifolia</i>
447	<i>Hieracium gronovii</i>	505	<i>Lythrum lineare</i>
386	<i>Hieracium venosum</i>	418	<i>Lythrum petiolatum</i>
589	<i>Holcus laxus</i>	214	<i>Lythrum verticillatum</i>
590	<i>Holcus striatus</i>	034	<i>Magnolia virginiana</i>
060	<i>Houstonia caerulea</i>	404	<i>Magnolia virginiana</i> var. <i>acuminata</i>
063	<i>Houstonia purpurea</i>	024	<i>Magnolia virginiana</i> var. <i>foetida</i>
079	<i>Hydrangea arborescens</i>	022	<i>Medeola virginiana</i>
429	<i>Hydrocotyle umbellata</i>	191	<i>Medicago virginica</i>
249	<i>Hydrophyllum virginianum</i>	422	<i>Melanthium virginicum</i>
376	<i>Hyoseris virginica</i>	198	<i>Melissa nepeta</i>
552	<i>Hypericum canadense</i>	514	<i>Melissa pulegioides</i>
232	<i>Hypericum mutilum</i>	134	<i>Melothria pendula</i>
135	<i>Hypericum setosum</i>	546	<i>Menispermum canadense</i>
168	<i>Hyssopus nepetoides</i>	425	<i>Menispermum virginicum</i>
000	<i>Ilex aquifolium</i>	654	<i>Mentha spicata</i> var. <i>viridis</i>
253	<i>Iris verna</i>	060	<i>Mespilus canadensis</i>
259	<i>Iris virginica</i>	295	<i>Mespilus canadensis</i>
480	<i>Itea virginica</i>	130	<i>Mimulus ringens</i>
556	<i>Itea virginica</i>	028	<i>Mitchella repens</i>
243	<i>Iva frutescens</i>	399	<i>Mollugo verticillata</i>
000	<i>Juglans alba</i>	412	<i>Monarda ciliata</i>
000	<i>Juglans nigra</i>	212	<i>Monarda clinopodia</i>
340	<i>Juncus bulbosus</i>	140	<i>Monarda punctata</i>
332	<i>Juncus campestris</i>	245	<i>Monotropa uniflora</i>
393	<i>Juncus effusus</i>	111	<i>Myosotis virginiana</i>
580	<i>Juncus filiformis</i>	684	<i>Myrica asplenifolia</i>
884	<i>Juniperus virginiana</i>	719	<i>Myrica asplenifolia</i>
021	<i>Kalmia angustifolia</i>	000	<i>Myrica cerifera</i>
331	<i>Lamium amplexicaule</i>	437	<i>Nepeta virginica</i>
000	<i>Laurus aestivalis</i>	000	<i>Nyssa aquatica</i>
520	<i>Laurus aestivalis</i>	049	<i>Nyssa aquatica</i>
054	<i>Laurus benzoin</i>	286	<i>Obolaria virginica</i>
000	<i>Laurus borbonia</i>	000	<i>Oenothera biennis</i>
485	<i>Laurus indica</i>	036	<i>Oenothera fruticosa</i>
056	<i>Laurus sassafra</i>	333	<i>Oenothera fruticosa</i>
610	<i>Lechea minor</i>	200	<i>Oenothera mollissima</i>
545	<i>Leontice thalictroides</i>	587	<i>Oldenlandia uniflora</i>
019	<i>Leontodon dandelion</i>	674	<i>Onoclea sensibilis</i>
383	<i>Leontodon dandelion</i>	714	<i>Onoclea sensibilis</i>
000	<i>Lepidium virginicum</i>	187	<i>Ophiorrhiza mitreola</i>
307	<i>Ligusticum scothieum</i>	658	<i>Ophrys lilifolia</i>
076	<i>Limodorum tuberosum</i>	708	<i>Ophrys lilifolia</i>
440	<i>Linum virginianum</i>	560	<i>Orchis ciliaris</i>
000	<i>Liquidambar styraciflua</i>	639	<i>Orchis flava</i>
487	<i>Liquidambar styraciflua</i>	668	<i>Orchis psycodes</i>
531	<i>Liquidambar styraciflua</i>	260	<i>Orchis spectabilis</i>
016	<i>Liriodendron tulipifera</i>	310	<i>Origanum vulgare</i>
647	<i>Lithospermum virginianum</i>	044	<i>Ornithogalum bivale</i>
005	<i>Lobelia cardinalis</i>	000	<i>Ornithogalum hirsutum</i>
196	<i>Lobelia cliffortiana</i>	387	<i>Orobanche uniflora</i>
000	<i>Lonicera marilandica</i>	604	<i>Orobanche virginiana</i>
201	<i>Lonicera symphoricarpos</i>	053	<i>Orontium aquaticum</i>
281	<i>Lonicera symphoricarpos</i>	000	<i>Osmunda claytoniana</i>
137	<i>Ludwigia alternifolia</i>	011	<i>Osmunda regalis</i>
000	<i>Lupinus perennis</i>	008	<i>Osmunda virginiana</i>
027	<i>Lycopodium alopecuroides</i>		
000	<i>Lycopsis virginica</i>		

138	<i>Osteospermum uvedalia</i>	339	<i>Pulmonaria virginica</i>
221	<i>Osteospermum uvedalia</i>	088	<i>Pyrola maculata</i>
741	<i>Oxalis longiflora</i>	000	<i>Pyrus coronaria</i>
474	<i>Oxalis stricta</i>	000	<i>Quercus alba</i>
000	<i>Oxalis violacea</i>	467	<i>Quercus alba</i>
000	<i>Panax quinquefolius</i>	000	<i>Quercus nigra</i>
329	<i>Panax trifolius</i>	000	<i>Quercus phellos</i>
454	<i>Panicum capillare</i>	000	<i>Quercus prinus</i>
458	<i>Panicum dichotomum</i>	000	<i>Quercus rubra</i>
579	<i>Panicum glaucum</i>	316	<i>Queria canadensis</i>
591	<i>Panicum italicum</i>	317	<i>Queria canadensis</i>
457	<i>Panicum sanguinale</i>	701	<i>Ranunculus abortivus</i>
578	<i>Panicum virgatum</i>	473	<i>Ranunculus repens</i>
606	<i>Panicum virgatum</i>	710	<i>Ranunculus repens</i>
263	<i>Parthenium integrifolium</i>	389	<i>Renealmia usneoides</i>
151	<i>Passiflora incarnata</i>	227	<i>Rhexia virginica</i>
118	<i>Passiflora lutea</i>	488	<i>Rhinanthus virginica</i>
158	<i>Penthorum sedoides</i>	728	<i>Rhus copallinum</i>
595	<i>Phalaris oryzoides</i>	492	<i>Rhus glabra</i>
297	<i>Phlox glaberrima</i>	238	<i>Rhus radicans</i>
129	<i>Phryma leptostachya</i>	000	<i>Rhus toxicodendron</i>
128	<i>Physalis viscosa</i>	479	<i>Rhus toxicodendron</i>
671	<i>Phytolacca americana</i>	681	<i>Rhus vernix</i>
547	<i>Pinus balsamea</i>	050	<i>Robinia pseudoacacia</i>
496	<i>Pinus taeda</i>	634	<i>Rubus caesius</i>
343	<i>Plantago virginica</i>	703	<i>Rubus fruticosus</i>
000	<i>Platanus occidentalis</i>	490	<i>Rudbeckia hirta</i>
471	<i>Platanus occidentalis</i>	539	<i>Rudbeckia laciniata</i>
580	<i>Poa capillaris</i>	609	<i>Rudbeckia oppositifolia</i>
581	<i>Poa capillaris</i>	417	<i>Rudbeckia purpurea</i>
273	<i>Poa flava</i>	490	<i>Rudbeckia purpurea</i>
255	<i>Podophyllum peltatum</i>	657	<i>Rudbeckia triloba</i>
249	<i>Polemonium caeruleum</i>	085	<i>Ruellia strepens</i>
556	<i>Polemonium dubium</i>	098	<i>Ruellia strepens</i>
157	<i>Polygala cruciata</i>	494	<i>Rumex acetosella</i>
414	<i>Polygala senega</i>	000	<i>Rumex britannica</i>
563	<i>Polygala verticillata</i>	000	<i>Rumex verticillatus</i>
000	<i>Polygala viridescens</i>	649	<i>Sagina virginica</i>
382	<i>Polygonum aviculare</i>	278	<i>Sagittaria sagittifolia</i>
000	<i>Polygonum convolvulus</i>	527	<i>Salicornia virginica</i>
670	<i>Polygonum persicaria</i>	667	<i>Salicornia virginica</i>
672	<i>Polygonum persicaria</i>	432	<i>Salsola kali</i>
000	<i>Polygonum sagittatum</i>	019	<i>Salvia lyrata</i>
000	<i>Polygonum scandens</i>	391	<i>Salvia lyrata</i>
183	<i>Polygonum virginianum</i>	292	<i>Salvia urticifolia</i>
322	<i>Polypodium lonchitis</i>	314	<i>Samolus valerandii</i>
768	<i>Polypremum procumbens</i>	247	<i>Sanguinaria canadensis</i>
087	<i>Pontederia cordata</i>	000	<i>Sanicula canadensis</i>
532	<i>Populus heterophylla</i>	000	<i>Sanicula europaea</i>
679	<i>Populus nigra</i>	028	<i>Sanicula marilandica</i>
410	<i>Portulaca oleracea</i>	660	<i>Saponaria officinalis</i>
664	<i>Potamogeton nutans</i>	110	<i>Sarothra gentianoides</i>
699	<i>Potentilla reptans</i>	559	<i>Sarracenia flava</i>
015	<i>Prenanthes alba</i>	717	<i>Sarracenia purpurea</i>
284	<i>Prenanthes alba</i>	197	<i>Satureja origanoides</i>
319	<i>Prenanthes alba</i>	141	<i>Satureja virginiana</i>
078	<i>Prinos verticillatus</i>	107	<i>Saururus cernuus</i>
000	<i>Prunus virginiana</i>	525	<i>Saxifraga nivalis</i>
627	<i>Prunus virginiana</i>	304	<i>Saxifraga pensylvanica</i>
682	<i>Pteris atropurpurea</i>	407	<i>Scandix cerefolium</i>

- | | | | |
|-----|---|-----|---|
| 407 | <i>Scandix procumbens</i> | 135 | <i>Teucrium canadense</i> |
| 033 | <i>Schwalbea americana</i> | 443 | <i>Teucrium chamaepitys</i> |
| 585 | <i>Schoenus glomeratus</i> | 117 | <i>Teucrium virginicum</i> |
| 380 | <i>Scirpus capitatus</i> | 291 | <i>Thapsia trifoliata</i> |
| 570 | <i>Scirpus glomeratus</i> | 554 | <i>Tiarella cordifolia</i> |
| 457 | <i>Scirpus retrofractus</i> | 000 | <i>Tilia americana</i> |
| 456 | <i>Scirpus spadicus</i> | 297 | <i>Tradescantia virginiana</i> |
| 220 | <i>Scrophularia marilandica</i> | 309 | <i>Tragopogon virginicum</i> |
| 000 | <i>Scrophularia nodosa</i> | 177 | <i>Trichostema dichotomum</i> |
| 261 | <i>Scutellaria hyssopifolia</i> | 092 | <i>Trifolium biflorum</i> |
| 105 | <i>Scutellaria integrifolia</i> | 289 | <i>Trifolium reflexum</i> |
| 280 | <i>Scutellaria lateriflora</i> | 390 | <i>Trifolium repens</i> |
| 249 | <i>Senecio aureus</i> | 000 | <i>Trillium sessile</i> |
| 286 | <i>Senecio aureus</i> | 626 | <i>Triosteum angustifolium</i> |
| 015 | <i>Serratula glauca</i> | 524 | <i>Ulmus americana</i> |
| 175 | <i>Serratula glauca</i> | 000 | <i>Uniola paniculata</i> |
| 000 | <i>Serratula scariosa</i> | 508 | <i>Urtica cylindrica</i> |
| 651 | <i>Serratula scariosa</i> | 246 | <i>Urtica pumila</i> |
| 237 | <i>Serratula spicata</i> | 515 | <i>Utricularia gibba</i> |
| 014 | <i>Serratula squarrosa</i> | 517 | <i>Utricularia gibba</i> |
| 441 | <i>Sida abutilon</i> | 031 | <i>Utricularia subulata</i> |
| 131 | <i>Sida rhombifolia</i> | 258 | <i>Uvularia perfoliata</i> |
| 511 | <i>Sigesbeckia occidentalis</i> | 537 | <i>Vaccinium frondosum</i> |
| 388 | <i>Silene antirrhina</i> | 042 | <i>Vaccinium stamineum</i> |
| 423 | <i>Silene virginica</i> | 043 | <i>Valeriana locusta</i> |
| 187 | <i>Silphium asteriscus</i> | | var. <i>radiata</i> |
| 610 | <i>Silphium helianthoides</i> | 299 | <i>Veratrum luteum</i> |
| 721 | <i>Sison canadense</i> | 448 | <i>Verbena nodiflora</i> |
| 528 | <i>Sisymbrium nasturtium-</i>
<i>aquaticum</i> | 000 | <i>Verbena spuria</i> |
| 018 | <i>Sisyrinchium bermudiana</i> | 431 | <i>Verbena urticifolia</i> |
| 279 | <i>Sium rigidius</i> | 163 | <i>Verbesina alba</i> |
| 541 | <i>Smilax herbacea</i> | 166 | <i>Verbesina virginica</i> |
| 082 | <i>Smilax lanceolata</i> | 161 | <i>Veronica anagallis-</i>
<i>aquatica</i> |
| 617 | <i>Smilax laurifolia</i> | 368 | <i>Veronica arvensis</i> |
| 082 | <i>Smilax pseudo-china</i> | 161 | <i>Veronica beccabunga</i> |
| 541 | <i>Smilax pseudo-china</i> | 226 | <i>Veronica marilandica</i> |
| 561 | <i>Smilax pseudo-china</i> | 367 | <i>Veronica serpyllifolia</i> |
| 630 | <i>Smilax pseudo-china</i> | 428 | <i>Veronica virginica</i> |
| 081 | <i>Smilax sarsaparilla</i> | 543 | <i>Viburnum acerifolium</i> |
| 464 | <i>Smyrniium aureum</i> | 064 | <i>Viburnum nudum</i> |
| 549 | <i>Smyrniium integerrimum</i> | 047 | <i>Viburnum prunifolium</i> |
| 430 | <i>Solanum nigrum</i>
var. <i>vulgare</i> | 793 | <i>Viola palmata</i> |
| 283 | <i>Solidago canadensis</i> | 254 | <i>Viola pedata</i> |
| 283 | <i>Solidago sempervirens</i> | 470 | <i>Viola primulifolia</i> |
| 733 | <i>Solidago sempervirens</i> | 506 | <i>Vitex agnus-castus</i> |
| 139 | <i>Sonchus canadensis</i> | 696 | <i>Vitis vinifera</i> |
| 071 | <i>Sophora tinctoria</i> | 502 | <i>Xanthium strumarium</i> |
| 434 | <i>Sparganium erectum</i> | 219 | <i>Xyris indica</i> |
| 302 | <i>Spiraea aruncus</i> | 270 | <i>Yucca filamentosa</i> |
| 421 | <i>Spiraea aruncus</i> | 574 | <i>Zizania aquatica</i> |
| 302 | <i>Spiraea opulifolia</i> | | |
| 290 | <i>Spiraea trifoliata</i> | | |
| 000 | <i>Staphylea trifolia</i> | | |
| 573 | <i>Statice limonium</i> | | |
| 621 | <i>Stipa avenacea</i> | | |
| 171 | <i>Swertia difformis</i> | | |
| 097 | <i>Tetragonotheca heli-</i>
<i>anthoides</i> | | |
| | | 000 | <i>Acer rubrum</i> |
| | | 000 | <i>Agrimonia eupatoria</i> |

**Species and Collection Numbers
of Vascular Plants Gathered by
John Clayton and Indirectly
Reported in Linnaeus' First
Edition of Species Plantarum**

000	<i>Agrostis virginica</i>	000	<i>Tilia americana</i>
000	<i>Amaranthus lividus</i>	000	<i>Trillium sessile</i>
000	<i>Andromeda arborea</i>	000	<i>Uniola paniculata</i>
000	<i>Andromeda calyculata</i>	000	<i>Verbena spuria</i>
000	<i>Arethusa bulbosa</i>	001	<i>Cynanchus suberosum</i>
000	<i>Aristolochia serpentaria</i>	003	<i>Drosera rotundifolia</i>
000	<i>Asclepias syriaca</i>	004	<i>Galax aphylla</i>
000	<i>Atriplex laciniata</i>	005	<i>Lobelia cardinalis</i>
000	<i>Betula lenta</i>	008	<i>Osmunda virginiana</i>
000	<i>Callicarpa americana</i>	009	<i>Aster rigidus</i>
000	<i>Carpinus ostrya</i>	009	<i>Gerardia flava</i>
000	<i>Celastrus scandens</i>	010	<i>Chelone glabra</i>
000	<i>Cercis canadensis</i>	010	<i>Gentiana saponaria</i>
000	<i>Chenopodium album</i>	011	<i>Acrostichum areolatum</i>
000	<i>Crataegus crus-galli</i>	011	<i>Osmunda regalis</i>
000	<i>Datura stramonium</i>	013	<i>Cicuta maculata</i>
000	<i>Dirca palustris</i>	013	<i>Helianthus angustifolius</i>
000	<i>Eupatorium perfoliatum</i>	014	<i>Acrostichum platyneuros</i>
000	<i>Euphorbia ipecacuanhae</i>	014	<i>Serratula squarrosa</i>
000	<i>Fagus pumila</i>	015	<i>Prenanthes alba</i>
000	<i>Fraxinus americana</i>	015	<i>Serratula glauca</i>
000	<i>Gerardia purpurea</i>	016	<i>Liriodendron tulipifera</i>
000	<i>Gleditsia triacanthos</i>	017	<i>Dracontium foetidum</i>
000	<i>Ilex aquifolium</i>	018	<i>Sisyrinchium bermudiana</i>
000	<i>Juglans alba</i>	019	<i>Leontodon dandelion</i>
000	<i>Juglans nigra</i>	019	<i>Salvia lyrata</i>
000	<i>Laurus aestivalis</i>	020	<i>Campanula perfoliata</i>
000	<i>Laurus borbonia</i>	021	<i>Kalmia angustifolia</i>
000	<i>Lepidium virginicum</i>	022	<i>Mediola virginiana</i>
000	<i>Liquidambar styraciflua</i>	024	<i>Magnolia virginiana</i> var. <i>foetida</i>
000	<i>Lonicera marilandica</i>	027	<i>Lycopodium alopecuroides</i>
000	<i>Lupinus perennis</i>	028	<i>Mitchella repens</i>
000	<i>Lycopsis virginica</i>	028	<i>Sanicula marilandica</i>
000	<i>Myrica cerifera</i>	030	<i>Andromeda mariana</i>
000	<i>Nyssa aquatica</i>	031	<i>Utricularia subulata</i>
000	<i>Oenothera biennis</i>	032	<i>Azalea viscosa</i>
000	<i>Ornithogalum hirsutum</i>	033	<i>Schwalbea americana</i>
000	<i>Osmunda claytoniana</i>	034	<i>Magnolia virginiana</i>
000	<i>Oxalis violacea</i>	035	<i>Convallaria racemosa</i>
000	<i>Panax quinquefolius</i>	036	<i>Oenothera fruticosa</i>
000	<i>Platanus occidentalis</i>	039	<i>Chelone hirsuta</i>
000	<i>Polygala viridescens</i>	040	<i>Cypripedium calceolus</i>
000	<i>Polygonum convolvulus</i>	042	<i>Aralia nudicaulis</i>
000	<i>Polygonum sagittatum</i>	042	<i>Vaccinium stamineum</i>
000	<i>Polygonum scandens</i>	043	<i>Crataegus coccinea</i>
000	<i>Prunus virginiana</i>	043	<i>Valeriana locusta</i> var. <i>radiata</i>
000	<i>Pyrus coronaria</i>	044	<i>Ornithogalum bivale</i>
000	<i>Quercus alba</i>	046	<i>Chionanthus virginica</i>
000	<i>Quercus nigra</i>	047	<i>Viburnum prunifolium</i>
000	<i>Quercus phellos</i>	049	<i>Nyssa aquatica</i>
000	<i>Quercus prinus</i>	050	<i>Robinia pseudoacacia</i>
000	<i>Quercus rubra</i>	052	<i>Azalea lutea</i>
000	<i>Rhus toxicodendron</i>	053	<i>Orontium aquaticum</i>
000	<i>Rumex britannica</i>	054	<i>Laurus benzoin</i>
000	<i>Rumex verticillatus</i>	055	<i>Crataegus tomentosa</i>
000	<i>Sanicula canadensis</i>	056	<i>Arabis lyrata</i>
000	<i>Sanicula europaea</i>	056	<i>Laurus sassafras</i>
000	<i>Scrophularia nodosa</i>	058	<i>Annona muricata</i>
000	<i>Serratula scariosa</i>		
000	<i>Staphylea trifolia</i>		

060	<i>Houstonia caerulea</i>	130	<i>Mimulus ringens</i>
060	<i>Mespilus canadensis</i>	131	<i>Sida rhombifolia</i>
063	<i>Houstonia purpurea</i>	133	<i>Cacalia atriplicifolia</i>
064	<i>Viburnum nudum</i>	134	<i>Melothria pendula</i>
065	<i>Asclepias nivea</i>	135	<i>Hypericum setosum</i>
066	<i>Arum triphyllum</i>	135	<i>Teucrium canadense</i>
067	<i>Coix lacryma-jobi</i>	136	<i>Helianthus atrorubens</i>
069	<i>Ceanothus americanus</i>	137	<i>Ludwigia alternifolia</i>
070	<i>Andropogon divaricatum</i>	138	<i>Osteospermum uvedalia</i>
071	<i>Sophora tinctoria</i>	139	<i>Sonchus canadensis</i>
072	<i>Aster dumosus</i>	140	<i>Monarda punctata</i>
072	<i>Aster linifolius</i>	141	<i>Satureja virginiana</i>
073	<i>Andromeda paniculata</i>	142	<i>Buchnera americana</i>
074	<i>Aletris farinosa</i>	143	<i>Aster divaricatus</i>
075	<i>Euonymus americanus</i>	146	<i>Cassia ligustrina</i>
076	<i>Limodorum tuberosum</i>	147	<i>Eupatorium scandens</i>
077	<i>Arethusa ophioglossoides</i>	148	<i>Elephantopus tomentosus</i>
078	<i>Prinos verticillatus</i>	151	<i>Passiflora incarnata</i>
079	<i>Hydrangea arborescens</i>	152	<i>Euphorbia polygonifolia</i>
080	<i>Diospyros virginiana</i>	155	<i>Euphorbia corollata</i>
081	<i>Smilax sarsaparilla</i>	156	<i>Cassia chamaecrista</i>
082	<i>Smilax lanceolata</i>	157	<i>Polygala cruciata</i>
082	<i>Smilax pseudo-china</i>	158	<i>Penthorum sedoides</i>
083	<i>Asclepias decumbens</i>	159	<i>Baccharis foetida</i>
085	<i>Hedysarum repens</i>	161	<i>Veronica anagallis-</i> <i>aquatica</i>
085	<i>Ruellia strepens</i>	161	<i>Veronica beccabunga</i>
087	<i>Pontederia cordata</i>	162	<i>Eupatorium purpureum</i>
088	<i>Pyrola maculata</i>	163	<i>Verbesina alba</i>
092	<i>Trifolium biflorum</i>	164	<i>Gratiola dubia</i>
093	<i>Commelina communis</i>	165	<i>Erigeron camphoratum</i>
093	<i>Commelina erecta</i>	166	<i>Verbesina virginica</i>
094	<i>Dioscorea villosa</i>	168	<i>Hyssopus nepetoides</i>
097	<i>Tetragonotheca heli-</i> <i>anthoides</i>	171	<i>Swertia difformis</i>
098	<i>Ruellia strepens</i>	174	<i>Hedysarum frutescens</i>
099	<i>Cactus opuntia</i>	175	<i>Serratula glauca</i>
100	<i>Bignonia crucigera</i>	177	<i>Trichostema dichotomum</i>
102	<i>Cracca virginiana</i>	179	<i>Eupatorium coelestinum</i>
103	<i>Hedysarum violaceum</i>	182	<i>Glycine comosa</i>
105	<i>Scutellaria integrifolia</i>	183	<i>Polygonum virginianum</i>
106	<i>Cephalanthus occidentalis</i>	184	<i>Hedysarum paniculatum</i>
107	<i>Saururus cernuus</i>	185	<i>Lycopus virginicus</i>
109	<i>Helianthus giganteus</i>	187	<i>Ophiorrhiza mitreola</i>
110	<i>Sarothra gentianoides</i>	187	<i>Silphium asteriscus</i>
111	<i>Myosotis virginiana</i>	190	<i>Hedysarum viridiflorum</i>
112	<i>Clitoria virginiana</i>	191	<i>Medicago virginica</i>
113	<i>Glycine tomentosa</i>	192	<i>Gerardia pedicularia</i>
116	<i>Hedera quinquefolia</i>	193	<i>Carduus virginianus</i>
117	<i>Teucrium virginicum</i>	194	<i>Aster ericoides</i>
118	<i>Passiflora lutea</i>	195	<i>Helianthus laevis</i>
120	<i>Chironia dodecandra</i>	196	<i>Lobelia cliffortiana</i>
121	<i>Dolichos regularis</i>	197	<i>Satureja origanoides</i>
122	<i>Hibiscus moscheutos</i>	198	<i>Melissa nepeta</i>
123	<i>Cicuta maculata</i>	199	<i>Ageratum altissimum</i>
124	<i>Hedysarum nudiflorum</i>	199	<i>Geum virginianum</i>
125	<i>Angelica sylvestris</i>	200	<i>Oenothera mollissima</i>
126	<i>Crotalaria sagittalis</i>	201	<i>Acalypha virginica</i>
127	<i>Glycine apios</i>	201	<i>Lonicera symphoricarpos</i>
128	<i>Physalis viscosa</i>	202	<i>Helenium autumnale</i>
129	<i>Phryma leptostachya</i>	203	<i>Gnaphalium obtusifolium</i>

- | | | | |
|-----|---------------------------------|-----|-----------------------------------|
| 206 | <i>Cenchrus tribuloides</i> | 279 | <i>Sium rigidius</i> |
| 208 | <i>Bupthalmum helianthoides</i> | 280 | <i>Scutellaria lateriflora</i> |
| 209 | <i>Hedysarum canescens</i> | 281 | <i>Lonicera symphoricarpos</i> |
| 212 | <i>Monarda clinopodia</i> | 282 | <i>Eryngium foetidum</i> |
| 214 | <i>Lythrum verticillatum</i> | 283 | <i>Solidago canadensis</i> |
| 215 | <i>Cicuta bulbifera</i> | 283 | <i>Solidago sempervirens</i> |
| 215 | <i>Cuscuta americana</i> | 284 | <i>Prenanthes alba</i> |
| 216 | <i>Asclepias verticillata</i> | 286 | <i>Obolaria virginica</i> |
| 219 | <i>Xyris indica</i> | 286 | <i>Senecio aureus</i> |
| 220 | <i>Scrophularia marilandica</i> | 287 | <i>Gnaphalium plantaginifolia</i> |
| 221 | <i>Osteospermum uvedalia</i> | 288 | <i>Asarum canadense</i> |
| 222 | <i>Asclepias incarnata</i> | 289 | <i>Trifolium reflexum</i> |
| 223 | <i>Cynanchum suberosum</i> | 290 | <i>Spiraea trifoliata</i> |
| 225 | <i>Bignonia radicans</i> | 291 | <i>Thapsia trifoliata</i> |
| 226 | <i>Veronica marilandica</i> | 292 | <i>Salvia urticifolia</i> |
| 227 | <i>Rhexia virginica</i> | 293 | <i>Bartsia coccinea</i> |
| 228 | <i>Arum virginicum</i> | 294 | <i>Anemone thalictroides</i> |
| 230 | <i>Ascyrum hypericoides</i> | 295 | <i>Mespilus canadensis</i> |
| 232 | <i>Hypericum mutilum</i> | 297 | <i>Phlox glaberrima</i> |
| 233 | <i>Aralia spinosa</i> | 297 | <i>Tradescantia virginiana</i> |
| 234 | <i>Eriocaulon decangulare</i> | 298 | <i>Chrysogonum virginianum</i> |
| 237 | <i>Serratula spicata</i> | 298 | <i>Coreopsis auriculata</i> |
| 238 | <i>Rhus radicans</i> | 299 | <i>Veratrum luteum</i> |
| 239 | <i>Aster grandiflorus</i> | 301 | <i>Heuchera americana</i> |
| 240 | <i>Baccharis halimifolia</i> | 302 | <i>Spiraea aruncus</i> |
| 242 | <i>Bupthalmum frutescens</i> | 302 | <i>Spiraea opulifolia</i> |
| 243 | <i>Iva frutescens</i> | 304 | <i>Anchusa virginiana</i> |
| 244 | <i>Aster novae-angliae</i> | 304 | <i>Saxifraga pensylvanica</i> |
| 245 | <i>Cucubalus stellatus</i> | 305 | <i>Actaea racemosa</i> |
| 245 | <i>Monotropa uniflora</i> | 307 | <i>Geranium maculatum</i> |
| 246 | <i>Urtica pumila</i> | 307 | <i>Ligusticum scothieum</i> |
| 247 | <i>Sanguinaria canadensis</i> | 308 | <i>Coreopsis verticillata</i> |
| 248 | <i>Burmannia biflora</i> | 309 | <i>Tragopogon virginicum</i> |
| 249 | <i>Hydrophyllum virginianum</i> | 310 | <i>Origanum vulgare</i> |
| 249 | <i>Polemonium caeruleum</i> | 311 | <i>Ceanothus americanus</i> |
| 249 | <i>Senecio aureus</i> | 313 | <i>Galium bermudense</i> |
| 250 | <i>Epigaea repens</i> | 314 | <i>Samolus valerandii</i> |
| 251 | <i>Claytonia virginica</i> | 316 | <i>Queria canadensis</i> |
| 253 | <i>Iris verna</i> | 317 | <i>Queria canadensis</i> |
| 254 | <i>Viola pedata</i> | 319 | <i>Prenanthes alba</i> |
| 255 | <i>Podophyllum peltatum</i> | 320 | <i>Adiantum pedatum</i> |
| 256 | <i>Amaryllis atamasca</i> | 321 | <i>Adiantum pedatum</i> |
| 256 | <i>Antirrhinum canadense</i> | 322 | <i>Polypodium lonchitis</i> |
| 257 | <i>Cynoglossum virginianum</i> | 328 | <i>Anemone hepatica</i> |
| 258 | <i>Uvularia perfoliata</i> | 329 | <i>Panax trifolius</i> |
| 259 | <i>Carex pseudocyperus</i> | 331 | <i>Lamium amplexicaule</i> |
| 259 | <i>Iris virginica</i> | 332 | <i>Juncus campestris</i> |
| 260 | <i>Orchis spectabilis</i> | 333 | <i>Oenothera fruticosa</i> |
| 261 | <i>Scutellaria hyssopifolia</i> | 335 | <i>Convallaria polygonatum</i> |
| 263 | <i>Asclepias rubra</i> | 338 | <i>Aquilegia canadensis</i> |
| 263 | <i>Parthenium integrifolium</i> | 339 | <i>Pulmonaria virginica</i> |
| 267 | <i>Dipsacus fullonum</i> | 340 | <i>Juncus bulbosus</i> |
| 269 | <i>Anthericum calyculatum</i> | 341 | <i>Equisetum arvense</i> |
| 270 | <i>Clematis vitalba</i> | 342 | <i>Cerastium semidecandrum</i> |
| 270 | <i>Yucca filamentosa</i> | 343 | <i>Plantago virginica</i> |
| 271 | <i>Betonica annua</i> | 367 | <i>Veronica serpyllifolia</i> |
| 273 | <i>Poa flava</i> | 368 | <i>Veronica arvensis</i> |
| 276 | <i>Chelidonium glaucium</i> | 372 | <i>Geranium carolinianum</i> |
| 277 | <i>Diodia virginiana</i> | 374 | <i>Aphanes arvensis</i> |
| 278 | <i>Sagittaria sagittifolia</i> | 376 | <i>Hyoseris virginica</i> |

378	<i>Callitriche palustris</i>	458	<i>Panicum dichotomum</i>
379	<i>Gratiola virginiana</i>	460	<i>Andropogon virginicum</i>
380	<i>Scirpus capitatus</i>	461	<i>Eriophorum virginicum</i>
382	<i>Polygonum aviculare</i>	462	<i>Cardamine virginica</i>
383	<i>Leontodon dandelion</i>	464	<i>Smyrnium aureum</i>
384	<i>Cupressus distichia</i>	467	<i>Quercus alba</i>
385	<i>Gnaphalium purpureum</i>	470	<i>Viola primulifolia</i>
386	<i>Hieracium venosum</i>	471	<i>Platanus occidentalis</i>
387	<i>Orobanche uniflora</i>	472	<i>Arethusa bulbosa</i>
388	<i>Silene antirrhina</i>	473	<i>Ranunculus repens</i>
389	<i>Renealmia usneoides</i>	474	<i>Oxalis stricta</i>
390	<i>Trifolium repens</i>	475	<i>Arenaria rubra</i>
391	<i>Aster vernus</i>		var. <i>marina</i>
391	<i>Salvia lyrata</i>	479	<i>Rhus toxicodendron</i>
393	<i>Juncus effusus</i>	480	<i>Itea virginica</i>
394	<i>Arabis lyrata</i>	485	<i>Laurus indica</i>
399	<i>Mollugo verticillata</i>	487	<i>Liquidambar styraciflua</i>
400	<i>Arabis canadensis</i>	488	<i>Rhinanthus virginica</i>
404	<i>Magnolia virginiana</i>	490	<i>Rudbeckia hirta</i>
	var. <i>acuminata</i>	490	<i>Rudbeckia purpurea</i>
407	<i>Scandix cerefolium</i>	492	<i>Rhus glabra</i>
407	<i>Scandix procumbens</i>	494	<i>Rumex acetosella</i>
408	<i>Dianthera americana</i>	496	<i>Pinus taeda</i>
410	<i>Portulaca oleracea</i>	498	<i>Agave virginica</i>
411	<i>Clematis viorna</i>	500	<i>Eryngium aquaticum</i>
412	<i>Monarda ciliata</i>	502	<i>Xanthium strumarium</i>
414	<i>Polygala senega</i>	504	<i>Convolvulus hederaceus</i>
417	<i>Rudbeckia purpurea</i>	505	<i>Lythrum lineare</i>
418	<i>Lythrum petiolatum</i>	506	<i>Vitex agnus-castus</i>
419	<i>Lysimachia quadrifolia</i>	507	<i>Agrostis virginica</i>
421	<i>Spiraea aruncus</i>	508	<i>Urtica cylindrica</i>
422	<i>Melanthium virginicum</i>	509	<i>Cyperus odoratus</i>
423	<i>Silene virginica</i>	510	<i>Hedysarum hirtum</i>
424	<i>Heuchera americana</i>	511	<i>Sigesbeckia occidentalis</i>
425	<i>Menispermum virginicum</i>	512	<i>Ambrosia elatior</i>
428	<i>Veronica virginica</i>	514	<i>Melissa pulegioides</i>
429	<i>Hydrocotyle umbellata</i>	515	<i>Utricularia gibba</i>
430	<i>Solanum nigrum</i>	516	<i>Hedysarum marilandicum</i>
	var. <i>vulgare</i>	517	<i>Utricularia gibba</i>
431	<i>Verbena officifolia</i>	520	<i>Laurus aestivalis</i>
432	<i>Salsola kali</i>	522	<i>Caltha palustris</i>
433	<i>Lysimachia punctata</i>	524	<i>Ulmus americana</i>
434	<i>Sparganium erectum</i>	525	<i>Draba verna</i>
435	<i>Antirrhinum elatine</i>	525	<i>Saxifraga nivalis</i>
437	<i>Nepeta virginica</i>	526	<i>Crataegus viridis</i>
438	<i>Apocynum cannabinum</i>	527	<i>Salicornia virginica</i>
439	<i>Eriocaulon decangulare</i>	528	<i>Sisymbrium nasturtium-</i>
440	<i>Linum virginianum</i>		aquaticum
441	<i>Sida abutilon</i>	529	<i>Alsine media</i>
442	<i>Amaranthus graecizans</i>	529	<i>Anemone virginiana</i>
443	<i>Teucrium chamaepitys</i>	530	<i>Acer negundo</i>
444	<i>Daucus carota</i>	531	<i>Liquidambar styraciflua</i>
446	<i>Elymus virginicus</i>	532	<i>Populus heterophylla</i>
447	<i>Hieracium gronovii</i>	537	<i>Vaccinium frondosum</i>
448	<i>Verbena nodiflora</i>	539	<i>Rudbeckia laciniata</i>
449	<i>Erigeron canadense</i>	541	<i>Smilax herbacea</i>
454	<i>Panicum capillare</i>	541	<i>Smilax pseudo-china</i>
456	<i>Scirpus spadicus</i>	543	<i>Viburnum acerifolium</i>
457	<i>Panicum sanguinale</i>	544	<i>Hamamelis virginiana</i>
457	<i>Scirpus retrofractus</i>	545	<i>Leontice thalictroides</i>

- | | | | |
|-----|----------------------------------|-----|---|
| 546 | <i>Menispermum canadense</i> | 624 | <i>Celtis occidentalis</i> |
| 547 | <i>Pinus balsamea</i> | 625 | <i>Fumaria cucullaria</i> |
| 548 | <i>Ferula canadensis</i> | 626 | <i>Triosteum angustifolium</i> |
| 549 | <i>Smyrniun integerrimum</i> | 627 | <i>Prunus virginiana</i> |
| 552 | <i>Hypericum canadense</i> | 630 | <i>Smilax pseudo-china</i> |
| 553 | <i>Convolvulus spithameus</i> | 632 | <i>Diospyros virginiana</i> |
| 554 | <i>Tiarella cordifolia</i> | 634 | <i>Rubus caesius</i> |
| 556 | <i>Itea virginica</i> | 635 | <i>Arethusa divaricata</i> |
| 556 | <i>Polemonium dubium</i> | 639 | <i>Orchis flava</i> |
| 559 | <i>Sarracenia flava</i> | 641 | <i>Convolvulus panduratus</i> |
| 560 | <i>Orchis ciliaris</i> | 646 | <i>Elatine hydropiper</i> |
| 561 | <i>Smilax pseudo-china</i> | 647 | <i>Lithospermum virginianum</i> |
| 562 | <i>Cyperus arundinacea</i> | 649 | <i>Sagina virginica</i> |
| 563 | <i>Polygala verticillata</i> | 651 | <i>Serratula squarrosa</i> |
| 564 | <i>Hedysarum virginicum</i> | 654 | <i>Mentha spicata</i> var. <i>viridis</i> |
| 565 | <i>Astragalus canadensis</i> | 655 | <i>Elephantopus scaber</i> |
| 567 | <i>Hibiscus virginicus</i> | 657 | <i>Equisetum hyemale</i> |
| 568 | <i>Dolichos polystachyus</i> | 657 | <i>Rudbeckia triloba</i> |
| 569 | <i>Amaranthus spinosus</i> | 658 | <i>Ophrys lilifolia</i> |
| 570 | <i>Scirpus glomeratus</i> | 660 | <i>Saponaria officinalis</i> |
| 571 | <i>Atriplex halimus</i> | 664 | <i>Potamogeton nutans</i> |
| 573 | <i>Statice limonium</i> | 665 | <i>Convolvulus repens</i> |
| 574 | <i>Zizania aquatica</i> | 667 | <i>Coreopsis angustifolia</i> |
| 576 | <i>Celosia paniculata</i> | 667 | <i>Salicornia virginica</i> |
| 577 | <i>Dactylis cynosuroides</i> | 668 | <i>Orchis psycodes</i> |
| 578 | <i>Panicum virgatum</i> | 670 | <i>Polygonum persicaria</i> |
| 579 | <i>Panicum glaucum</i> | 671 | <i>Phytolacca americana</i> |
| 580 | <i>Juncus filiformis</i> | 672 | <i>Polygonum persicaria</i> |
| 580 | <i>Poa capillaris</i> | 673 | <i>Hamamelis virginiana</i> |
| 581 | <i>Arundo phragmites</i> | 674 | <i>Onoclea sensibilis</i> |
| 581 | <i>Poa capillaris</i> | 679 | <i>Populus nigra</i> |
| 582 | <i>Briza eragrostis</i> | 681 | <i>Rhus vernix</i> |
| 583 | <i>Dactylis cynosuroides</i> | 682 | <i>Pteris atropurpurea</i> |
| 585 | <i>Schoenus glomeratus</i> | 684 | <i>Myrica asplenifolia</i> |
| 586 | <i>Epilobium hirsutum</i> | 685 | <i>Acrostichum polypodoides</i> |
| 587 | <i>Oldenlandia uniflora</i> | 688 | <i>Betula nigra</i> |
| 589 | <i>Holcus laxus</i> | 691 | <i>Erythronium dens-canis</i> |
| 590 | <i>Holcus striatus</i> | 696 | <i>Vitis vinifera</i> |
| 591 | <i>Panicum italicum</i> | 699 | <i>Potentilla reptans</i> |
| 592 | <i>Glycine bracteata</i> | 701 | <i>Ranunculus abortivus</i> |
| 594 | <i>Gratiola dubia</i> | 703 | <i>Rubus fruticosus</i> |
| 595 | <i>Phalaris oryzoides</i> | 704 | <i>Asarum virginicum</i> |
| 599 | <i>Acnida cannabina</i> | 708 | <i>Ophrys lilifolia</i> |
| 600 | <i>Andropogon divaricatum</i> | 710 | <i>Ranunculus repens</i> |
| 601 | <i>Andropogon altopecuroides</i> | 714 | <i>Onoclea sensibilis</i> |
| 602 | <i>Andropogon hirtum</i> | 717 | <i>Sarracenia purpurea</i> |
| 603 | <i>Eupatorium aromaticum</i> | 719 | <i>Myrica asplenifolia</i> |
| 604 | <i>Orobanche virginiana</i> | 721 | <i>Sison canadense</i> |
| 605 | <i>Gentiana villosa</i> | 723 | <i>Alisma subulata</i> |
| 606 | <i>Andropogon virginicum</i> | 724 | <i>Ambrosia trifida</i> |
| 606 | <i>Panicum virgatum</i> | 728 | <i>Rhus copallinum</i> |
| 607 | <i>Aster concolor</i> | 732 | <i>Bunias cakile</i> |
| 609 | <i>Rudbeckia oppositifolia</i> | 733 | <i>Solidago sempervirens</i> |
| 610 | <i>Lechea minor</i> | 741 | <i>Oxalis longiflora</i> |
| 610 | <i>Silphium helianthoides</i> | 745 | <i>Arabis canadensis</i> |
| 614 | <i>Hedysarum virginicum</i> | 767 | <i>Aster divaricatus</i> |
| 617 | <i>Smilax laurifolia</i> | 768 | <i>Polypremum procumbens</i> |
| 620 | <i>Eupatorium trifoliatum</i> | 774 | <i>Ammannia ramosior</i> |
| 621 | <i>Andropogon nutans</i> | 793 | <i>Viola palmata</i> |
| 621 | <i>Stipa avenacea</i> | 884 | <i>Juniperus virginiana</i> |

**Species of Vascular Plants
from Temperate North America
Attributed to Peter Kalm by
Linnaeus in Species Plantarum
or in the Linnaean Herbarium**

Acalypha virginica	Cephalanthus occidentalis
Acer negundo	Cercis canadense
Acer pensylvanicum	Chaerophyllum arborescens
Acer rubrum	Chelone glabra
Acer saccharinum	Chenopodium anthelminticum
Adiantum pedatum	Chenopodium virginicum
Agrostis virginica	Chironia angularis
Allium canadense	Chironia campanulata
Amaranthus retroflexus	Chrysocoma graminifolia
Andromeda mariana	Chrysosplenium oppositifolium
Andromeda paniculata	Cicuta bulbifera
Andromeda racemosa	Cicuta maculata
Andropogon nutans	Cinna arundinacea
Anemone quinquefolia	Cistus canadensis
Anemone thalictroides	Claytonia virginica
Anemone virginiana	Collinsonia canadensis
Angelica sylvestris	Convolvulus spithameus
Antirrhinum canadense	Conyza asteroides
Arabis canadensis	Cornus canadensis
Arabis lyrata	Cornus florida
Aralia spinosa	Cracca virginiana
Arenaria rubra var. marina	Crataegus coccinea
Arethusa bulbosa	Crataegus crus-galli
Arethusa ophioglossoides	Crataegus tomentosa
Aristolochia serpentaria	Crotalaria sagittalis
Asarum canadense	Cucubalus stellatus
Asclepias nivea	Cupressus thyoides
Asclepias tuberosa	Cuscuta americana
Asplenium rhizophyllum	Cynoglossum virginianum
Aster concolor	Cyperus strigosus
Aster cordifolius	Dalibarda repens
Aster laevis	Datisca hirta
Aster linariifolius	Dianthera americana
Aster novi-belgii	Diospyros virginiana
Aster puniceus	Dracocephalum virginianum
Aster undulatus	Dracontium foetidum
Astragalus canadensis	Elymus canadensis
Avena pensylvanica	Erigeron philadelphicus
Avena spicata	Eupatorium rotundifolium
Azalea lutea	Euphorbia corollata
Azalea viscosa	Euphorbia ipecacuanhae
Bartsia coccinea	Euphorbia polygonifolia
Briza eragrostis	Euphorbia portulacoides
Bromus ciliatus	Euonymus americanus
Bromus purgans	Fagus pumila
Cacalia atriplicifolia	Galium tinctorium
Carex folliculata	Galium trifidum
Carex squarrosa	Gentiana quinquefolia
Carpinus betulus	Gentiana saponaria
Cassia nictitans	Geranium carolinianum
Ceanothus americanus	Gerardia flava
Celastrus myrtifolius	Gerardia pedicularia
Celtis occidentalis	Gerardia purpurea
Cenchrus tribuloides	Gleditsia triacanthos
	Glycine apios
	Glycine bracteata
	Glycine comosa
	Gnaphalium margaritaceum
	Gnaphalium obtusifolium
	Gnaphalium plantaginifolium

Gnaphalium purpureum	Ophrys cernua
Gratiola virginiana	Ophrys lilifolia
Hamamelis virginiana	Orchis psycodes
Hedera quinquefolia	Origanum vulgare
Hedysarum hirtum	Ornithogalum hirsutum
Hedysarum violaceum	Orobanche virginiana
Helianthus decapetalus	Osmunda cinnamomea
Heuchera americana	Osmunda regalis
Hibiscus palustris	Osmunda virginiana
Hieracium gronovii	Oxalis stricta
Hieracium kalmii	Oxalis violacea
Hieracium paniculatum	Panax quinquefolius
Hippophae canadensis	Panicum clandestinum
Hordeum jubatum	Panicum crusgalli
Houstonia caerulea	Panicum dichotomum
Houstonia purpurea	Panicum dissectum
Hydrangea arborescens	Panicum filiforme
Hydrocotyle americana	Panicum latifolium
Hydrocotyle umbellata	Phlox maculata
Hyoseris virginica	Phlox subulata
Hypericum canadense	Pinus strobus
Hypericum kalmianum	Pinus taeda
Hypericum mutilum	Plantago virginica
Impatiens noti-tangere	Poa capillaris
Iris verna	Polygala incarnata
Juglans alba	Polygala lutea
Lactuca canadensis	Polygala sanguinea
Laurus sassafra	Polygala verticillata
Lilium canadense	Polygala viridescens
Limodorum tuberosum	Polygonum articulatum
Linum virginianum	Polygonum erectum
Liquidambar peregrina	Polygonum pensylvanicum
Liquidambar styraciflua	Polypodium marginale
Liriodendron tulipifera	Polypodium noveboracense
Lobelia cliffortiana	Polymnia canadensis
Lobelia kalmii	Pontederia cordata
Ludwigia alternifolia	Potentilla canadensis
Lupinus perennis	Prenanthes alba
Lycopodium alopecuroides	Prinos glaber
Lycopodium apodum	Prinos verticillatus
Lycopodium complanatum	Pteris atropurpurea
Lycopodium obscurum	Pulmonaria virginica
Lycopodium rupestre	Pyrola maculata
Magnolia virginiana	Pyrola umbellata
var. glauca	Quercus alba
Medeola virginica	Quercus nigra
Melanthium virginicum	Quercus phellos
Melissa pulegioides	Quercus prinus
Menispermum canadense	Quercus rubra
Mentha canadensis	Ranunculus abortivus
Mespilus arbutifolia	Rhexia virginica
Mespilus canadensis	Rhus glabra
Mitella diphylla	Rhus radicans
Mollugo verticillata	Rhus toxicodendron
Monotropa uniflora	Rhus vernix
Morus rubra	Ribes cynosbati
Nymphaea lotus	Rubus canadensis
Nymphaea lutea	Rubus hispidus
Nyssa aquatica	Rubus occidentalis
Oenothera fruticosa	Rumex persicarioides

Salvia lyrata
Sambucus canadensis
Sanguinaria canadensis
Sanguisorba canadensis
Sarothra gentianoides
Satureja origanoides
Saxifraga nivalis
Saxifraga pensylvanica
Schoenus glomeratus
Scirpus capitatus
Scutellaria hyssopifolia
Scutellaria integrifolia
Senecio canadensis
Serratula praealta
Serratula spicata
Silene nocturna
Silene virginica
Sison canadense
Sisyrinchium bermudiana
Smilax caduca
Smilax pseudo-china
Smilax rotundifolia
Smilax tamnoides
Smyrnium aureum
Solidago canadensis
Solidago flexicaulis
Solidago lateriflora
Sonchus canadensis
Sophora tinctoria
Spiraea opulifolia
Spiraea tomentosa
Spiraea trifoliata
Swertia corniculata
Thalictrum dioicum
Thapsia trifoliata
Thesium umbellatum
Thuja occidentalis
Trogopogon virginicum
Trifolium arvense
Trifolium biflorum
Trillium cernuum
Trillium erectum
Ulmus americana
Uniola spicata
Urtica capitata
Urtica cylindrica
Urtica divaricata
Urtica pumila
Utricularia subulata
Uvularia perfoliata
Uvularia sessilifolia
Vaccinium album
Vaccinium corymbosum
Vaccinium frondosum
Vaccinium hispidulum
Vaccinium ligustrinum
Vaccinium mucronatum
Vaccinium stamineum
Veratrum luteum
Viburnum acerifolium

Viburnum lentago
Viburnum prunifolium
Viola canadensis
Viola lanceolata
Viola pedata
Viscum terrestre
Vitis labrusca
Vitis laciniosa
Vitis vinifera
Vitis vulpina
Xyris indica

**Selected Literature Cited by
 Linnaeus in the First Edition
 of Species Plantarum for Seem-
 ingly Temperate North American
 Vascular Plants**

Bauhin

Acnida cannabina
Adiantum pedatum
Amaranthus lividus
Cornus canadensis
Diospyros virginiana
Helianthus multiflorus
Hibiscus palustris
Laurus sassafras
Othonna cineraria
Rhus glabra
Rudbeckia laciniata
Sarracenia flava
Sarracenia purpurea
Smilax bona-nox
Spiraea hypericifolia
Tradescantia virginiana
Trillium erectum
Uvularia perfoliata
Verbena nodiflora
Vitis labrusca

Catesby

Acer rubrum
Amaryllis atamasca
Andromeda arborea
Andromeda paniculata
Annona glabra
Annona triloba
Arethusa divaricata
Aristolochia serpentaria
Azalea viscosa
Bignonia caerulea
Bignonia radicans
Bignonia sempervirens
Callicarpa americana
Chionanthus virginica
Cissampelos smilacina
Clethra alnifolia
Cornus florida
Cupressus distichia
Diospyros virginiana

Dodecatheon meadia
 Erythrina herbacea
 Fagus pumila
 Fraxinus americana
 Gentiana saponaria
 Gleditsia triacanthos
 Hamamelis virginiana
 Ilex cassine
 Ipomoea carolina
 Juglans alba
 Juglans nigra
 Kalmia angustifolia
 Kalmia latifolia
 Laurus borbonia
 Laurus sassafras
 Lilium canadense
 Liriodendron tulipifera
 Lonicera marilandica
 Magnolia virginiana
 var. acuminata
 Magnolia virginiana
 var. foetida
 Magnolia virginiana
 var. glauca
 Magnolia virginiana
 var. tripetala
 Mitchella repens
 Monotropa uniflora
 Myrica cerifera
 Nyssa aquatica
 Panax quinquefolius
 Philadelphus inodorus
 Platanus occidentalis
 Podophyllum peltatum
 Populus balsamifera
 Prunus virginiana
 Quercus alba
 Quercus nigra
 Quercus phellos
 Quercus prinus
 Quercus rubra
 Rhododendron maximum
 Robinia pseudoacacia
 Rudbeckia purpurea
 Rumex sanguineus
 Sarracenia flava
 Sarracenia purpurea
 Sloanea emarginata
 Smilax laurifolia
 Smilax tannoides
 Stewartia malacodendron
 Trillium cernuum
 Trillium sessile
 Uniola paniculata
 Viscum purpureum
 Viscum rubrum

Colden

Acer rubrum
 Ambrosia elatior

Aralia racemosa
 Azalea lutea
 Azalea viscosa
 Collinsonia canadensis
 Cucubalus stellatus
 Dracontium foetidum
 Eupatorium perfoliatum
 Eupatorium purpureum
 Eupatorium scandens
 Gaultheria procumbens
 Hamamelis virginiana
 Hieracium venosum
 Holosteum succulentum
 Kalmia angustifolia
 Laurus sassafras
 Leontice thalictroides
 Mitella diphylla
 Monarda didyma
 Myrica asplenifolia
 Pinus strobus
 Pinus taeda
 Polygonum sagittatum
 Polygonum scandens
 Pontederia cordata
 Rhus glabra
 Rumex britannica
 Saxifraga pensylvanica
 Staphylea trifolia
 Trillium cernuum
 Uvularia perfoliata
 Uvulariasessilifolia
 Veronica virginica

Cornuti

Actaea spicata var. alba
 Adiantum pedatum
 Ageratum altissimum
 Angelica atropurpurea
 Angelica lucida
 Aquilegia canadensis
 Aralia racemosa
 Asarum canadense
 Asclepias incarnata
 Asclepias syriaca
 Aster annuus
 Bignonia radicans
 Convallaria racemosa
 Convallaria stellata
 Eupatorium purpureum
 Fumaria cucullaria
 Fumaria sempervirens
 Glycine apios
 Hedera quinquefolia
 Hedysarum canadense
 Helenium autumnale
 Hibiscus moscheutos
 Monarda fistulosa
 Polypodium bulbiferum
 Rhus radicans
 Rubus odoratus

Rudbeckia laciniata
Sanguinaria canadensis
Sanguisorba canadensis
Solidago sempervirens
Thalictrum cornutii
Trillium erectum
Uvularia perfoliata

Dillenius

Actaea racemosa
Asclepias amoena
Asclepias purpurascens
Asclepias tuberosa
Asclepias variagata
Aster ericoides
Aster grandiflorus
Aster miser
Astragalus canadensis
Astragalus carolinianus
Baccharis foetida
Bidens nivea
Bidens pilosa
Canna glauca
Carduus altissimus
Clematis crispa
Clematis viorna
Clitoria virginiana
Commelina communis
Commelina erecta
Convolvulus carolinus
Coreopsis lanceolata
Crotalaria perfoliata
Erythrina herbacea
Eupatorium coelestinum
Eupatorium hyssopifolium
Geranium carolinianum
Glycine tomentosa
Gnaphalium obtusifolium
Gnaphalium purpureum
Hedysarum marilandicum
Hedysarum volubile
Helianthus atrorubens
Horminum virginicum
Ipomoea lacunosa
Ipomoea tannifolia
Iris versicolor
Lonicera symphoricarpos
Lycopodium alopecuroides
Lycopodium apodum
Lycopodium carolinianum
Lycopodium obscurum
Magnolia virginiana
var. glauca
Malva caroliniana
Mensipermum virginicum
Parthenium integrifolium
Phaseolus helvulus
Phlox glaberrima
Polemonium rubrum
Polygonum scandens

Ptelea trifoliata
Rhus glabra
Rhus radicans
Ribes oxycanthoides
Rosa carolina
Rubus occidentalis
Rudbeckia hirta
Ruellia biflora
Ruellia strepens
Sanguinaria canadensis
Saxifraga pensylvanica
Serratula glauca
Serratula praealta
Serratula spicata
Serratula squarrosa
Silene antirrhina
Silphium asteriscus
Solanum carolinense
Solanum virginianum
Solidago altissima
Solidago caesia
Spermacoce tenuior
Tetragonotheca helianthoides
Trichostema brachiatum
Triosteum perfoliatum

Gronovius

Acer negundo
Acer rubrum
Acnida cannabina
Acrostichum areolatum
Acrostichum platyneuros
Actaea racemosa
Adiantum pedatum
Agave virginica
Ageratum altissimum
Alettris farinosa
Alisma subulata
Amaranthus graecizans
Amaranthus lividus
Amaryllis atamasca
Ambrosia elatior
Ambrosia trifida
Anchusa virginiana
Andromeda arborea
Andromeda mariana
Andromeda paniculata
Andropogon alopecuroides
Andropogon divaricatum
Andropogon virginicum
Anemone thalictroides
Anemone virginiana
Antirrhinum canadense
Apocynum cannabinum
Aquilegia canadensis
Arabis canadensis
Arabis lyrata
Aralia spinosa
Arethusa bulbosa
Arethusa divaricata

<i>Arethusa ophioglossoides</i>	<i>Coreopsis angustifolia</i>
<i>Aristolochia serpentaria</i>	<i>Coreopsis auriculata</i>
<i>Arum virginicum</i>	<i>Coreopsis verticillata</i>
<i>Asarum canadense</i>	<i>Cornus florida</i>
<i>Asarum virginicum</i>	<i>Cracca virginiana</i>
<i>Asclepias decumbens</i>	<i>Crataegus coccinea</i>
<i>Asclepias incarnata</i>	<i>Crataegus crus-galli</i>
<i>Asclepias rubra</i>	<i>Crataegus tomentosa</i>
<i>Asclepias syriaca</i>	<i>Crataegus viridis</i>
<i>Asclepias verticillata</i>	<i>Cucubalus stellatus</i>
<i>Ascyrum hypericoides</i>	<i>Cupressus distichia</i>
<i>Aster concolor</i>	<i>Cuscuta americana</i>
<i>Aster divaricatus</i>	<i>Cynoglossum virginianum</i>
<i>Aster dumosus</i>	<i>Cyperus arundinacea</i>
<i>Aster ericoides</i>	<i>Cyperus odoratus</i>
<i>Aster grandiflorus</i>	<i>Dianthera americana</i>
<i>Aster linifolius</i>	<i>Diodia virginiana</i>
<i>Aster novae-angliae</i>	<i>Dioscorea villosa</i>
<i>Aster rigidus</i>	<i>Diospyros virginiana</i>
<i>Aster vernus</i>	<i>Dirca palustris</i>
<i>Azalea lutea</i>	<i>Dolichos polystachyus</i>
<i>Azalea viscosa</i>	<i>Dolichos regularis</i>
<i>Baccharis foetida</i>	<i>Dracontium foetidum</i>
<i>Baccharis halimifolia</i>	<i>Elephantopus tomentosus</i>
<i>Bartsia coccinea</i>	<i>Elymus virginicus</i>
<i>Betula lenta</i>	<i>Epigaea repens</i>
<i>Betula nigra</i>	<i>Erigeron camphoratum</i>
<i>Bignonia radicans</i>	<i>Eriocaulon decangulare</i>
<i>Buchnera americana</i>	<i>Eriophorum virginicum</i>
<i>Buphthalmum helianthoides</i>	<i>Eryngium aquaticum</i>
<i>Burmanna biflora</i>	<i>Eupatorium aromaticum</i>
<i>Cacalia atriplicifolia</i>	<i>Eupatorium coelestinum</i>
<i>Callicarpa americana</i>	<i>Eupatorium perfoliatum</i>
<i>Campanula perfoliata</i>	<i>Eupatorium purpureum</i>
<i>Cardamine virginica</i>	<i>Eupatorium scandens</i>
<i>Carduus virginianus</i>	<i>Eupatorium trifoliatum</i>
<i>Cassia marilandica</i>	<i>Euphorbia corollata</i>
<i>Ceanothus americanus</i>	<i>Euphorbia ipeacacuanhae</i>
<i>Celastrus scandens</i>	<i>Euphorbia polygonifolia</i>
<i>Celosia paniculata</i>	<i>Euonymus americanus</i>
<i>Celtis occidentalis</i>	<i>Fagus pumila</i>
<i>Cenchrus tribuloides</i>	<i>Ferula canadensis</i>
<i>Cephalanthus occidentalis</i>	<i>Fraxinus americana</i>
<i>Cercis canadensis</i>	<i>Fumaria cucullaria</i>
<i>Chelone glabra</i>	<i>Galax aphylla</i>
<i>Chelone hirsuta</i>	<i>Galium bermudense</i>
<i>Chionanthus virginica</i>	<i>Gentiana saponaria</i>
<i>Chironia dodecandra</i>	<i>Gentiana villosa</i>
<i>Chrysogonum virginianum</i>	<i>Geranium carolinianum</i>
<i>Cicuta bulbifera</i>	<i>Gerardia flava</i>
<i>Cicuta maculata</i>	<i>Gerardia pedicularia</i>
<i>Claytonia virginica</i>	<i>Gerardia purpurea</i>
<i>Clematis viorna</i>	<i>Geum virginianum</i>
<i>Clitoria virginiana</i>	<i>Gleditsia triacanthos</i>
<i>Commelina communis</i>	<i>Glycine apios</i>
<i>Commelina erecta</i>	<i>Glycine bracteata</i>
<i>Convallaria racemosa</i>	<i>Glycine comosa</i>
<i>Convolvulus panduratus</i>	<i>Glycine tomentosa</i>
<i>Convolvulus repens</i>	<i>Gnaphalium obtusifolium</i>
<i>Convolvulus spithameus</i>	<i>Gnaphalium plantaginifolium</i>

Gnaphalium purpureum	Lycopodium alopecuroides
Gratiola dubia	Lycopsis virginica
Gratiola virginiana	Lycopus virginicus
Hamamelis virginiana	Lysimachia quadrifolia
Hedera quinquefolia	Lythrum lineare
Hedysarum frutescens	Lythrum petiolatum
Hedysarum hirtum	Lythrum verticillatum
Hedysarum marilandicum	Magnolia virginiana
Hedysarum nudiflorum	Magnolia virginiana
Hedysarum paniculatum	var. acuminata
Hedysarum repens	Magnolia virginiana
Hedysarum violaceum	var. foetida
Hedysarum virginicum	Medeola virginiana
Hedysarum viridiflorum	Medicago virginica
Helenium autumnale	Melanthium virginicum
Helianthus angustifolius	Melissa pulegioides
Helianthus atrorubens	Menispermum canadense
Helianthus giganteus	Menispermum virginicum
Helianthus laevis	Mespilus canadensis
Heuchera americana	Mimulus ringens
Hibiscus moscheutos	Mitchella repens
Hibiscus virginicus	Monarda ciliata
Hieracium gronovii	Monarda clinopodia
Hieracium venosum	Monarda punctata
Holcus laxus	Monotropa uniflora
Holcus striatus	Myosotis virginiana
Houstonia caerulea	Myrica asplenifolia
Houstonia purpurea	Myrica cerifera
Hydrangea arborescens	Nepeta virginica
Hydrocotyle umbellata	Nyssa aquatica
Hydrophyllum virginianum	Obolaria virginica
Hyoseris virginica	Oenothera fruticosa
Hypericum mutilum	Oldenlandia uniflora
Hypericum setosum	Onoclea sensibilis
Hyssopus nepetoides	Ophiorrhiza mitreola
Iris verna	Orchis ciliaris
Iris virginica	Orchis flava
Itea virginica	Orchis pycodes
Juglans alba	Orchis spectabilis
Juglans nigra	Ornithogalum bivale
Juniperus virginiana	Ornithogalum hirsutum
Kalmia angustifolia	Orobanche uniflora
Kalmia latifolia	Orobanche virginiana
Laurus aestivalis	Orontium aquaticum
Laurus benzoin	Osmunda virginiana
Laurus borbonia	Osteospermum uvedalia
Laurus sassafras	Oxalis stricta
Lechea minor	Oxalis violacea
Leontice thalictroides	Panax quinquefolius
Leontodon dandelion	Panax trifolius
Limodorum tuberosum	Panicum dichotomum
Linum virginianum	Panicum virgatum
Liriodendron tulipifera	Parthenium integrifolium
Lithospermum virginianum	Penthorum sedoides
Lobelia cardinalis	Phalaris oryzoides
Lobelia cliffortiana	Phlox glaberrima
Lonicera marilandica	Phryma leptostachya
Lonicera symphoricarpos	Pinus balsamea
Ludwigia alternifolia	Pinus taeda
Lupinus perennis	Plantago virginica

<i>Platanus occidentalis</i>	<i>Scutellaria hyssophifolia</i>
<i>Poa capillaris</i>	<i>Scutellaria integrifolia</i>
<i>Poa flava</i>	<i>Scutellaria lateriflora</i>
<i>Podophyllum peltatum</i>	<i>Senecio aureus</i>
<i>Polemonium dubium</i>	<i>Serratula glauca</i>
<i>Polygala cruciata</i>	<i>Serratula scariosa</i>
<i>Polygala senega</i>	<i>Serratula spicata</i>
<i>Polygala verticillata</i>	<i>Serratula squarrosa</i>
<i>Polygala viridescens</i>	<i>Sigesbeckia occidentalis</i>
<i>Polygonum sagittatum</i>	<i>Silene antirrhina</i>
<i>Polygonum scandens</i>	<i>Silene virginica</i>
<i>Polygonum virginianum</i>	<i>Silphium asteriscus</i>
<i>Populus heterophylla</i>	<i>Silphium helianthoides</i>
<i>Pontederia cordata</i>	<i>Sison canadense</i>
<i>Prenanthes alba</i>	<i>Sium rigidus</i>
<i>Prinos verticillatus</i>	<i>Smilax herbacea</i>
<i>Prunus virginiana</i>	<i>Smilax lanceolata</i>
<i>Pteris atropurpurea</i>	<i>Smilax laurifolia</i>
<i>Pulmonaria virginica</i>	<i>Smyrniium aureum</i>
<i>Pyrola maculata</i>	<i>Smyrniium integerrimum</i>
<i>Pyrus coronaria</i>	<i>Solidago canadensis</i>
<i>Quercus alba</i>	<i>Solidago sempervirens</i>
<i>Quercus nigra</i>	<i>Sonchus canadensis</i>
<i>Quercus phellos</i>	<i>Sophora tinctoria</i>
<i>Quercus prinus</i>	<i>Spiraea opulifolia</i>
<i>Quercus rubra</i>	<i>Spiraea trifoliata</i>
<i>Queria canadensis</i>	<i>Staphylea trifolia</i>
<i>Ranunculus abortivus</i>	<i>Stipa avenacea</i>
<i>Rhexia virginica</i>	<i>Swertia difformis</i>
<i>Rhinanthus virginica</i>	<i>Tetragonotheca helianthoides</i>
<i>Rhus copallinum</i>	<i>Teucrium canadense</i>
<i>Rhus glabra</i>	<i>Teucrium virginicum</i>
<i>Rhus radicans</i>	<i>Thapsia trifoliata</i>
<i>Rhus toxicodendron</i>	<i>Tilia americana</i>
<i>Robinia pseudoacacia</i>	<i>Tradescantia virginiana</i>
<i>Rudbeckia hirta</i>	<i>Trigopogon virginicum</i>
<i>Rudbeckia laciniata</i>	<i>Trichostema dichotomum</i>
<i>Rudbeckia oppositifolia</i>	<i>Trifolium biflorum</i>
<i>Rudbeckia purpurea</i>	<i>Trifolium reflexum</i>
<i>Rudbeckia triloba</i>	<i>Trillium sessile</i>
<i>Ruellia strepens</i>	<i>Triosteum angustifolium</i>
<i>Rumex britannica</i>	<i>Ulmus americana</i>
<i>Rumex verticillatus</i>	<i>Uniola paniculata</i>
<i>Salvia lyrata</i>	<i>Urtica pumila</i>
<i>Salvia urticifolia</i>	<i>Utricularia gibba</i>
<i>Sanguinaria canadensis</i>	<i>Utricularia subulata</i>
<i>Sanicula canadensis</i>	<i>Uvularia perfoliata</i>
<i>Sanicula marilandica</i>	<i>Vaccinium frondosum</i>
<i>Sarothra gentianoides</i>	<i>Vaccinium stamineum</i>
<i>Sarracenia flava</i>	<i>Valeriana locusta</i>
<i>Sarracenia purpurea</i>	var. <i>radiata</i>
<i>Satureja origanoides</i>	<i>Veratrum luteum</i>
<i>Satureja virginiana</i>	<i>Verbena nodiflora</i>
<i>Saururus cernuus</i>	<i>Verbena spuria</i>
<i>Saxifraga pensylvanica</i>	<i>Verbena urticifolia</i>
<i>Scandix procumbens</i>	<i>Verbesina virginica</i>
<i>Schwalbea americana</i>	<i>Veronica marilandica</i>
<i>Schoenus glomeratus</i>	<i>Veronica virginica</i>
<i>Scirpus capitatus</i>	<i>Viburnum acerifolium</i>
<i>Scrophularia marilandica</i>	<i>Viburnum nudum</i>

Viburnum prunifolium
 Viola palmata
 Viola pedata
 Yucca filamentosa

Hermann

Acer rubrum
 Ambrosia elatior
 Anemone virginiana
 Arum dracontium
 Asclepias purpurascens
 Asclepias tuberosa
 Aster dumosus
 Aster novae-angliae
 Aster novi-belgii
 Aster puniceus
 Aster undulatus
 Baccharis halimifolia
 Bidens bipinnata
 Campanula americana
 Cynanchum hirtum
 Eupatorium purpureum
 Geum virginianum
 Helianthus strumosus
 Heuchera americana
 Juglans nigra
 Liriodendron tulipifera
 Mespilus arbutifolia
 Mitella diphylla
 Napaea hermaphrodita
 Ranunculus abortivus
 Satureja virginiana
 Senecio hieracifolius
 Solidago flexicaulis
 Solidago rigida
 Solidago sempervirens
 Staphylea trifolia
 Verbena hastata

Linnaeus Hort. Cliff.

Acer negundo
 Ageratum altissimum
 Amaranthus hypocondriacus
 Amaryllis atamasca
 Ambrosia trifida
 Amorpha fruticosa
 Andromeda paniculata
 Angelica lucida
 Apocynum androsaemifolium
 Aralia racemosa
 Aralia spinosa
 Arethusa ophioglossoides
 Aristolochia arborescens
 Arum virginicum
 Asclepias incarnata
 Asclepias syriaca
 Asclepias tuberosa
 Ascyrum hypericoides
 Aster annuus
 Aster dumosus

Aster linifolius
 Aster novae-angliae
 Aster novi-belgii
 Aster puniceus
 Aster tradescantii
 Aster undulatus
 Azalea lutea
 Baccharis halimifolia
 Bartsia coccinea
 Bidens pilosa
 Bignonia capreolata
 Bignonia radicans
 Burmannia biflora
 Cacalia porophyllum
 Cactus pentagonus
 Campanula perfoliata
 Canna glauca
 Cassia marilandica
 Cassia nictitans
 Ceanothus americanus
 Cephalanthus occidentalis
 Cercis canadensis
 Chelone glabra
 Chionanthus virginica
 Chrysanthemum serotinum
 Chrysogonum virginianum
 Coix dactyloides
 Collinsonia canadensis
 Commelina communis
 Commelina erecta
 Convallaria racemosa
 Coreopsis lanceolata
 Cornus florida
 Cracca virginiana
 Crataegus coccinea
 Crotalaria alba
 Cupressus distichia
 Cynanchum hirtum
 Diodia virginiana
 Diospyros virginiana
 Dracocephalum virginianum
 Erythrina herbacea
 Eupatorium perfoliatum
 Eupatorium purpureum
 Eupatorium scandens
 Euonymus americanus
 Fumaria cucullaria
 Fumaria sempervirens
 Geum virginianum
 Gleditsia triacanthos
 Glycine apios
 Glycine frutescens
 Hedera quinquefolia
 Hedysarum canadense
 Hedysarum volubile
 Helenium autumnale
 Helianthus multiflorus
 Helianthus strumosus
 Heuchera americana
 Hibiscus moscheutos

Houstonia caerulea	Serratula squarrosa
Hydrophyllum virginianum	Silphium asteriscus
Hyssopus nepetoides	Sison canadense
Ilex cassine	Smilax lanceolata
Juglans nigra	Smyrnium aureum
Juniperus virginiana	Solanum carolinense
Jussiaea erecta	Solanum diphyllum
Laurus benzoin	Solanum tomentosum
Laurus borbonia	Solidago canadensis
Laurus sassafras	Solidago rigida
Liriodendron tulipifera	Spiraea hypericifolia
Lobelia cardinalis	Spiraea opulifolia
Lobelia cliffortiana	Spiraea trifoliata
Lobelia inflata	Staphylea trifolia
Lobelia siphilitica	Tradescantia virginiana
Lonicera symphoricarpos	Trichostema dichotomum
Ludwigia alternifolia	Uniola paniculata
Lycopodium alopecuroides	Uvularia perfoliata
Magnolia virginiana	Verbena nodiflora
Malva caroliniana	Verbena urticifolia
Menispermum canadense	Veronica virginica
Mespilus arbutifolia	
Mitella diphylla	Linnaeus Hort. Upsal.
Monarda didyma	Acer rubrum
Monarda fistulosa	Ageratum altissimum
Monarda punctata	Amaranthus hypocondriacus
Myrica asplenifolia	Ambrosia elatior
Myrica cerifera	Ambrosia trifida
Napaea hermaphrodita	Amorpha fruticosa
Nepeta virginica	Angelica lucida
Nyssa aquatica	Aquilegia canadensis
Obolaria virginica	Aralia racemosa
Onoclea sensibilis	Asclepias syriaca
Ophiorrhiza mitreola	Aster annuus
Osteospermum uvedalia	Aster novi-belgii
Parthenium integrifolium	Aster tradescantii
Phlox glaberrima	Bidens pilosa
Platanus occidentalis	Bignonia radicans
Podophyllum peltatum	Bupthalmum helianthoides
Polygonum sagittatum	Cacalia suaveolens
Polygonum virginianum	Campanula perfoliata
Populus balsamifera	Cassia marilandica
Pontederia cordata	Cassia nictitans
Prenanthes alba	Ceanothus americanus
Ptelea trifoliata	Cercis canadensis
Quercus prinus	Commelina communis
Quercus rubra	Commelina erecta
Rhus radicans	Coreopsis alternifolia
Robinia pseudoacacia	Coreopsis tripteris
Rubus odoratus	Cornus florida
Rudbeckia laciniata	Crataegus coccinea
Ruellia strepens	Cucubalus stellatus
Rumex sanguineus	Cupressus distichia
Sanguinaria canadensis	Erigeron camphoratum
Sanguisorba canadensis	Eupatorium altissimum
Sarracenia flava	Eupatorium perfoliatum
Sarracenia purpurea	Eupatorium scandens
Satureja virginiana	Eupatorium sessilifolium
Saururus cernuus	Euonymus americanus
Senecio hieracifolius	Fumaria sempervirens

Gleditsia triacanthos
Glycine apios
Helenium autumnale
Helianthus multiflorus
Hibiscus moscheutos
Hyssopus nepetoides
Juglans nigra
Jussiaea erecta
Liriodendron tulipifera
Lobelia cardinalis
Lobelia inflata
Ludwigia alternifolia
Malva caroliniana
Menispermum canadense
Monarda fistulosa
Monarda punctata
Myrica cerifera
Osteospermum uvedalia
Podophyllum peltatum
Ribes oxyacanthoides
Robinia pseudoacacia
Rubus odoratus
Rudbeckia laciniata
Rudbeckia triloba
Ruellia strepens
Rumex sanguineus
Sanicula marilandica
Saururus cernuus
Scrophularia marilandica
Senecio hieracifolius
Solidago altissima
Solidago canadensis
Spiraea hypericifolia
Spiraea trifoliata
Tradescantia virginiana
Verbena hastata
Verbena spuria
Verbena urticifolia

Linnaeus Amoen. Acad.

Bartsia coccinea
Cornus canadensis
Polygala cruciata
Polygala lutea
Polygala senega
Polygala verticillata
Polygala viridescens
Rhododendron maximum
Trillium erectum
Uvularia perfoliata
Viola canadensis

Linnaeus Materia Medica

Actaea racemosa
Aristolochia serpentaria
Ilex cassine
Laurus benzoin
Laurus sassafras
Panax quinquefolius
Polygala senega

Rhus copallinum
Rhus vernix
Rumex britannica
Rumex sanguineus

Linnaeus Nova Plantarum Genera

Agave virginica
Arethusa bulbosa
Arethusa ophioglossoides
Cracca virginiana
Elymus canadensis
Epigaea repens
Gaultheria procumbens
Gaura biennis
Helonias bullata
Holosteum succulentum
Kalmia angustifolia
Kalmia latifolia
Lechea major
Lechea minor
Mitchella repens
Mitella diphylla
Napaea dioica
Onoclea sensibilis
Orontium aquaticum
Phryma leptostachya
Sarothra gentianoides
Veratrum luteum

Linnaeus Virid. Cliffort.

Aralia racemosa
Aralia spinosa
Asclepias incarnata
Asclepias syriaca
Bignonia capreolata
Commelina communis
Commelina erecta
Hyssopus nepetoides
Monarda fistulosa
Rudbeckia laciniata
Solanum carolinense
Solanum diphyllum

Martyn

Aster grandiflorus
Buphthalmum helianthoides
Cassia marilandica
Coreopsis lanceolata
Crotalaria alba
Limodorum tuberosum
Malva caroliniana
Solidago altissima

Mitchell

Acnida cannabina
Callicarpa americana
Chelone penstemon
Cracca virginiana
Galax aphylla
Hamamelis virginiana

Hedera quinquefolia
 Itea virginica
 Mitchellia repens
 Onoclea sensibilis
 Phryma leptostachya
 Proserpinaca palustris
 Stewartia malacodendron

Morison

Acrostichum platyneuros
 Actaea spicata var. alba
 Adiantum pedatum
 Ageratum altissimum
 Alisma coridifolia
 Amaryllis atamasca
 Ambrosia artemisiifolia
 Ambrosia elatior
 Ambrosia trifida
 Anchusa virginiana
 Angelica lucida
 Apocynum androsaemifolium
 Apocynum cannabinum
 Aquilegia canadensis
 Aralia racemosa
 Arethusa ophioglossoides
 Aristolochia serpentaria
 Asarum canadense
 Asarum virginicum
 Aster annuus
 Aster cordifolius
 Aster linifolius
 Aster puniceus
 Aster tradescantii
 Aster undulatus
 Baccharis halimifolia
 Bartsia coccinea
 Bidens bipinnata
 Bidens frondosa
 Bignonia radicans
 Buphthalmum helianthoides
 Cacalia atriplicifolia
 Cacalia porophyllum
 Campanula perfoliata
 Carduus virginianus
 Cenchrus tribuloides
 Chrysanthemum serotinum
 Cicuta maculata
 Coix dactyloides
 Convallaria racemosa
 Convallaria stellata
 Coreopsis alternifolia
 Coreopsis auriculata
 Coreopsis tripteris
 Cynanchum hirtum
 Cyperus arundinacea
 Dracocephalum virginianum
 Elymus canadensis
 Eriophorum virginicum
 Eupatorium altissimum
 Eupatorium aromaticum

Eupatorium hyssopifolium
 Eupatorium perfoliatum
 Eupatorium purpureum
 Eupatorium rotundifolium
 Eupatorium sessilifolium
 Fumaria sempervirens
 Gentiana saponaria
 Glycine apios
 Gnaphalium obtusifolium
 Gnaphalium purpureum
 Hedysarum canadense
 Helenium autumnale
 Helianthus divaricatus
 Helianthus giganteus
 Helianthus strumosus
 Helonias bullata
 Hibiscus moscheutos
 Houstonia caerulea
 Hydrophyllum virginianum
 Hyssopus nepetoides
 Lilius canadense
 Lithospermum virginianum
 Lobelia cardinalis
 Lobelia siphilitica
 Lupinus perennis
 Lycopodium alopecuroides
 Monarda ciliata
 Monotropa uniflora
 Nepeta virginica
 Obolaria virginica
 Onoclea sensibilis
 Orchis ciliaris
 Orobanche virginiana
 Osteospermum uvedalia
 Oxalis stricta
 Panicum latifolium
 Plantago virginica
 Poa capillaris
 Polygonum virginianum
 Polypodium bulbiferum
 Polypodium virginianum
 Pontederia cordata
 Pulmonaria virginica
 Rudbeckia hirta
 Rudbeckia laciniata
 Rudbeckia purpurea
 Rudbeckia triloba
 Salvia lyrata
 Salvia urticifolia
 Sanguisorba canadensis
 Sarracenia flava
 Sarracenia purpurea
 Satureja origanoides
 Satureja virginiana
 Scandix procumbens
 Scutellaria lateriflora
 Senecio aureus
 Serratula spicata
 Serratula squarrosa
 Silphium trifoliatum

Sison canadense
 Sium rigidius
 Smyrnium aureum
 Solidago canadensis
 Solidago sempervirens
 Spiraea trifoliata
 Staphylea trifoliata
 Thalictrum cornutii
 Tradescantia virginiana
 Trillium erectum
 Uvularia perfoliata
 Verbena urticifolia
 Yucca filamentosa

Parkinson

Juglans alba
 Platanus occidentalis
 Thalictrum cornutii

Petiver

Acrostichum areolatum
 Asclepias verticillata
 Baccharis halimifolia
 Eriocaulon decangulare
 Glycine comosa
 Helianthus angustifolius
 Lysimachia quadrifolia
 Mitchella repens
 Myrica asplenifolia
 Ornithogalum hirsutum
 Osmunda cinnamomea
 Osmunda virginiana
 Poa flava
 Polypodium virginianum
 Polyplemum procumbens
 Pontederia cordata
 Pyrola maculata
 Rudbeckia purpurea
 Scutellaria hirsutifolia
 Trifolium comosum

Plukenet

Acer negundo
 Acer rubrum
 Acrostichum platyneuros
 Actaea racemosa
 Adiantum pedatum
 Aletris farinosa
 Amaryllis atamasca
 Ambrosia artemisiifolia
 Anchusa virginiana
 Andromeda mariana
 Andromeda paniculata
 Anemone quinquefolia
 Anemone thalictroides
 Apocynum cannabinum
 Arabis canadensis
 Aralia spinosa
 Arethusa bulbosa
 Arethusa ophioglossoides

Aristolochia arborescens
 Aristolochia serpentaria
 Arum dracontium
 Asarum virginicum
 Asclepias variegata
 Asclepias verticillata
 Ascyrum crux-andreae
 Ascyrum hypericoides
 Ascyrum villosum
 Aster divaricatus
 Aster dumosus
 Aster linariifolius
 Aster mutabilis
 Aster tenuifolius
 Azalea lutea
 Azalea viscosa
 Baccharis halimifolia
 Bartsia coccinea
 Betula nigra
 Bidens bipinnata
 Bignonia sempervirens
 Buphthalmum helianthoides
 Cacalia atriplicifolia
 Cacalia porophyllum
 Callicarpa americana
 Cardamine virginica
 Cassia nictitans
 Ceanothus americanus
 Celastrus bullatus
 Celosia paniculata
 Cenchrus tribuloides
 Cephalanthus occidentalis
 Chaerophyllum arborescens
 Chelone glabra
 Chelone hirsuta
 Chrysanthemum serotinum
 Chrysogonum virginianum
 Cicuta maculata
 Claytonia virginica
 Clethra alnifolia
 Coix dactyloides
 Convallaria racemosa
 Conyza asteroides
 Conyza bifrons
 var. flocculosa
 Conyza linifolia
 Coreopsis alternifolia
 Coreopsis auriculata
 Coreopsis verticillata
 Cornus florida
 Cracca virginiana
 Crataegus coccinea
 Cupressus distichia
 Cupressus thyoides
 Cyperus arundinacea
 Epigaea repens
 Eriocaulon decangulare
 Eriophorum virginicum
 Eryngium aquaticum
 Eupatorium aromaticum

Eupatorium coelestinum	Orobanche uniflora
Eupatorium hyssopifolium	Osteospermum uvedalia
Eupatorium perfoliatum	Oxalis violacea
Eupatorium rotundifolium	Panax trifolius
Eupatorium scandens	Parthenium integrifolium
Euphorbia maculata	Phlox divaricata
Euonymus americanus	Phlox ovata
Fagus pumila	Phlox paniculata
Fumaria cucullaria	Phlox pilosa
Galium bermudense	Phlox setacea
Gaura biennis	Phlox subulata
Gerardia flava	Pinus balsamea
Gerardia pedicularia	Pinus strobus
Gerardia purpurea	Pinus taeda
Gleditsia triacanthos	Plantago virginica
Gnaphalium plantaginifolium	Polygala cruciata
Hamamelis virginiana	Polygala lutea
Hedysarum canescens	Polygala verticillata
Hedysarum paniculatum	Polygonum arifolium
Helianthus giganteus	Polygonum sagittatum
Helonias bullata	Polygonum scandens
Heuchera americana	Polypodium bulbiferum
Hibiscus virginicus	Pontederia cordata
Hieracium venosum	Prenanthes alba
Horminum virginicum	Prenanthes altissima
Houstonia caerulea	Prunus virginiana
Houstonia purpurea	Ptelea trifoliata
Hypericum setosum	Pulmonaria virginica
Hyssopus nepetoides	Pyrola maculata
Ilex cassine	Quercus prinus
Iris verna	Quercus rubra
Juglans alba	Rhexia mariana
Jussiaea erecta	Rhexia virginica
Kalmia angustifolia	Rhus copallinum
Kalmia latifolia	Rhus vernix
Laurus benzoin	Ribes oxyacanthoides
Laurus indica	Robinia pseudoacacia
Laurus sassafras	Rudbeckia hirta
Liriodendron tulipifera	Rudbeckia purpurea
Ludwigia alternifolia	Rudbeckia triloba
Lysimachia quadrifolia	Sarothra gentianoides
Magnolia virginiana	Sarracenia flava
var. glauca	Sarracenia purpurea
Medeola virginiana	Satureja origanoides
Medicago virginica	Satureja virginiana
Menispermum canadense	Saururus cernuus
Mitchella repens	Saxifraga pensylvanica
Monarda ciliata	Schwalbea americana
Monarda punctata	Scirpus retrofractus
Monotropa uniflora	Scutellaria integrifolia
Morus rubra	Serratula glauca
Myosotis virginiana	Serratula praealta
Myrica asplenifolia	Serratula scariosa
Myrica cerifera	Serratula spicata
Nepeta virginica	Silphium helianthoides
Nyssa aquatica	Smilax bona-nox
Obolaria virginica	Smilax herbacea
Oldenlandia uniflora	Smilax lanceolata
Onoclea sensibilis	Smyrniium aureum
Ornithogalum hirsutum	Solanum diphyllum

<i>Solanum verbascifolium</i>	<i>Diospyros virginiana</i>
<i>Solanum virginianum</i>	<i>Dodecatheon meadia</i>
<i>Solidago flexicaulis</i>	<i>Elymus canadensis</i>
<i>Solidago latifolia</i>	<i>Eupatorium altissimum</i>
<i>Solidago sempervirens</i>	<i>Eupatorium hyssopifolium</i>
<i>Spermacoce tenuior</i>	<i>Eupatorium perfoliatum</i>
<i>Spiraea hypericifolia</i>	<i>Eupatorium purpureum</i>
<i>Spiraea trifoliata</i>	<i>Eupatorium sessilifolium</i>
<i>Thesium umbellatum</i>	<i>Eupatorium trifoliatum</i>
<i>Tilia americana</i>	<i>Euphorbia polygonifolia</i>
<i>Trifolium reflexum</i>	<i>Euonymus americanus</i>
<i>Trillium sessile</i>	<i>Fumaria cucullaria</i>
<i>Triosteum angustifolium</i>	<i>Galium bermudense</i>
<i>Uniola paniculata</i>	<i>Gerardia pedicularia</i>
<i>Urtica divaricata</i>	<i>Hedysarum canescens</i>
<i>Vaccinium hispidulum</i>	<i>Hedysarum marilandicum</i>
<i>Vaccinium stamineum</i>	<i>Helenium autumnale</i>
<i>Viburnum prunifolium</i>	<i>Houstonia caerulea</i>
<i>Viola palmata</i>	<i>Houstonia purpurea</i>
<i>Viola pedata</i>	<i>Juniperus virginiana</i>
<i>Vitis arborea</i>	<i>Lactuca canadensis</i>
<i>Vitis labrusca</i>	<i>Lechea minor</i>
<i>Vitis vulpina</i>	<i>Lonicera marilandica</i>
<i>Yucca filamentosa</i>	<i>Lycopodium alopecuroides</i>
Ray - Hist. Pl.	<i>Magnolia virginiana</i>
<i>Acrostichum platyneuros</i>	var. <i>glauca</i>
<i>Adiantum pedatum</i>	<i>Magnolia virginiana</i>
<i>Amaranthus hybridus</i>	var. <i>grisea</i>
<i>Ambrosia artemisiifolia</i>	<i>Mimulus ringens</i>
<i>Ambrosia elatior</i>	<i>Obolaria virginica</i>
<i>Ambrosia trifida</i>	<i>Orchis ciliaris</i>
<i>Aristolochia serpentaria</i>	<i>Orchis psychodes</i>
<i>Asclepias incarnata</i>	<i>Orobanche virginiana</i>
<i>Ascyrum crux-andreae</i>	<i>Osmunda cinnamomea</i>
<i>Ascyrum hypericoides</i>	<i>Phlox maculata</i>
<i>Aster linariifolius</i>	<i>Phlox pilosa</i>
<i>Baccharis halimifolia</i>	<i>Pinus balsamea</i>
<i>Bartsia coccinea</i>	<i>Pinus taeda</i>
<i>Betula nigra</i>	<i>Plantago virginica</i>
<i>Bignonia capreolata</i>	<i>Plantago incarnata</i>
<i>Bignonia sempervirens</i>	<i>Polygala lutea</i>
<i>Bupthalmum helianthoides</i>	<i>Polygala senega</i>
<i>Carduus virginianus</i>	<i>Polygala verticillata</i>
<i>Ceanothus americanus</i>	<i>Polygonum virginianum</i>
<i>Celtis occidentalis</i>	<i>Polypodium virginianum</i>
<i>Cenchrus tribuloides</i>	<i>Quercus nigra</i>
<i>Cercis canadensis</i>	<i>Quercus phellos</i>
<i>Chaerophyllum arborescens</i>	<i>Quercus prinus</i>
<i>Chelone glabra</i>	<i>Rhus toxicodendron</i>
<i>Chrysanthemum serotinum</i>	<i>Robinia pseudoacacia</i>
<i>Chrysogonum virginianum</i>	<i>Rubus occidentalis</i>
<i>Cicuta bulbifera</i>	<i>Rudbeckia hirta</i>
<i>Coreopsis alternifolia</i>	<i>Rudbeckia purpurea</i>
<i>Coreopsis auriculata</i>	<i>Sanguinaria canadensis</i>
<i>Coreopsis tripteris</i>	<i>Sanicula marilandica</i>
<i>Cucubalus stellatus</i>	<i>Scrophularia marilandica</i>
<i>Cyperus odoratus</i>	<i>Scutellaria integrifolia</i>
<i>Dioscorea villosa</i>	<i>Senecio aureus</i>
<i>Diospyros virginiana</i>	<i>Silphium trifoliatum</i>
	<i>Smilax herbacea</i>

Smilax lanceolata
 Smyrniun aureum
 Solidago caesia
 Spiraea trifoliata
 Trichostema dichotomum
 Vaccinium hispidulum
 Valeriana locusta
 var. radiata
 Vitis vulpina

Ray - Banister, 1688

Chelone hirsuta
 Clematis viorna
 Gerardia flava
 Gerardia purpurea
 Hieracium venosum
 Napaea dioica
 Serratula scariosa
 Serratula spicata

Ray - Petiver Hort. Sicc.

Asclepias decumbens
 Chionanthus virginica
 Clematis viorna
 Clitoria mariana
 Polygala lutea
 Trichostema dichotomum
 Trifolium biflorum
 Viola pedata

Royen

Acer negundo
 Adiantum pedatum
 Amaryllis atamasca
 Ambrosia trifida
 Amorpha fruticosa
 Andropogon virginicum
 Anemone virginiana
 Angelica atropurpurea
 Angelica lucida
 Apocynum androsaemifolium
 Aralia racemosa
 Aralia spinosa
 Arum dracontium
 Asclepias incarnata
 Asclepias syriaca
 Asclepias tuberosa
 Aster annuus
 Aster cordifolius
 Aster dumosus
 Aster ericoides
 Aster grandiflorus
 Aster linifolius
 Aster novae-angliae
 Aster puniceus
 Aster tradescantii
 Aster undulatus
 Astragalus carolinianus
 Baccharis halimifolia
 Bidens nivea

Bignonia capreolata
 Bignonia sempervirens
 Cactus pentagonus
 Campanula americana
 Campanula perfoliata
 Cassia marilandica
 Ceanothus americanus
 Cephalanthus occidentalis
 Cercis canadensis
 Chionanthus virginica
 Chrysanthemum serotinum
 Coix dactyloides
 Commelina communis
 Commelina erecta
 Convallaria racemosa
 Coreopsis alternifolia
 Coreopsis lanceolata
 Coreopsis tripteris
 Cornus florida
 Crataegus coccinea
 Crotalaria alba
 Cupressus distichia
 Cynanchum hirtum
 Cyperus odoratus
 Diospyros virginiana
 Dracocephalum virginianum
 Erythrina herbacea
 Eupatorium perfoliatum
 Eupatorium purpureum
 Eupatorium scandens
 Fagus pumila
 Fraxinus americana
 Fumaria cucullaria
 Fumaria sempervirens
 Geranium carolinianum
 Glycine apios
 Glycine frutescens
 Gnaphalium purpureum
 Gomphrena interrupta
 Hedera quinquefolia
 Hedysarum canadense
 Hedysarum canescens
 Hedysarum volubile
 Helenium autumnale
 Helianthus multiflorus
 Helianthus strumosus
 Heuchera americana
 Hibiscus moscheutos
 Hydrophyllum virginianum
 Hyssopus nepetoides
 Ilex cassine
 Juglans nigra
 Juniperus virginiana
 Laurus benoin
 Laurus borbonia
 Laurus sassafras
 Limodorum tuberosum
 Liriodendron tulipifera
 Lobelia cardinalis
 Lobelia inflata

Lobelia siphilitica
 Lonicera symphoricarpos
 Ludwigia alternifolia
 Lupinus perennis
 Magnolia virginica
 Mespilus arbutifolia
 Mitella diphylla
 Monarda fistulosa
 Monarda punctata
 Myrica cerifera
 Napaea hermaphrodita
 Orchis ciliaris
 Ornithogalum hirsutum
 Osteosperma uvedalia
 Phaseolus helvulus
 Phlox glaberrima
 Phryma leptostachya
 Platanus occidentalis
 Podophyllum peltatum
 Polygonum sagittatum
 Polygonum virginianum
 Polypodium bulbiferum
 Pontederia cordata
 Populus balsamifera
 Prunus virginiana
 Quercus prinus
 Quercus rubra
 Ranunculus abortivus
 Rhus copallinum
 Rhus radicans
 Rhus vernix
 Robinia pseudoacacia
 Rubus odoratus
 Rudbeckia laciniata
 Ruellia strepens
 Rumex sanguineus
 Sanguisorba canadensis
 Satureja virginiana
 Saururus cernuus
 Scutellaria lateriflora
 Senecio hieracifolius
 Serratula squarrosa
 Silene antirrhina
 Silphium asteriscus
 Silphium trifoliatum
 Sison canadense
 Smyrnum aureum
 Solanum carolinense
 Solanum diphyllum
 Solanum tomentosum
 Solanum virginianum
 Solidago canadensis
 Solidago flexicaulis
 Solidago rigida
 Sonchus floridanus
 Spermocoe tenuior
 Spiraea hypericifolia
 Spiraea opulifolia
 Spiraea trifoliata
 Staphylea trifoliata

Thalictrum cornutii
 Tradescantia virginiana
 Trichostema dichotomum
 Trifolium comosum
 Uniola paniculata
 Uvularia perfoliata
 Verbena hastata
 Verbena nodiflora
 Verbena urticifolia
 Veronica virginica

Tournefort

Amaranthus hypocondriacus
 Aster novae-angliae
 Astragalus canadensis
 Bidens frondosa
 Chelone glabra
 Circaea lutetiana
 var. canadensis
 Lactuca canadensis
 Liliun canadense
 Lobelia kalmii
 Menispermum canadense
 Mentzelia aspera
 Oenothera fruticosa
 Oxalis stricta
 Polygonum scandens
 Rhus toxicodendron
 Teucrium canadense

Vaillant

Alsima cordifolia
 Bidens nivea
 Coreopsis verticillata
 Panax quinquefolius
 Panax trifolius
 Prenanthes alba
 Sanicula canadensis
 Sonchus floridanus
 Viburnum prunifolium

Synonymies for Some of the Vascular Plants Reported for Temperate North America in Linnaeus' First Edition of Species Plantarum

Acrostichum areolatum =
 Woodwardia areolata
 Acrostichum platyneuros =
 Asplenium platyneuron
 Acrostichum polypodioides =
 Polypodium polypodioides
 Actaea racemosa =
 Cimicifuga racemosa
 Actaea spicata var. alba =
 Actaea pachypoda
 Agrostis virginica =
 Sporobolus virginicus
 Alisma cordifolia =
 Echinodorus cordifolius

- Alisma subulata =
 Sagittaria subulata
 Alsine media =
 Stellaria media
 Amaryllis atamasca =
 Zephyranthes atamasca
 Ambrosia elatior =
 Ambrosia artemisiifolia
 var. elatior
 Ammannia ramosior =
 Rotala ramosior
 Andromeda arborea =
 Oxydendrum arboreum
 Andromeda calyculata =
 Chamaedaphne calyculata
 Andromeda mariana =
 Lyonia mariana
 Andromeda paniculata =
 Leucothoe racemosa
 Andromeda racemosa =
 Leucothoe racemosa
 Andropogon alopecuroides =
 Erianthus alopecuroides
 Andropogon divaricatum =
 Erianthus alopecuroides
 Andropogon hirtum =
 Hyparrhenia hirta
 Andropogon nutans =
 Sorghastrum nutans
 Anemone hepatica =
 Hepatica nobilis
 Anemone thalictroides =
 Anemonella thalictroides
 Annona triloba =
 Asimina triloba
 Antirrhinum canadense =
 Linaria canadensis
 Antirrhinum elatine =
 Kickxia elatine
 Aphanes arvensis =
 Alchemilla arvensis
 Arbutus uva-ursi =
 Arctostaphylos uva-ursi
 Arenaria rubra var. marina
 = Spergularia marina
 Arethusa divaricata =
 Cleistes divaricata
 Arethusa ophioglossoides =
 Pogonia ophioglossoides
 Arenica maritima =
 Senecio pseudoarnica
 Arum dracontium =
 Arisaema dracontium
 Arum triphyllum =
 Arisaema triphyllum
 Arum virginicum =
 Peltandra virginica
 Arundo phragmites =
 Phragmites australis
 Asclepias amoena =
 Asclepias purpurascens
 Asclepias decumbens =
 Asclepias tuberosa
 Ascyrum crux-andreae =
 Hypericum crux-andreae
 Ascyrum hypericoides =
 Hypericum hypericoides
 Ascyrum villosum =
 Hypericum setosum
 Aster annuus =
 Erigeron annuus
 Aster vernus =
 Erigeron vernus
 Avena pensylvanica =
 Trisetum pensylvanicum
 Avena spicata =
 Trisetum spicatum
 Azalea lutea =
 Rhododendron nudiflorum
 Azalea viscosa =
 Rhododendron viscosum
 Baccharis foetida =
 Pluchea foetida
 Bartsia coccinea =
 Castilleja coccinea
 Betonica annua =
 Stachys annua
 Bidens nivea =
 Melanthera nivea
 Bignonia catalpa =
 Catalpa bignonioides
 Bignonia crucigera =
 Bignonia capreolata
 Bignonia radicans =
 Campsis radicans
 Bignonia sempervirens =
 Gelsemium sempervirens
 Briza eragrostis =
 Eragrostis cilianensis
 Bunias cakile =
 Cakile maritima
 Bupthallum frutescens =
 Borrichia frutescens
 Bupthallum helianthoides =
 Kallias helianthoides
 Cacalia atriplicifolia =
 Arnoglossum atriplici-
 folium
 Cacalia porophyllum =
 Porophyllum ruderale
 Cacalia suaveolens =
 Hasteola suaveolens
 Cactus opuntia =
 Opuntia vulgaris
 Cactus pentagonus =
 Cereus pentagonus
 Campanula perfoliata =
 Triodanis perfoliata
 Cardamine virginica =
 Sibara virginica

Carduus altissimus =	Cracca virginiana =
Cirsium altissimum	Tephrosia virginiana
Carduus virginianus =	Crotalaria alba =
Cirsium virginianum	Baptisia alba
Carpinus ostrya =	Crotalaria perfoliata =
Ostrya virginiana	Baptisia perfoliata
Celastrus bullatus =	Cucubalus stellatus =
Celastrus scandens	Silene stellata
Celosia paniculata =	Cupressus distichia =
Iresine paniculata	Taxodium distichium
Chaerophyllum arborescens =	Cupressus thysoides =
Aralia spinosa	Chamaecyparis thyoidea
Chelidonium glaucium =	Cynanchum suberosum =
Glaucium flavum	Matelea suberosa
Chelone hirsuta =	Cynosurus aegyptius =
Penstemon hirsutus	Dactyloctenium aegyptium
Chelone penstemon =	Cyperus arundinacea
Penstemon laevigatus	Dulichium arundinaceum
Chenopodium anthelminticum =	Dactylis cynosuroides =
Chenopodium ambrosioides	Spartina cynosuroides
Chironia angularis =	Datisca hirta =
Sabatia angularis	Rhus hirta
Chironia campanulata =	Dianthera americana =
Sabatia campanulata	Justicia americana
Chironia dodecandra =	Dolichos polystachyus =
Sabatia dodecandra	Phaseolus polystachyus
Chrysocoma graminifolia =	Dolichos regularis =
Euthamia graminifolia	Galactia regularis
Cissampelos smilacina =	Dracocephalum virginianum =
Menispermum canadense	Physostegia virginiana
Cistus canadensis =	Dracontium foetidum =
Helianthemum canadense	Symplocarpus foetidus
Clinopodium incanum =	Elymus hystrix =
Pycnanthemum incanum	Hystrix patula
Clitoria virginiana =	Erigeron camphoratus =
Centrosema virginiana	Pluchea camphorata
Coix dactyloides =	Erigeron canadense =
Tripsacum dactyloides	Conyza canadensis
Convallaria polygonatum =	Eupatorium altissimum =
Polygonatum officinale	Ageratina altissima
Convallaria racemosa =	Eupatorium aromaticum =
Smilacina racemosa	Ageratina aromatica
Convallaria stellata =	Eupatorium coelestinum =
Smilacina stellata	Conoclinium coelestinum
Convolvulus carolinus =	Eupatorium scandens =
Ipomoea trichocarpa	Mikania scandens
Convolvulus panduratus =	Eupatorium trifoliatum =
Ipomoea pandurata	Eupatorium purpureum
Convolvulus repens =	Euphorbia portulacoides =
Calystegia sepium	Euphorbia ipecacuanhae
Convolvulus spithameae =	Fagus pumila =
Calystegia spithameae	Castanea pumila
Conyza asteroides =	Ferula canadensis =
Sericocarpus asteroides	Ligusticum canadense
Conyza linifolia =	Fumaria cucullaria =
Sericocarpus linifolius	Dicentra cucullaria
Coreopsis alternifolia =	Fumaria sempervirens =
Verbesina alternifolia	Corydalis sempervirens
Coreopsis angustifolia =	Gentiana quinquefolia =
Helianthus angustifolia	Gentianeella quinquefolia

Gerardia flava =	Helleborus trifolius =
Aureolaria flava	Coptis trifolia
Gerardia pedicularia =	Hibiscus palustris =
Aureolaria pedicularia	Hibiscus moscheutos
Gerardia purpurea =	Hibiscus virginicus =
Agalinis purpurea	Kosteletzkya virginica
Glycine apios =	Hippophae canadensis =
Apios americana	Shepherdia canadensis
Glycine bracteata =	Holcus laxus =
Amphicarpa bracteata	Uniola laxa
Glycine comosa =	Holcus striatus =
Amphicarpa bracteata	Sacciolepis striata
Glycine frutescens =	Holosteam succulentum =
Wisteria frutescens	Honkenya peploides
Glycine tomentosa =	Horminum virginicum =
Rhynchosia tomentosa	Salvia lyrata
Gnaphalium margaritaceum =	Hyoseris virginica =
Anaphalis margaritacea	Krigia virginica
Gnaphalium plantaginifolium =	Hypericum lasianthus =
Antennaria plantagini-	Gordonia lasianthus
folia	Hyssopus nepetoides =
Gnaphalium purpureum =	Agastache nepetoides
Gamochaeta purpurea	Ipomoea nyctelea =
Gomphrena interrupta =	Ellisianyctelea
Froelichia interrupta	Ipomoea tamnifolia =
Gratiola dubia =	Jacquemontia tamnifolia
Lindernia dubia	Juglans alba =
Guilandina dioica =	Carya tomentosa
Gymnocladus dioica	Juncus campestris =
Hedera quinquefolia =	Luzula campestris
Parthenocissus quinque-	Jussiaea erecta =
folia	Ludwigia erecta
Hedysarum canadense =	Laurus aestivalis =
Desmodium canadense	Litsea aestivalis
Hedysarum canescens =	Laurus benzoin =
Desmodium canescens	Lindera benzoin
Hedysarum frutescens =	Laurus borbonia =
Lespedeza violacea	Persea borbonia
Hedysarum hirtum =	Laurus indica =
Lespedeza hirta	Persea indica
Hedysarum marilandicum =	Laurus sassafras =
Desmodium marilandicum	Sassafras albidum
Hedysarum nudiflorum =	Laurus winterana =
Desmodium nudiflorum	Canella winterana
Hedysarum paniculatum =	Lechea major =
Desmodium paniculatum	Helianthemum canadense
Hedysarum repens =	Leontice thalictroides =
Lespedeza repens	Caulophyllum thalictro-
Hedysarum violaceum =	ides
Lespedeza violacea	Leontodon dandelion =
Hedysarum virginicum =	Krigia dandelion
Lespedeza virginica	Lilium camschatcense =
Hedysarum viridiflorum =	Fritillaria camschatc-
Desmodium viridiflorum	ensis
Hedysarum volubile =	Limodorum tuberosum =
Galactia volubilis	Calopogon tuberosus
Helianthus laevis =	Liquidambar peregrina =
Bidens laevis	Comptonia peregrina
Helianthus multiflorus =	Lithospermum virginianum =
Helianthus decapetalus	Onosmodium virginianum

Lonicera marilandica =	Spigelia marilandica	Cynoctonum mitreola
Lonicera symphoricarpos =	Symphoricarpos orbiculata	Ophrys lilifolia =
Lycopodium apodum =	Selaginella apoda	Liparis linifolia
Lycopsis virginica =	Myosotis verna	Orchis ciliaris =
Lythrum petiolatum =	Cuphea petiolata	Platanthera ciliaris
Magnolia virginiana	var. acuminata =	Orchis flava =
Magnolia virginiana	Magnolia acuminata	Platanthera flava
Magnolia virginiana	var. foetida =	Orchis psychodes =
Magnolia virginiana	Magnolia grandiflora	Platanthera psychodes
Magnolia virginiana	var. glauca =	Orchis spectabilis =
Magnolia virginiana	Magnolia virginiana	Galearis spectabilis
Magnolia virginiana	var. grisea =	Ornithogalum bivale =
Magnolia virginiana	Magnolia virginiana	Allium bivale
Magnolia virginiana	var. tripetala =	Ornithogalum hirsutum
Magnolia virginiana	Magnolia tripetala	Hypoxis hirsuta
Malva caroliniana =	Modiola caroliniana	Orbanche virginiana =
Medicago virginica =	Lespedeza virginica	Epifagus virginiana
Melissa nepeta =	Calamintha nepeta	Osmunda virginiana =
Melissa pulegioides =	Hedeoma pulegioides	Botrychium virginianum
Menispermum carolinum =	Cocculus carolinus	Osteospermum uvedalia =
Menispermum virginicum =	Cocculus carolinus	Smallanthus uvedalia
Mentha spicata var. viridis =	Mentha spicata	Oxalis longiflora =
Mespilus arbutifolia =	Aronia arbutifolia	Oxalis violacea
Mespilus canadensis =	Amelanchier canadensis	Pancreatium carolinianum =
Monarda ciliata =	Blephilia ciliata	Hymenocallis caroliniana
Monotropa hypopithys =	Hypopithys monotropa	Panicum clandestinum =
Myosotis virginiana	Hackelia virginiana	Dichanthelium clandestinum
Myrica asplenifolia =	Comptonia cerifera	Panicum crusgalli =
Myrica cerifera =	Comptonia cerifera	Echinochloa crusgalli
Napaea hermaphrodita =	Sida hermaphrodita	Panicum dichotomum =
Nymphaea lutea =	Nuphar luteum	Dichanthelium dichotomum
Nymphaea nelumbo =	Nelumbo lutea (ours)	Panicum dissectum =
Ophiorrhiza mitreola =		Paspalum dissectum
		Panicum filiforme =
		Digitaria filiformis
		Panicum glaucum =
		Setaria glauca
		Panicum italicum =
		Setaria italica
		Panicum latifolium =
		Dichanthelium latifolium
		Panicum sanguinale =
		Digitaria sanguinalis
		Phalaris oryzoides =
		Leersia oryzoides
		Phaseolus helvulus =
		Strophostyles helvola
		Phlox ovata =
		Ruellia caroliniensis
		Phlox setacea =
		Phlox subulata var. setacea
		Pinus balsamea =
		Abies balsamea
		Pistacia simaruba =
		Bursera simaruba
		Poa capillaris =
		Eragrostis capillaris
		Poa flava =

- Tridens flavus =
 Podophyllum diphyllum =
 Jeffersonia diphylla
 Polemonium dubium =
 Phacelia dubia
 Polygala viridescens =
 Polygala sanguinea
 Polygonum articulatum =
 Polygonella articulata
 Polypodium bulbiferum =
 Cystopteris bulbifera
 Polypodium lonchitis =
 Polystichum lonchitis
 Polypodium marginale =
 Dryopteris marginalis
 Polypodium noveboracense =
 Thelypteris noveboracensis
 Polypodium phegopteris =
 Thelypteris phegopteris
 Prinos glaber =
 Ilex glabra
 Prinos verticillatus =
 Ilex verticillata
 Pteris atropurpurea =
 Pellaea atropurpurea
 Pulmonaria virginica =
 Mertensia virginica
 Pyrus coronaria =
 Malus coronaria
 Queria canadensis =
 Paronychia canadensis
 Renealmia usneoides =
 Tillandsia usneoides
 Rhinanthus virginica =
 Aureolaria virginica
 Rhus radicans =
 Toxicodendron radicans
 Rhus toxicodendron =
 Toxicodendron quercifolia
 Rhus vernix =
 Toxicodendron vernix
 Rudbeckia oppositifolia =
 Kalliashelianthoides
 Rudbeckia purpurea =
 Echinacea purpurea
 Sagina virginica =
 Bartonia virginica
 Salsola prostrata =
 Kochia prostrata
 Sarothra gentianoides =
 Hypericum gentianoides
 Satureja origanoides =
 Cunila origanoides
 Satureja virginiana =
 Pycnanthemum virginianum
 Scandix cerefolium =
 Anthriscus cerefolium
 Scandix procumbens =
 Chaerophyllum procumbens
 Schoenus glomeratus =
 Rhyschospora glomerata
 Scirpus retrofractus =
 Cyperus retrofractus
 Scirpus spadicus =
 Fimbristylis spadicea
 Scutellaria hyssopifolia =
 Scutellaria integrifolia
 Senecio hieracifolius =
 Erechtites hieracifolia
 Serratula glauca =
 Vernonia glauca
 Serratula noveboracensis =
 Vernonia noveboracensis
 Serratula praealta =
 Vernonia noveboracensis
 Serratula scariosa =
 Liatris scariosa
 Serratula spicata =
 Liatris spicata
 Serratula squarrosa =
 Liatris squarrosa
 Sida abutilon =
 Abutilon theophrastii
 Sida crispa =
 Malva crispa
 Sigesbeckia occidentalis =
 Verbesina occidentalis
 Silphium helianthoides =
 Kalliashelianthoides
 Silphium solidaginoides =
 Kalliashelianthoides
 Sison canadense =
 Cryptotaenia canadensis
 Sisymbrium nasturtium-aquaticum =
 Nasturtium officinale
 Sium rigidius =
 Oxypholis rigidus
 Smilax caduca =
 Smilax rotundifolia
 Smyrnum aureum =
 Zizia aurea
 Smyrnum integerrimum =
 Taenidia integerrima
 Solidago altissima =
 Solidago canadensis
 Solidago lateriflora =
 Aster lateriflorus
 Solidago latifolia =
 Solidagoflexicaulis
 Sonchus floridanus =
 Lactuca floridana
 Sophora tinctoria =
 Baptisia tinctoria
 Spiraea aruncus =
 Aruncus sylvester
 Spiraea opulifolia =
 Physocarpus opulifolius
 Spiraea trifoliata =

Porteranthus trifoliatus
 Statice armeria =
 Armeria maritima
 Swertia difformis =
 Sabatia difformis
 Teucrium chamaepitys =
 Ajuga chamaepitys
 Teucrium virginicum =
 Teucrium canadense
 Thapsia trifoliata =
 Thaspium trifoliatum
 Thesium umbellatum =
 Comandra umbellata
 Tragopogon virginicum =
 Krigia virginica
 Trifolium biflorum =
 Stylosanthes biflora
 Uniola spicata =
 Distichlis spicata
 Urtica canadensis =
 Laportea canadensis
 Urtica cylindrica =
 Boehmeria cylindrica
 Urtica divaricata =
 Laportea canadensis
 Urtica pumila =
 Pilea pumila
 Vaccinium frondosum =
 Gaylussacia frondosa
 Vaccinium hispidulum =
 Gaultheria hispidula
 Vaccinium ligustrinum =
 Lyonia ligustrina
 Vaccinium mucronatum =
 Nemopanthus mucronata
 Valeriana locusta var. radiata
 = Valerianella radiata
 Veratrum luteum =
 Chamaelirium luteum
 Verbena nodiflora =
 Lippa nodiflora
 Verbena spuria =
 Verbena officinalis
 Verbesina alba =
 Eclipta alba
 Veronica marilandica =
 Polypremum procumbens
 Veronica virginica =
 Veronicastrum virginicum
 Viscum rubrum =
 Phoradendron rubrum
 Viscum terrestre =
 Lysimachia terrestris
 Vitis arborea =
 Ampelopsis arborea
 Vitis laciniosa =
 Vitis vulpina

From the raw data above, it can be noted that Linnaeus

mentions 889 species from temperate North America. Of that number 13 are varieties leaving a total of 876 species. Two of the varieties, *Magnolia virginiana* var. *glauca* and *Solanum nigrum* var. *vulgare*, can be considered equivalent to the modern declarations of var. *virginiana* and var. *nigrum* in each instance. One species, *Ornithogalum canadense*, is based solely upon South African material and Linnaeus misnamed it. Most likely he meant "capense".

Linnaeus gives habitat references for nearly all of his species. In the above review, 108 were referred to "America". This was variously defined by Linnaeus to include North America or the Americas, meaning both North and South America. Those species which are strictly stated by Linnaeus to be only from South America (but have synonymies based on North American elements) are not included. Canada is given for 162 species, while 80 (including five varieties of *Magnolia virginiana*) are given as occurring in the Carolinas. Maryland is credited with a total of 18 species with Pennsylvania (or Philadelphia) given for 44 species. Linnaeus assigns four species to Florida, two to Mississippi, five to New York, and one each to New England and New Jersey. The largest number of species are referred to Virginia. He refers 421 species and varieties to Virginia.

To be sure Linnaeus often referred a species to more than a single area, and thus the total is greater than the number of species considered, by Linnaeus, to be strictly from these areas. Even so, fully 47% of the species considered by Linnaeus to be from temperate North America (even if not stated as such) were considered to occur in Virgi-

nia. Only 18% of the plants were mentioned as being from Canada, while 12% came from America.

A number of temperate North American species were mentioned in Linnaeus' first edition of *Species plantarum* in synonymy, but the geographic range given by him did not include an area within temperate North America. Linnaeus gives the range of 47 species as occurring somewhere in the New World but not specifically in the temperate region. For 86 species Linnaeus gives the range as occurring in both temperate North America and somewhere else in the World. This usually refers to species which Linnaeus believed occurred in both Europe and the Americas. In 79 instances, synonyms based on temperate North American species are cited, but the geographical range does not even mention the New World or temperate North America. Mostly, Linnaeus is referring to weed species gathered in temperate North America prior to 1753.

Linnaeus named 14 species for America, 37 for Canada, 10 for the Carolinas, nine for Maryland, four for New York, five for Pennsylvania or Philadelphia, and 63 for Virginia. He named one species each for Florida and New England.

Linnaeus named some North American species for early explorers. Three species were named for Peter Kalm (*Hieracium kalmii*, *Hypericum kalmianum* and *Lobelia kalmii*), and one each for Clayton (*Osmunda claytoniana*) and Tradescant (*Aster tradescantii*). He did name species for others, of which three may be mentioned: *Hieracium gronovii*, *Lobelia cliffortiana* and *Thalictrum cornutii*.

EVALUATION OF SPECIES PUBLISHED IN SPECIES PLANTARUM

In attempting to evaluate the contents of Carl Linnaeus' first edition of *Species plantarum* for those species of vascular plants he attributes to temperate North America it is necessary to consider the conditions under which Linnaeus labored, and his abilities to resolve problems long associated with efforts to produce a world's flora. To begin, it is useful to start with the status of the collections available to him, and then turn to the literature already published which Linnaeus had access to. As shall become obvious, the two are not entirely mutually inclusive.

Over Linnaeus' life time he gathered a herbarium of some 16,000 specimens. Today specimens once in his possession are located in England, Sweden and France. The gathering of plants with the intention of making permanent collections of dried specimens began in the early 1500s and was a well established practice when Linnaeus began his own botanical explorations on an expedition to Lapland in 1732. The older herbaria consisted largely of bound volumes, or *hortus siccus*, into which specimens were glued in a predetermined order. Typically the plant specimens were arranged according to some system of classification, according to geographical location, or even randomly. A few workers maintained their collections loose between sheets of brown paper. In either cases, notations as to collector, place of origin, or even the scientific name may or may not be affixed to the specimen. In the better collections, such data were retained. Unfortunately, few such collections exist. As

the herbarium developed, more and more workers realized the inherent difficulties of the inflexibility of bound volumes and moved to individual sheets. As workers had done with the bound volumes, often several species were placed on a single sheet, but in time, a single specimen was placed on a sheet so as to allow absolute freedom in rearranging the entire collection as the newer systems of classifications were developed.

Most of the herbaria made prior to the Linnaean era were arranged in bound volumes. Linnaeus departed from this practice, as did Sherard and Dillenius for example. Unlike his Oxford colleagues who placed one or more collections on the same sheet, Linnaeus usually glued a single collection to a sheet. None of these men was consistent in their annotations, although Sherard, Dillenius and Gronovius in Holland were far more consistent in this practice than Linnaeus. While most workers wrote onto a label who collected each specimen, and from where it came, this was not at all a consistent practice. Dale and Du Bois were far more consistent in this practice than most. Petiver, in spite of his chaotic collections, maintained good records but apparently was of the practice of giving away specimens. To record what he had seen he would add the collector and his data to preexisting labels compounding the identification of many his specimens. Sloane, Ray, and Plukenet were not at all consistent in their herbarium skills. Sloane kept only the briefest records of his own collections, and retained almost none for the specimens he received from others. Ray failed to retain nearly all of the data he might have received and was constantly rear-

ranging his collections to fit into his latest scheme of classification. As for Plukenet, he kept some records, but for the most part, his bound volumes of dried plants are remarkably void of collecting data. This is somewhat overcome by the identifications associated with each specimen (when annotated) as these can be used to trace a collection to a particular collector.

For Linnaeus, few of the extant collections made by other botanists prior to 1753 were consulted. He examined the collection made by Joachim Burser, but these were only of Old World species. Linnaeus also examined some of the collections in the Sherardian and Dillenian herbaria at Oxford University, but the extent to which he reviewed these holdings is unknown. Linnaeus did not see the vast holdings owned by Sir Hans Sloane when he visited England in 1736. Thus, the rich array of specimens gathered by Plukenet, Petiver, Sloane and many others were not studied by him. Likewise, the large Catesby collections ultimately obtained by Sloane were not examined in London although Linnaeus certainly saw duplicates sent to Sherard and Dillenius at Oxford. Linnaeus states that he saw the collections of Tournefort, Vaillant and Jussieu in France, but it is unlikely that he spent much time evaluating their contents. Linnaeus also visited Phillip Miller in London and Peter Collinson, but it is not known to what extent Linnaeus examined their gardens or reviewed their herbaria. Most likely, any such examination was highly superfluous. However, in many instances, Linnaeus received specimens from these men, or exchanged plant collections with them so that Linnaeus had access to many of their more unique species.

During the Linnaean years abroad in Holland he worked closely with three men and was able to examine not only their libraries and gardens, but in some cases, exotic collections of dried herbarium material. With the assistance of George Clifford, Linnaeus was able to study the Clifford garden in great detail, make herbarium specimens, and to prepare an illustrated volume treating the species in and around the Clifford estate. This book was published in 1738. The specimens remained with Clifford and eventually came to the British Museum (Natural History) in London where they may be examined today. Thus, when Linnaeus was preparing *Species plantarum* in 1751 and 1752, he did not have direct access to these specimens.

Linnaeus also aided Adriaan van Royen, a professor of botany at Leyden. His book, *Florae leydenensis prodromus*, was published in 1740. It is unclear what role Linnaeus played in the production of this work, but it is certain that Linnaeus must have aided him in the assignment of synonymies and the literature he cites.

The third man was Gronovius, an amateur botanist, with a large library and herbarium. In assisting Gronovius with his *Flora virginica*, Linnaeus was able to examine many of the temperate North American collections of John Clayton. It is likely that Linnaeus saw most of the species reported in the first part of that work (published in 1739), but it is uncertain to what extent Linnaeus examined Clayton material cited in the second part (published in 1743). Gronovius sent a number of Clayton specimens to Linnaeus, and these are extant in the Linnaean Herbarium in London.

Significant collections of vascular plants came to Linnaeus from a number of sources. For temperate North America the most significant holdings available to Linnaeus while writing *Species plantarum* were those provided by Peter Kalm. Linnaeus heavily relied upon these specimens to determine species circumscriptions. In some cases, the Virginia specimens Linnaeus examined earlier proved to be somewhat different from the specimens found by Kalm, and Linnaeus was not in a position to differentiate between them inasmuch as he no longer had access to Clayton material.

Of the two collectors, Clayton and Kalm, it can be seen from the listings given above that Linnaeus cited John Clayton material more frequently than he did that gathered by Peter Kalm. In part this is due to the greater number of species found by Clayton compared to Kalm, but an additional factor is that many of the Kalm collections in the Linnaean Herbarium in London likely are not identified as such and cannot be presently assigned to him. Continued studies of the herbarium holdings available to Linnaeus will likely result in an increase in the number of Kalm specimens. This can be accomplished, in part, by examining the Kalm collections found elsewhere. It is interesting to note that when one compares the Kalm species with those cited by Linnaeus as originating from Canada or Pennsylvania there is a close relationship. In some instances, the Canada reference is based on a Cornuti polynomial or some other name that clearly establishes Canada as the source of the name.

In a number of instances, Linnaeus saw both Clayton and Kalm material of plants which

he took to represent the same species. For the most part, Linnaeus cites both Virginia and Canada as the sources of the species, but in some instances he gives only Virginia. In the following listing, species of temperate North American vascular plants are given which the available evidence indicates were collected by both men. In addition, the geographic source(s) of the species is given only as they are applicable to those parts of temperate North America where the two men botanized (A= America; C= Canada; NY= New York; P= Pennsylvania; V= Virginia; N= none of the above; X= no geographic data given):

- Acalypha virginica* - V
Acer negundo - V
Acer rubrum - P, V
Adiantum pedatum - C, V
Agrostis virginica - V
Andromeda mariana - V
Andromeda paniculata - V
Andropogon nutans - V
Anemone thalictroides - C, V
Anemone virginiana - V
Angelica sylvestris - N
Antirrhinum canadense - C, V
Apocynum cannabinum - C, V
Arabis canadensis - A
Arabis lyrata - C
Aralia spinosa - V
Arenaria rubra var. *marina* - N
Arethusa bulbosa - C, V
Arethusa ophioglossoides - C, V
Aristolochia serpentaria - V
Asarum canadense - C
Asclepias nivea - V
Aster concolor - V
Astragalus canadensis - C, V
Azalea viscosa - V
Bartsia coccinea - NY, V
Briza eragrostis - N
Cacalia atriplicifolia - C, V
Ceanothus americanus - V
Celtis occidentalis - V
Centurus tribuloides - V
Cephalanthus occidentalis - A
Cercis canadensis - V
Chaerophyllum arborescens - V
Chelone glabra - C, V
Cicuta bulbifera - C, V
Cicuta maculata - V
Claytonia virginica - V
Convolvulus spithameus - V
Cornus florida - V
Cracca virginiana - C, V
Crataegus coccinea - C, V
Crataegus crus-galli - X
Crataegus tomentosa - V
Crotalaria sagittalis - V
Cucubalus stellatus - C, V
Cuscuta americana - V
Dianthera americana - V
Diospyros virginiana - A
Dracontium foetidum - V
Euphorbia corollata - C, V
Euphorbia ipecacuanhae - C, V
Euphorbia polygonifolia - C, V
Euonymus americanus - V
Fagus pumila - A
Geranium carolinianum - V
Gerardia flava - C, V
Gerardia pedicularia - C, V
Gerardia purpurea - C, V
Gleditsia triacanthos - V
Glycine apios - V
Glycine bracteata - V
Glycine comosa - V
Gnaphalium obtusifolium - P, V
Gnaphalium plantaginifolium - V
Gnaphalium purpureum - P, V
Gratiola virginiana - V
Hamamelis virginiana - V
Hedera quinquefolia - C
Hedysarum hirtum - V
Hedysarum violaceum - V
Heuchera americana - V
Hibiscus palustris - C, V
Hieracium gronovii - P, V
Houstonia caerulea - V
Houstonia purpurea - V
Hydrangea arborescens - V
Hydrocotyle umbellata - A
Hyoseris virginica - V
Hypericum canadense - C
Hypericum mutilum - C, V
Iris verna - V
Juglans alba - V
Laurus sassafras - V
Limodorum tuberosum - A
Liquidambar styraciflua - V
Liriodendron tulipifera - A
Lobelia cliffortiana - C, V
Ludwigia alternifolia - V
Lupinus perennis - V
Medeola virginiana - V
Melanthium virginicum - V
Melissa pulegioides - C, V
Menispermum canadense - C, V
Mespilus canadensis - C, V

Mollugo verticillata - V
 Monotropa uniflora - C, V
 Nyssa aquatica - A
 Oenothera fruticosa - V
 Ophrys lilifolia - C, V
 Orchis psychodes - C
 Ornithogalum hirsutum - C, V
 Orobanche virginiana - V
 Osmunda regalis - V
 Osmunda virginiana - A
 Oxalis stricta - V
 Oxalis violacea - C, V
 Panax quinquefolius - C, P, V
 Panicum dichotomum - V
 Pinus taeda - C, V
 Plantago virginica - V
 Poa capillaris - C, V
 Polygala verticillata - V
 Polygala viridescens - V
 Pontedaria cordata - V
 Prenanthes alba - P, V
 Prinos verticillatus - V
 Pteris atropurpurea - V
 Pulmonaria virginica - V
 Pyrola maculata - A
 Quercus alba - V
 Quercus nigra - A
 Quercus phellos - A
 Quercus prinus - A
 Quercus rubra - V
 Ranunculus abortivus - C, V
 Rhexia virginica - V
 Rhus glabra - A
 Rhus radicans - C, V
 Rhus toxicodendron - C, V
 Rhus vernix - A
 Salvia lyrata - V
 Sanguinaria canadensis - A
 Sarothra gentianoides - P, V
 Satureja origanoides - V
 Saxifraga nivalis - C, V
 Saxifraga pennsylvanica - C, P,
 V
 Schoenus glomeratus - V
 Scirpus capitatus - V
 Scutellaria hyssopifolia - V
 Scutellaria integrifolia - C,
 V
 Serratula spicata - A
 Silene virginica - V
 Sison canadense - A
 Sisyrinchium bermudiana - V
 Smilax pseudo-china - V
 Smyrniolum aureum - A
 Solidago canadensis - C, V
 Sonchus canadensis - C
 Sophora tinctoria - V
 Spiraea opulifolia - C, V
 Spiraea trifoliata - C, V
 Thapsia trifoliata - V

Tragopogon virginicum - C, V
 Trifolium biflorum - C, V
 Ulmus americana - V
 Uniola spicata - A
 Urtica cylindrica - C, V
 Urtica pumila - C
 Utricularia subulata - V
 Uvularia perfoliata - C, V
 Vaccinium frondosum - A
 Vaccinium stamineum - A
 Veratrum luteum - C, V
 Viburnum acerifolium - V
 Viburnum prunifolium - C, V
 Viola pedata - V
 Vitis vinifera - N
 Xyris indica - N

As can be seen from the above summary, Linnaeus had both Clayton and Kalm material for 170 species, of which 8 were referred only to Canada, 79 only to Virginia, and only 22 to America. A total of 47 species were referred to Canada and Virginia, 7 to Pennsylvania and Virginia, and one to New York and Virginia. As Kalm material came into Linnaeus' possession well after he had formulated many of his opinions regarding species, it is not unusual to see a debate in the literature over the selection of a Clayton specimen or a Kalm collection as a suitable lectotype. This cannot be resolved at this point, but it is well to consider that while in many instances Linnaeus does not give Canada or Pennsylvania as the only location for a combination Clayton-Kalm species, he does give many species with Virginia as the only location. It is possible, therefore, that Linnaeus had already established the definition of the species on material other than the Kalm collection in his herbarium. Thus, while he used Kalm specimens to assist him in his characterization, he likely did not base his circumscription initially upon that Kalm collection.

When Linnaeus wrote *Species plantarum* in 1753, some

species from temperate North America were described based only upon a single reference and/or collection. Of these, the majority are, as one would suspect, Clayton or Kalm specimens. The following breakdown is based solely upon single author or collector references in *Species plantarum*. In those instances where there is a specimen in the Linnaean Herbarium in London that seems to be different from the author of the synonym or the collector mentioned in the reference given by Linnaeus, a star (*) is given. The reader is urged to consult *Species plantarum* for the places of publication.

Bauhin

Othonna cineraria

Catesby

Annona glabra
*Annona triloba**
Bignonia caerulea
Cissampelos smilacina
Ipomoea carolina
Magnolia virginiana
 var. *tripetala*
Pancratium carolinianum
Philadelphus inodorus
Sloanea emarginata
*Smilax tannoides**
*Viscum purpureum**
Viscum rubrum

Clayton

*Agrostis virginica**
Alisma subulata
Amaranthus graecizans
Ammannia ramosa
Andropogon divaricatum
*Antirrhinum canadense**
*Arabis lyrata**
Asclepias rubra
*Aster concolor**
Aster rigidus
Aster vernus
*Betula lenta**
Buchnera americana
Chironia dodecandra
Convolvulus panduratus
*Convolvulus spithameus**
Coreopsis angustifolia
*Crateagus tomentosa**
Crateagus viridis

*Cynoglossum virginianum**
*Dactylis cynosuroides**
*Dianthera americana**
*Dirca palustris**
Dolichos polystachyus
Dolichos regularis
Elephantopus tomentosus
Elymus hystrix
Ferula canadensis
*Glycine bracteata**
Gratiola dubia
*Gratiola virginiana**
Hedysarum frutescens
*Hedysarum hirtum**
Hedysarum nudiflorum
*Hedysarum repens**
*Hedysarum violaceum**
*Hedysarum virginicum**
Hedysarum viridiflorum
Helianthus laevis
*Hieracium gronovii**
Holcus laxus
Holcus striatus
*Hydrangea arborescens**
*Hyoseris virginica**
*Hypericum mutilum**
*Ipomoea nyctelea**
*Iris virginica**
Laurus aestivalis
Leontodon dandelion
*Linum virginianum**
Lycopsis virginica
*Lycopus virginicus**
Lythrum lineare
Lythrum petiolatum
*Lythrum verticillatum**
*Melanthium virginicum**
Melissa pulegioides
*Mespilus canadensis**
*Monarda clinopodia**
Orchis flava
Orchis spectabilis
Ornithogalum bivale
Osmunda claytoniana
*Panicum dichotomum**
Panicum virgatum
*Phalaris oryzoides**
Polemonium dubium
*Prinos verticillatus**
*Pteris atropurpurea**
*Pyrus coronaria**
*Queria canadensis**
*Rhinanthus virginica**
Rudbeckia oppositifolia
Rumex verticillatus
Sagina virginica
Salicornia virginica
*Schoenus glomeratus**
*Scirpus capitatus**
Sigesbeckia occidentalis

- Silene virginica**
Silphium solidaginoides
Smyrniun integerrimum
*Sonchus canadensis**
Stipa avenacea
Swertia difformis
Teucrium virginicum
*Thapsia trifoliata**
*Tragopogon virginicum**
*Ulmus americana**
*Uniola spicata**
*Urtica pumila**
*Utricularia gibba**
Utricularia subulata
*Vaccinium frondosum**
Verbesina virginica
Veronica marilandica
*Viburnum acerifolium**
Viburnum nudum
*Viola primulifolia**
*Xyris indica**
- Collinson**
*Podophyllum diphyllum**
- Colden**
*Uvularia sessilifolia**
- Dillenius**
Asclepias amoena
Aster miser
Carduus altissimus
*Chenopodium anthelminticum**
*Clematis crispa**
Convolvulus carolinus
Crotalaria perfoliata
Ipomoea lacunosa
Ipomoea tamnifolia
*Lycopodium apodum**
Lycopodium carolinianum
*Lycopodium obscurum**
*Polemonium rubrum**
*Rosa carolina**
Ruellia biflora
Solanum nigrum
 var. *virginicum*
*Trichostema brachiatum**
Triosteum perfoliatum
- Kalm**
Acer pennsylvanicum
Acer saccharinum
Allium canadense
Amaranthus retroflexus
Andromeda racemosa
Aster laevis
Avena pennsylvanica
Avena spicata
Bromus ciliatus
Carex folliculata
- Carex squarrosa*
Chenopodium virginicum
Chironia angularis
Chironia campanulata
Chrysocoma graminifolia
Cinna arundinacea
Cistus canadensis
Dalibarda repens
Datisca hirta
Erigeron philadelphicum
Galium tinctorium
Galium trifidum
Gentiana quinquefolia
Gentiana villosa
Helianthus decapetalus
Hieracium kalmii
Hieracium paniculatum
Hippophae canadensis
Hordeum jubatum
Hydrocotyle americana
Hypericum canadense
Hypericum kalmianum
Liquidambar peregrina
Mentha canadensis
Ophrys cernua
Panicum filiforme
Polygala sanguinea
Polygonum articulatum
Polygonum erectum
Polygonum pensylvanicum
Polypodium marginale
Polypodium noveboracense
Potentilla canadensis
Prinos glaber
Ribes cynosbati
Rubus canadensis
Rubus hispidus
Sambucus canadensis
Senecio canadensis
Smilax caduca
Smilax rotundifolia
Solidago lateriflora
Thalictrum dioicum
Urtica capitata
Vaccinium album
Vaccinium corymbosum
Vaccinium ligustrinum
Vaccinium mucronatum
Viburnum lentago
Viscum terrestre
- Linnaeus**
Cacalia suaveolens
Gomphrena serrata
Guilandina dioica
Helleborus trifolius
Lechea major
Lilium camtschatcensis
Menispermum carolinum
*Oxalis longiflora**

Physalis pruinosa
*Polymnia canadensis**
Rumex persicarioides
Salsola prostrata
Solidago noveboracensis
*Swertia corniculata**
Thalictrum purpurascens
Viburnum dentatum
*Viola canadensis**
*Viola lanceolata**

Mitchell

*Chelone penstemon**

Morison

*Helianthus divaricatus**

Petiver

Clitoria mariana

Plukenet

*Anemone quinquefolia**
Ascyrum villosum
Aster mutabilis
*Aster tenuifolius**
*Conyza asteroides**
Conyza bifrons var. *flosculosa*
Conyza linifolia
*Cupressus thyioides**
Euphorbia maculata
*Morus rubra**
*Phlox divaricata**
*Phlox ovata**
*Phlox paniculata**
Phlox setacea
Phlox subulata
Rhexia mariana
*Scirpus retrofractus**
*Solidago latifolia**
Spiraea tomentosa
*Thesium umbellatum**
*Urtica divaricata**

Plumier

Silphium laciniatum

Ray

Magnolia virginiana
 var. *grisea*
*Phlox maculata**
*Polygala incarnata**

Tournefort

*Lobelia kalmii**

van Royen

Bidens bullata
Gomphrena interrupta

Of the some 889 species

credited to temperate North America by Linnaeus in the first edition of *Species plantarum*, 241, or about 27% were based on a single reference. In giving a single reference, Clayton material is mentioned for 100 species and Kalm for 60 species. For 49 of the 100 Clayton species, Linnaeus had other specimens. In nearly all instances, the additional material was a Kalm collection which is now found in Linnaeus' herbarium in London.

For the remaining names based on a single reference, the majority were established on illustrations in books published by the authors mentioned above. Linnaeus credits Catesby as the basis of 12 of his species. In three instances Linnaeus had other material in his herbarium to augment the illustrations published by Catesby. As for Dillenius, who Linnaeus mentions as his only reference for 18 species, he had other material for seven of those species. Plukenet is mentioned as Linnaeus' only source of information for 21 species. Of that number, however, Linnaeus had additional material in only 11 instances. All of these authors illustrated their works, true, but it is significant that of these particular authors -- Catesby, Dillenius and Plukenet -- Linnaeus may well have been able to examine actual specimens as well as the illustrations. Collections at the University of Oxford are extant with many of the polynomials mentioned by Linnaeus in synonymy, and it is likely that Linnaeus was able to associate specific names with certain species only as a result of his examination of the Sherardian and Dellinian herbaria at Oxford.

It is significant that by the late 1740s when Mitchell published his work on new

genera that Linnaeus had additional information relative to all but one. Mitchell is cited 13 times in the first edition of *Species plantarum*, but only once is he the sole source of information. This is the case of *Chelone penstemon*, a plant Mitchell proposed to recognize as a new genus, *Penstemon*. Even so, Linnaeus had other specimens at hand and did not need to rely only upon Mitchell's publications. As for van Royen who Linnaeus assisted, he gives him as the only reference in two species; in neither case did Linnaeus have supplementary material in his herbarium.

As already noted, Linnaeus was unable to see the original herbarium material gathered by a number of workers. Thus, he had to rely upon the illustrations prepared by these workers for the characterization and identification of many species.

For his study of temperate North American species, the work of Gronovius proved particularly significant. Not only did Linnaeus examine the collections made by Clayton which were described in the 1739 part of *Flora virginica*, but he aided Gronovius in the determination of synonyms. Additionally, Linnaeus received many subsequent specimens from Gronovius of Clayton's Virginia discoveries, and these too were evaluated by him and synonymies determined. Of specific interest is the relationship between the Gronovius flora and those works on American botany published prior to 1739.

In the following listing, Gronovius citations given by Linnaeus in the first edition of *Species plantarum* are reviewed, and the synonyms cited by both Gronovius and Linnaeus are listed for each species.

The following abbreviations are used: BR = Breynius; CA = Catesby; CM = Commelinus; CO = Cornuti; D = Dillenius; H = Hermann; LA = Lafitau; L = Linnaeus' *Hortus cliffortianus*; LT = Leat; LV = Linnaeus' *Viridarium cliffortianum*; MA = Martyn; MO = Morison; PE = Petiver; PL = Plukenet; PU = Plumier; RA = Ray; RO = Royen; S = Sloane; T = Tourenfort; V = Vaillant. Only those species considered by Linnaeus to be strictly New World will be evaluated. A star (*) indicates when Gronovius and the additional reference(s) cited by Linnaeus are the only synonyms given in the first edition of *Species plantarum*. Those works published after 1739 or 1743, depending on the page in Gronovius, are not considered in this review.

Acer negundo - L
Acer rubrum - CA, PL
Acrostichum areolatum - PE*
Acrostichum platyneuros - PL
Actaea racemosa - D
Adiantum pedatum - CO
Ageratum altissimum - L
Aletris farinosa - PL*
Amaryllis atamasca - MO
Ambrosia trifida - L
Anchusa virginiana - MO, PL*
Andromeda arborea - CA*
Andromeda mariana - PL*
Andromeda paniculata - L, PL
Andropogon virginicum - RO*
Anemone virginiana - RO
Annona muricata - L, PL
Apocynum cannabinum - PL
Aquilegia canadensis - CO
Aralia spinosa - L, LV
Arethusa bulbosa - PL*
Arethusa divaricata - CA*
Arethusa ophioglossoides, L
 PL
Aristolochia serpentaria - CA,
 PL
Arum triphyllum - MO
Arum virginicum - L*
Asarum canadense - CO
Asarum virginicum - PL
Asclepias decumbens - PE*
Asclepias incarnata - CO, L
Asclepias nivea - D
Asclepias syriaca - CO, L
Asclepias verticillata - PE,

- PL*
Ascyrum hypericoides - PL,
 RA
Aster divaricatus - PL*
Aster dumosus - L
Aster ericoides - D*
Aster grandiflorus - D
Aster novae-angliae - H, L*
Azalea lutea - L
Baccharis halimifolia - L
Bartsia coccinea - L, MO
Bignonia radicans - CA, L
Buphthalmum frutescens - CA,
 D, L
Buphthalmum helianthoides - PL
Burmanna biflora - L*
Cacalia atriplicifolia - PL
Callicarpa americana - CA
Campanula perfoliata - L, MO*
Carduus virginianus - MO
Cassia chamaecrista - BR, L
Cassia ligustrina - D, L
Ceanothus americanus - CM, L
Celosia paniculata - S
Celtis occidentalis - RA*
Cenchrus tribuloides - MO
Cephalanthus occidentalis - L
Cercis canadensis - L
Chelone glabra - L, T
Chelone hirsuta - PL, RA*
Chionanthus virginica - L
Chrysogonum virginianum - L
Cicuta bulbifera - RA*
Cicuta maculata - MO, PL*
Claytonia virginica - PL
Clematis viorna - D
Clitoria virginiana - D*
Commelina communis - L
Convallaria racemosa - L
Convolvulus repens - PU*
Coreopsis auriculata - PL
Coreopsis verticillata - PL,
 V*
Cornus florida - L
Cracca virginiana - L, PL*
Crataegus coccinea - L
Crotalaria sagittalis - L, PL
Cucubalus stellatus - RA*
Cupressus distichia - CA, L
Cuscuta americana - S*
Cynanchum suberosum - D, L
Cyperus arundinacea - MO, PL*
Cyperus odoratus - RO
Diodia virginiana - L*
Dioscorea villosa - PL*
Diospyros virginiana - CA, L
Epigaea repens - PL
Erigeron canadense - L, MO
Eriocaulon decangulare - PE,
 PL*
Eriophorum virginicum - MO,
 PL*
Eryngium aquaticum - PL*
Eryngium foetidum - L
Eupatorium aromaticum - PL
Eupatorium coelestinum - D
Eupatorium perfoliatum - L
Eupatorium purpureum - CO
Eupatorium scandens - L, PL*
Euphorbia polygonifolia - RA*
Euonymus americanus - PL
Fagus pumila - CA, PL*
Fraxinus americana - CA*
Fumaria cucullaria - L
Galium bermudense - PL, RA*
Gentiana saponaria - CA, MO*
Geranium carolinianum - D*
Geranium maculatum - D*
Gerardia flava - RA
Gerardia pedicularia - PL
Gerardia purpurea - PL, RA*
Geum virginianum - H, L*
Gleditsia triacanthos - L
Glycine apios - CO, L
Glycine comosa - PE*
Glycine tomentosa - D*
Gnaphalium plantaginifolium -
 PL*
Gnaphalium purpureum - D, RO
Hamamelis virginiana - PL*
Hedera quinquefolia - CO, L
Hedysarum canescens - L
Hedysarum paniculatum - PL*
Helenium autumnale - L, PL
Helianthus angustifolius - PE*
Helianthus atrorubens - D*
Heuchera americana - L
Hibiscus moscheutos - L
Hieracium venosum - PL, RA*
Houstonia caerulea - L
Houstonia purpurea - RA
Hydrophyllum virginianum - L
Hyssopus nepetoides - L, LV
Iris verna - PL*
Iva frutescens - L
Juglans nigra - L
Juniperus virginiana - L
Kalmia angustifolia - PL*
Kalmia latifolia - PL
Laurus benzoin - L
Laurus borbonia - CA, L*
Laurus indica - L
Laurus sassafras - L, PL
Lechea minor - RA*
Ligusticum scothieum - L
Limodorum tuberosum - MA*
Liquidambar styraciflua - CA,
 L
Liriodendron tulipifera - H, L
Lithospermum virginianum - MO*

- Lobelia cardinalis* - L
Lobelia cliffortiana - L
Lonicera marilandica - CA, RA*
Lonicera symphoricarpos - D, L
Ludwigia alternifolia - L
Lupinus perennis - MO, RO*
Lycopodium alopecuroides - L,
 MO
Lysimachia quadrifolia - PE
Magnolia virginiana - L*
Magnolia virginiana var.
acuminata - CA*
Magnolia virginiana var.
foetida - CA
Medeola virginiana - PL*
Medicago virginica - PL*
Melothria pendula - L
Menispermum canadense - L
Menispermum virginicum - D*
Mitchella repens - CA, PE,
 PL*
Mollugo verticillata - PL
Monarda ciliata - MO, PL*
Monarda punctata - L
Monotropa uniflora - CA, MO
Myosotis virginiana - PL*
Myrica asplenifolia - L
Myrica cerifera - CA, L, PL*
Nepeta virginica - L
Nyssa aquatica - CA, L
Obolaria virginica - L, PL
Oldenlandia uniflora - PL*
Oncoclea sensibilis - L, PL
Ophiorrhiza mitreola - L*
Orchis ciliaris - RO
Orobanche uniflora - PL*
Orobanche virginiana - MO, RA*
Osteospermum uvedalia - L
Oxalis stricta - T
Panax quinquefolius - LA, V
Panax trifolius - PL, V*
Panicum capillare - S*
Parthenium integrifolium - D,
 L
Passiflora lutea - L
Phlox glaberrima - D, L*
Physalis viscosa - D, L*
Phytolacca americana - L
Pinus balsamea - PL
Pinus taeda - PL
Plantago virginica - MO, PL,
 RA*
Platanus occidentalis - CA, L
Poa flava - PE*
Podophyllum peltatum - CA, L
Polygala senega - RA*
Polygala verticillata, PL,
 RA*
Polygonum sagittatum - L, LA
Polygonum virginianum - L
Pontederia cordata - L
Prenanthes alba - L
Pulmonaria virginica - MO, PL*
Pyrola maculata - PL
Quercus alba - CA*
Quercus nigra - CA, RA*
Quercus phellos - CA, RA*
Quercus prinus - CA, L
Quercus rubra - CA, PL
Ranunculus abortivus - H
Renalmia usneoides - L, PE, S
Rhus copallinum - PL
Rhus glabra - D, L
Rhus radicans - D, L
Rhus toxicodendron - T
Rhus vernix - D, L
Robinia pseudoacacia - L
Rudbeckia laciniata - CO, L,
 LV
Rudbeckia purpurea - CA, PL
Ruellia strepens - D, L*
Salvia urticifolia - MO*
Sanguinaria canadensis D, L
Sanicula canadensis - T
Sanicula marilandica - RA
Sarothra gentianoides - PL*
Sarracenia flava - CA, L
Sarracenia purpurea - CA, L
Satureja origanoides - MO, PL*
Satureja virginiana - H, L
Saururus cernuus - L
Saxifraga pensylvanica - D
Scandix procumbens - MO*
Schwalbea americana - PL*
Scirpus spadicus - S*
Scrophularia marilandica - RA*
Scutellaria hyssopifolia - PE*
Scutellaria integrifolia - PL,
 RA*
Scutellaria lateriflora - MO*
Senecio aureus - MO, RA*
Serratula glauca - D
Serratula spicata - D
Serratula squarrosa - D
Silene antirrhina - D*
Silphium asteriscus - D, L*
Silphium helianthoides - PL*
Sison canadense - L
Sisyrinchium bermudiana - D, L
Sium rigidius - MO*
Smilax lanceolata - L
Smilax laurifolia - CA*
Smilax pseudo-china - L
Smilax sarsaparilla - L
Smyrniolum aureum - PL
Solidago canadensis - L
Solidago sempervirens - H
Sophora tinctoria - PL*
Spiraea opulifolia - L
Spiraea trifoliata - L, PL

Tetragonotheca helianthoides - D*
 Tiarella cordifolia - L
 Tradescantia virginiana - L
 Trifolium reflexum - PL*
 Trillium sessile - CA, PL*
 Vaccinium stamineum - PL*
 Valeriana locusta var. radiata - RA*
 Verbena urticifolia - L
 Verbesina alba - D, L
 Veronica virginica - L*
 Viola palmata - PL*
 Viola pedata - PE, PL*
 Yucca filamentosa - MO, PL*

When Linnaeus proposed species in the first edition of *Species plantarum* he used, to a significant degree, information already published by others who proceeded him. During an examination of the Linnaean references relating to the State of Maryland published in the first two editions of his book, it was discovered that Linnaeus used the herbarium collections at Oxford to augment his understanding of temperate North American botany to a degree not previously realized. The knowledge which Linnaeus gained at Oxford was almost immediately expressed in Gronovius' *Flora virginica* which Linnaeus worked on soon after his departure from England in the late summer of 1736. This is probably equally true for regions in North America other than Maryland. It is not the purpose of this review to attempt a combined examination of this problem using both Linnaeus' published works and the Sherardian and Dillenian herbaria at the University of Oxford. However, the following list may prove helpful for those who are investigating potential lectotypes for species which might be found at Oxford. This in no way should be construed as a statement of affirmation that such authentic material is actually present. It should be remember that other Linnaean material

is located in Sweden and in the British Museum (Natural History) in London. These collections are not considered in this listing. Many of these species are based, in part, upon Clayton material. Authentic Clayton specimens are to be sought at the British Museum (Natural History).

A star (*) is added to those species which are also based on Clayton collections cited by Gronovius.

Acrostichum areolatum*
 Acrostichum platyneuros*
 Actaea spicata var. alba
 Andromeda arborea*
 Annona glabra
 Annona triloba
 Asarum virginicum*
 Asclepias amoena
 Asclepias decumbens*
 Asclepias variegata
 Ascyrum villosum
 Aster miser
 Aster mutabilis
 Astragalus carolinianus
 Baccharis foetida*
 Bignonia caerulea
 Bignonia sempervirens
 Carduus altissimus
 Carduus virginianus*
 Cissampelos smilacina
 Clinopodium rugosum
 Clitoria mariana
 Clitoria virginiana*
 Convallaria stellata
 Convolvulus carolinus
 Conyza bifrons var. flosculosa
 Conyza linifolia
 Coreopsis auriculata*
 Cyperus arundinacea*
 Dioscorea villosa*
 Eupatorium hyssopifolium
 Euphorbia maculata
 Galium bermudense*
 Glycine tomentosa*
 Helianthus angustifolius*
 Hieracium venosum*
 Horminum virginicum
 Hypericum setosum*
 Ipomoea carolina
 Ipomoea lacunosa
 Ipomoea tannifolia
 Iris versicolor
 Lonicera marilandica*
 Lycopodium carolinianum

Magnolia virginiana
 var. *acuminata**
Magnolia virginiana
 var. *foetida**
Magnolia virginiana
 var. *grisea*
Magnolia virginiana
 var. *tripetala*
*Medicago virginica**
*Monarda ciliata**
*Orobancha uniflora**
Pancreatium carolinianum
Phaseolus helvulus
Philadelphus inodorus
Phlox pilosa
Phlox setacea
Pistacia simaruba
*Poa flava**
Polemonium rubrum
*Polygala cruciata**
Polypodium virginianum
*Prenanthes alba**
Rhexia mariana
Ribes oxycanthoides
*Sanicula marilandica**
*Scandix procumbens**
*Schwalbea americana**
*Serratula glauca**
*Silphium helianthoides**
Sloanea emarginata
Smilax bona-nox
Solanum nigrum
 var. *virginicum*
Solidago caesia
Thalictrum cornutii
*Tilia americana**
Trifolium cosmosum
*Triosteum angustifolium**
Triosteum perfoliatum
Valeriana locusta
 var. *radiata**
Viscum rubrum
*Yucca filamentosa**

Of the 81 entities that might have authentic material found in the Sherardian and Dillenian herbaria at the University of Oxford, 33 are also based on Clayton specimens. Three major pre-Linnaean authors have major sets of their collections at Oxford. Of the three, Catesby, Dillenius and Morison, Catesby and Dillenius are the only authors or herbaria Linnaeus seems to have consulted (both of these men have their collections in the Sherardian Herbarium). The

following 12 species cited by Linnaeus might have voucher specimens at Oxford to go with the illustrations published by Catesby:

Annona glabra
Annona triloba
Bignonia caerulea
Bignonia sempervirens
Cissampelos smilacina
Ipomoea carolina
Magnolia virginiana
 var. *tripetala*
Pancreatium carolinianum
Philadelphus inodorus
Pistacia simaruba
Sloanea emarginata
Viscus rubrum

Dillenius specimens should be consulted for the following species which should be considered in addition to the published illustrations when proposing lectotypes. A total of 19 species are listed:

Asclepias amoena
Asclepias variegata
Aster miser
Astragalus carolinianus
Carduus altissimus
Clinopodium rugosum
Convolvulus carolinus
Eupatorium hyssopifolium
Horminum virginicum
Ipomoea lacunosa
Ipomoea tamnifolia
Iris versicolor
Lycopodium carolinianum
Phaseolus helvulus
Polemonium rubrum
Ribes oxycanthoides
Solidago caesia
Solanum nigrum
 var. *virginicum*
Triosteum perfoliatum

Morison (and in the case of most temperate North American species, this includes Bobart) the following species should be considered in addition to the published illustrations. A total of 5 species are listed below.

Actaea spicata var. *alba*
Convallaria stellata

Eupatorium hyssopifolium
 Polypodium virginianum
 Thalictrum cornutii

Other authors occasionally sent specimens to Sherard and Dillenius, or these two men annotated specimens they received from various naturalists, with polynomials that could have been observed when Linnaeus examined the Oxford collections. Polynomials published by Ray, Plukenet and Petiver are on such specimens. The following Linnaean species have polynomials proposed by these men given in synonymy, and such names may be on specimens found in the Sherardian and Dillenian herbaria at the University of Oxford. PE= Petiver; PL= Plukenet; R= Ray.

Asclepias variegata - PL
 Ascyrum villosum - PL
 Aster mutabilis - PL
 Bignonia sempervirens - PL, R
 Clinopodium rugosum - PL, R
 Clitoria mariana - PE
 Conyza bifrons var. floscula
 - PL
 Conyza linifolia - PL
 Eupatorium hyssopifolium -
 PL, R
 Euphorbia maculata - PL
 Horminum virginicum - PL
 Magnolia virginiana
 var. grisea - R
 Phlox pilosa - PL, R
 Phlox setacea - PL
 Polypodium virginianum - PE,
 R
 Rhexia mariana - PL
 Ribes oxycanthoides - PL
 Smilax bona-nox - PL
 Solidago caesia - R
 Trifolium comosum - PE

For Linnaeus, the botanical explorations that occurred in temperate North America prior to 1753 were significant to him. He never saw the New World himself and had to rely upon the efforts of others to provide him with information. This came in the form of dried herbarium specimens, seeds and fruits that resulted in culti-

vated plants he could examine, and publications he could read and study. Many of the herbarium specimens seen by Linnaeus were gathered by John Clayton and Peter Kalm. Of the 889 species and varieties described by Linnaeus in the first edition of *Species plantarum* in 1753, at least 586 entities, or 66%, were based on their collections. This includes 551 species for which there are Clayton specimens, 205 species for which there are Kalm specimens, and 170 species which were gathered by both men. Of the remaining 303 species, Linnaeus has material (mostly garden specimens or collections which cannot be determined as to collector) for some 250 of those species. The following species do not seem to have obvious specimens immediately available to Linnaeus. A star (*) indicates those species which are not illustrated.

Annona glabra
 Arnica maritima*
 Asclepias amoena
 Asclepias variegata
 Aster miser
 Aster mutabilis
 Astragalus carolinianus
 Bignonia caerulea
 Bignonia sempervirens
 Carduus altissimus
 Cissampelos smilacina
 Clinopodium rugosum
 Clitoria mariana*
 Convallaria stellata
 Convolvulus carolinus
 Conyza bifrons
 var. flosculosa
 Conyza linifolia
 Eupatorium hyssopifolium
 Euphorbia maculata
 Gomphrena interrupta*
 Gomphrena serrata*
 Guilandina dioica*
 Helleborus scaber*
 Horminum virginicum
 Hypericum lasianthus
 Ipomoea carolina
 Ipomoea lacunosa
 Ipomoea tamnifolia
 Iris versicolor

*Lilium camschatcense**
Lycopodium carolinianum
Magnolia virginiana
 var. *grisea**
Magnolia virginiana
 var. *tripetala*
*Othonna cinneraria**
Pancratium carolinianum
Phaseolus helvulus
Philadelphus inodorus
Phlox pilosa
Phlox setacea
Pistacia simaruba
Polemonium rubrum
Polypodium virginianum
Prenanthes altissima
Rhexia mariana
Ribes oxyacanthoides
Sloanea emarginata
Smilax bona-nox
Solanum nigrum
 var. *virginicum*
Solidago caesia
Thalictrum cornutii
*Trifolium comosum**
Triosteum perfoliatum
Viscum rubrum

CONCLUSIONS

Linnaeus left an impressive record of accomplishments when he died in 1778. His impact has extended far beyond his own life time, and today, the works of Carl Linnaeus are as critical to botanists now as they were at the time they were published. In the present review of just one publication -- the first edition of *Species plantarum* -- it can be seen that not only is this publication itself important, but the basis upon which it was established reflects Linnaeus' premier position within the botanical community. This becomes obvious when one realizes that Linnaeus had access, directly through specimens and indirectly through the literature, to a large percentage of

the world's flora in 1753. No doubt Linnaeus was a man of tremendous industry. He had the time and opportunity to work on projects he deemed important. He had the support of a large number of his colleagues, although many did not accept his general views as expressed in his sexual system of classification. Linnaeus revolutionized systematic botany by his consistent use of binomials. Not only did Linnaeus make it simpler to index names -- as was his intention -- he also made it more convenient for people of lesser intellect to remember scientific names.

What can be seen from this review of the temperate North American flora as reported in 1753 by Linnaeus is that he depended upon a great number of people to provide him with descriptions, illustrations, seeds and specimens. His place in the history of science is well established and shall not be diminished with time. That position does rest, however, upon the energies and even the lives of a host of men and women willing to face the unknown to discover the unseen so that he might inform the world of the existence of new species of plants. Time has not changed that fact. Today, we too fully depend upon the efforts of the many past generations of naturalists for our understanding of the living world. It is hoped that this review will remind the modern plant taxonomist that the history of systematic botany began long before 1753 and the first edition of *Species plantarum*.

New York Botanical Garden Library



3 5185 00288 2718

Inasmuch as we do no editing, papers accepted for publication *must* be submitted in *exactly* the form that the author wants to have them published. They will then be photographed and printed by photo-offset in exactly the form as submitted except that we will add page numbers and running-heads.

Typescripts should be prepared single-spaced on clean white heavy bond smooth and opaque paper. Elite type is probably the most space-economical. Typescript text must not exceed a rectangle $5\frac{1}{2}$ inches wide (horizontal) by $8\frac{1}{2}$ inches high (vertical), not including the running-head and page number.

The title of the paper should be typed in all uppercase (capital) letters with 2 blank lines above the title and one beneath; then the name of the author in ordinary upper-and lower-case letters, along with his address (if so desired); followed by 2 blank lines; then the first line of text. It is usually best to leave a blank line between paragraphs.

All scientific plant and animal names *and* group names should be typed either in italic type (if available) or underscored. Any corrections in the text made by the author must be complete and neat as they will be photographed as they are.

The finished typescript as submitted by the author will be reduced from the $8\frac{1}{2}$ x $5\frac{1}{2}$ inch size as submitted to $6\frac{2}{8}$ x 4 inches by the printer. It is therefore advisable to place a centimeter or millimeter scale on all text figures and plates included.

Use a *new* heavily inked black typewriter ribbon and be sure to *clean* the type on the typewriter after each several pages of typing.

Cost of publication at present is \$10.00 US per page, with no subsequent rebates, but this rate may vary depending on inflation and costs, so it is best to inquire as to current rates. The page charges are due *with* the typescript and no paper will be published before payment is received in full. Each author will receive gratis a proportionate share of the printed copies remaining after paid subscriptions are filled, but if separates (reprints or offprints) are desired, these will be charged extra in accord with the current rate for offprints provided by the printer. The cost of all such separates ordered must also be paid for in advance at the time the typescript is sent. No orders for separates will be accepted later, nor can additions or corrections be accepted.

Authors are asked to indicate in light pencil on the *reverse* side of each page of their typescript the page number so that no mistakes in sequence occur.

Each number consists of not less than 32 pages. All manuscript accepted will be published in the next issue, so that the size of numbers may vary greatly. A volume will contain 512 pages. This plan insures prompt publication of all accepted manuscript.

Illustrations will be published according to the desires of the authors. No extra charge is made for line drawings, such as are ordinarily reproduced in zinc, or for diagrams, tables, or charts, provided they conform to certain limitations of size and proportion. An extra charge will be made for halftones, depending on their size, as fixed by the engraver.

Articles dealing with research in all lines of botany and plant ecology, in any reasonable length, biographical sketches, and critical reviews and summaries of literature will be considered for publication.